


Math 3322 Test II  
DeMaio Spring 2010

Name \_\_\_\_\_


**Instructions.** Show all your work. Credit cannot and will not be awarded for work not shown. **Where appropriate, simplify all answers to a single decimal expansion.**

- (30 points) Recall our 20 person club with 13 women and 7 men. How many ways can we
  - elect a President and Vice-President;  $20 * 19 = 380$
  - form a committee of three people (where each member has equal rank and power);  $\binom{20}{3} = 1140$
  - form a committee of four people and elect one of those four members to be the chairperson;  $\binom{20}{4} * 4 = 19380$
  - form a committee of two people of opposite gender;  $13 * 7 = 91$
  - form a committee of two people of the same gender;  $\binom{13}{2} + \binom{7}{2} = 99$  or  $\binom{20}{2} - 91 = 99$
  - form a committee of six people with at least one member of each gender?  $\binom{20}{6} - \binom{13}{6} - \binom{7}{6} = 37037$
- (15 points) How many ways can we rearrange the letters in the word
  - vampire;  $7! = 5040$
  - werewolf;  $\frac{8!}{2! * 2!} = 10080$
  - mummy?  $\frac{5!}{3!} = 20$
- (10 points) Let  $D = \{1, 2, 3, 4, 5\}$  and  $R = \{a, b, c, d, e, f, g, h\}$ .  
How many functions  $f : D \rightarrow R$  exist?  $8^5 = 32768$   
How many one-to-one functions  $f : D \rightarrow R$  exist?  $8 * 7 * 6 * 5 * 4 = \frac{8!}{3!} = 6720$
- (10 points) Use the Binomial Theorem to expand  $(3x - 2)^5$ . You must show all the details of your work.  
 $(3x - 2)^5 =$   
 $\binom{5}{5} (3x)^5 (-2)^0 + \binom{5}{4} (3x)^4 (-2)^1 + \binom{5}{3} (3x)^3 (-2)^2 + \binom{5}{2} (3x)^2 (-2)^3 + \binom{5}{1} (3x)^1 (-2)^4 + \binom{5}{0} (3x)^0 (-2)^5$   
 $= 243x^5 - 5 * 81 * 2x^4 + 10 * 27 * 4x^3 - 10 * 9 * 8x^2 + 5 * 3 * 16x - 32$   
 $= 243x^5 - 810x^4 + 1080x^3 - 720x^2 + 240x - 32$
- (10 points) John has 13 ordinary coins (cent, nickel, dime, quarter) in his pocket.  
At least how many of the same coin must John have?  $\lceil \frac{13}{4} \rceil = 4$   
At least how many quarters must John have? None.
- (10 points) What is the minimum number of students, each of whom comes from one of the 50 states, who must be enrolled in a university to guarantee that there are at least 100 who come from the same state?  $50 * 99 + 1 = 4951$ .
- (15 points) Prove  $\sum_{i=1}^n F_{2i-1} = F_{2n}$  for  $n \geq 1$  for the Fibonacci sequence,  $F_n$ . First show that  $S(1)$  is true. So,  $\sum_{i=1}^1 F_{2i-1} = F_{2*1-1} = F_1 = 1$ . Since,  $F_{2*1} = F_2 = 1$ ,  $S(1)$  is true. Second, assume  $\sum_{i=1}^n F_{2i-1} = F_{2n}$  and show  $\sum_{i=1}^{n+1} F_{2i-1} = F_{2(n+1)} = F_{2n+2}$ .  
 $\sum_{i=1}^{n+1} F_{2i-1} = \sum_{i=1}^n F_{2i-1} + F_{2(n+1)-1} = \sum_{i=1}^n F_{2i-1} + F_{2n+1}$ . By the inductive assumption,  $\sum_{i=1}^n F_{2i-1} + F_{2n+1} = F_{2n} + F_{2n+1}$ . Of course, by the definition of the Fibonacci sequence,  $F_{2n} + F_{2n+1} = F_{2n+2}$ .


8. (15 points) Chairs in a Row: the Couples version: In this version, we have a row of  $n$  chairs. People arrive in twos and want to sit next to their partner. For instance, a row of 4 chairs can be filled in the following 5 ways:



Find all the ways a row of 2 chairs can be filled with couples or be left empty?



Find all the ways a row of 3 chairs can be filled with couples or be left empty?



Conjecture and prove a formula for the number of ways a row of  $n$  chairs can be filled with couples or be left empty? Let  $S_n$  be the number of ways a row of  $n$  chairs can be filled with couples or be left empty. Given that  $S_2 = 2$ ,  $S_3 = 3$  and  $S_4 = 5$  it would appear that  $S_n = F_{n+1}$ . So, is  $S_n = S_{n-1} + S_{n-2}$ ? Partition all  $S_n$  rows into two sets, those that end in an empty chair and those that end in a person. There are  $S_{n-1}$  that end in an empty chair because you can add an empty chair on the end of every row of length  $n - 1$ . A row that ends in a person, actually ends in two people since couples sit together. This means there are  $S_{n-2}$  rows that end in a person since you can add a couple to the end of every row of length  $n - 2$ . Thus,  $S_n = S_{n-1} + S_{n-2}$  and  $S_n = F_{n+1}$ .