

April 17 Math 2306 sec. 53 Spring 2019

Section 17: Fourier Series: Trigonometric Series

Consider the following problem:

An undamped spring mass system has a mass of 2 kg attached to a spring with spring constant 128 N/m. The mass is driven by an external force $f(t) = 2t$ for $-1 < t < 1$ that is 2-periodic so that $f(t + 2) = f(t)$ for all $t > 0$.




Figure:
$$2 \frac{d^2 x}{dt^2} + 128x = f(t)$$

Common Models of Periodic Sources (e.g. Voltage)




Figure: We'd like to solve, or at least approximate solutions, to ODEs and PDEs with periodic *right hand sides*.

Series Representations for Functions

The goal is to represent a function by a series

$$f(x) = \sum_{n=1}^{\infty} (\text{some simple functions})$$

In calculus, you saw power series $f(x) = \sum_{n=0}^{\infty} a_n(x - c)^n$ where the simple functions were powers $(x - c)^n$.

Here, you will see how some functions can be written as series of trigonometric functions

$$f(x) = \sum_{n=0}^{\infty} (a_n \cos nx + b_n \sin nx)$$

We'll move the $n = 0$ to the front before the rest of the sum.

Some Preliminary Concepts

Suppose two functions f and g are integrable on the interval $[a, b]$. We define the **inner product** of f and g on $[a, b]$ as

$$\langle f, g \rangle = \int_a^b f(x)g(x) dx.$$

$\langle f, g \rangle$ is a number

We say that f and g are **orthogonal** on $[a, b]$ if

$$\langle f, g \rangle = 0.$$

The product depends on the interval, so the orthogonality of two functions depends on the interval.

Properties of an Inner Product

Let f , g , and h be integrable functions on the appropriate interval and let c be any real number. The following hold

$$(i) \quad \langle f, g \rangle = \langle g, f \rangle$$

$$(ii) \quad \langle f, g + h \rangle = \langle f, g \rangle + \langle f, h \rangle$$

$$(iii) \quad \langle cf, g \rangle = c \langle f, g \rangle$$

$$(iv) \quad \langle f, f \rangle \geq 0 \text{ and } \langle f, f \rangle = 0 \text{ if and only if } f = 0$$

Orthogonal Set

A set of functions $\{\phi_0(x), \phi_1(x), \phi_2(x), \dots\}$ is said to be **orthogonal** on an interval $[a, b]$ if

$$\langle \phi_m, \phi_n \rangle = \int_a^b \phi_m(x) \phi_n(x) dx = 0 \quad \text{whenever} \quad m \neq n.$$

Note that any function $\phi(x)$ that is not identically zero will satisfy

$$\langle \phi, \phi \rangle = \int_a^b \phi^2(x) dx > 0.$$

Hence we define the **square norm** of ϕ (on $[a, b]$) to be

$$\|\phi\| = \sqrt{\int_a^b \phi^2(x) dx}.$$

An Orthogonal Set of Functions

Consider the set of functions

$$\{1, \cos x, \cos 2x, \cos 3x, \dots, \sin x, \sin 2x, \sin 3x, \dots\} \quad \text{on } [-\pi, \pi].$$

Evaluate $\langle \cos(nx), 1 \rangle$ and $\langle \sin(mx), 1 \rangle$. here n, m are any positive integer.

By definition

$$\langle \cos(nx), 1 \rangle = \int_{-\pi}^{\pi} \cos(nx) \cdot 1 \, dx$$

$$\begin{aligned} &= \frac{1}{n} \sin(nx) \Big|_{-\pi}^{\pi} = \frac{1}{n} \sin(n\pi) - \frac{1}{n} \sin(-n\pi) \\ &= 0 - 0 = 0 \end{aligned}$$

$$\langle \cos(nx), 1 \rangle = 0$$

$\cos(nx)$ is orthogonal to 1 for all n .

Now compute $\langle \sin(mx), 1 \rangle$

$$\begin{aligned}\langle \sin(mx), 1 \rangle &= \int_{-\pi}^{\pi} \sin(mx) \cdot 1 \, dx \\ &= \frac{-1}{m} \cos(mx) \Big|_{-\pi}^{\pi} = \frac{-1}{m} \cos(m\pi) - \frac{-1}{m} \cos(-m\pi)\end{aligned}$$

$\cos(-\theta) = \cos \theta$ by even symmetry $= 0$

$\langle \sin(mx), 1 \rangle = 0$ for all $m = 1, 2, 3, \dots$

so $\sin(mx)$ is orthogonal to 1 for all m
on $[-\pi, \pi]$.

An Orthogonal Set of Functions

Consider the set of functions

$$\{1, \cos x, \cos 2x, \cos 3x, \dots, \sin x, \sin 2x, \sin 3x, \dots\} \quad \text{on} \quad [-\pi, \pi].$$

It can easily be verified that

$$\int_{-\pi}^{\pi} \cos nx \, dx = 0 \quad \text{and} \quad \int_{-\pi}^{\pi} \sin mx \, dx = 0 \quad \text{for all} \quad n, m \geq 1,$$

$$\int_{-\pi}^{\pi} \cos nx \sin mx \, dx = 0 \quad \text{for all} \quad m, n \geq 1, \quad \text{and}$$

$$\int_{-\pi}^{\pi} \cos nx \cos mx \, dx = \int_{-\pi}^{\pi} \sin nx \sin mx \, dx = \begin{cases} 0, & m \neq n \\ \pi, & n = m \end{cases},$$

An Orthogonal Set of Functions on $[-\pi, \pi]$

These integral values indicated that the set of functions

$$\{1, \cos x, \cos 2x, \cos 3x, \dots, \sin x, \sin 2x, \sin 3x, \dots\}$$

is an orthogonal set on the interval $[-\pi, \pi]$.

Key Point: This means that if we take any two functions f and g from **this set**, then

$$\int_{-\pi}^{\pi} f(x)g(x) dx = 0 \quad \text{if } f \text{ and } g \text{ are different functions!}$$

Fourier Series

Suppose $f(x)$ is defined for $-\pi < x < \pi$. We would like to know how to write f as a series **in terms of sines and cosines**.

Task: Find coefficients (numbers) a_0, a_1, a_2, \dots and b_1, b_2, \dots such that¹

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx).$$

¹We'll write $\frac{a_0}{2}$ as opposed to a_0 purely for convenience.

Fourier Series

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx).$$

The question of convergence naturally arises when we wish to work with infinite series. To highlight convergence considerations, some authors prefer not to use the equal sign when expressing a Fourier series and instead write

$$f(x) \sim \frac{a_0}{2} + \dots$$

Herein, we'll use the equal sign with the understanding that equality may not hold at each point.

Convergence will be address later.

Finding an Example Coefficient

Let's find the coefficient b_4 .

Start with the series $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$, and multiply both sides by $\sin(4x)$.

$$f(x) \sin(4x) = \frac{a_0}{2} \sin(4x) + \sum_{n=1}^{\infty} (a_n \cos nx \sin(4x) + b_n \sin nx \sin(4x)).$$

We'll integrate both sides from $-\pi$ to π , we'll assume that it's safe to switch the order of integration and summation.

$$\int_{-\pi}^{\pi} f(x) \sin(4x) dx = \int_{-\pi}^{\pi} \frac{a_0}{2} \sin(4x) dx + \sum_{n=1}^{\infty} \left[\int_{-\pi}^{\pi} a_n \cos(nx) \sin(4x) dx \right]$$

$$+ \int_{-\pi}^{\pi} b_n \sin(nx) \sin(4x) dx$$

$$\int_{-\pi}^{\pi} f(x) \sin(4x) dx = \frac{a_0}{2} \int_{-\pi}^{\pi} \overset{=0}{\sin(4x)} dx +$$

$$\sum_{n=1}^{\infty} \left[a_n \int_{-\pi}^{\pi} \overset{=0}{\cos(nx) \sin(4x)} dx + b_n \int_{-\pi}^{\pi} \sin(nx) \sin(4x) dx \right]$$

Recall $\int_{-\pi}^{\pi} \sin(mx) dx = 0$ and $\int_{-\pi}^{\pi} \cos(mx) \sin(mx) dx = 0$

for all n, m

So we have

$$\int_{-\pi}^{\pi} f(x) \sin(4x) dx = \sum_{n=1}^{\infty} b_n \int_{-\pi}^{\pi} \sin(nx) \sin(4x) dx$$

$$\int_{-\pi}^{\pi} \sin(nx) \sin(4x) dx = \begin{cases} 0, & n \neq 4 \\ \pi, & n = 4 \end{cases}$$

$$\begin{aligned} \int_{-\pi}^{\pi} f(x) \sin(4x) dx &= 0 + 0 + 0 + b_4 \pi + 0 + \dots \\ &= b_4 \pi \end{aligned}$$

Finally,

$$b_4 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(4x) dx$$

Finding Fourier Coefficients

Note that there was nothing special about seeking the 4th sine coefficient b_4 . We could have just as easily sought b_m for any positive integer m . We would simply start by introducing the factor $\sin(mx)$.

Moreover, using the same orthogonality property, we could pick on the a 's by starting with the factor $\cos(mx)$ —including the constant term since $\cos(0 \cdot x) = 1$. The only minor difference we want to be aware of is that

$$\int_{-\pi}^{\pi} \cos^2(mx) dx = \begin{cases} 2\pi, & m = 0 \\ \pi, & m \geq 1 \end{cases}$$

Careful consideration of this sheds light on why it is conventional to take the constant to be $\frac{a_0}{2}$ as opposed to just a_0 .

The Fourier Series of $f(x)$ on $(-\pi, \pi)$

The **Fourier series** of the function f defined on $(-\pi, \pi)$ is given by

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx).$$

Where

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx,$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx, \quad \text{and}$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$