

## Section 1.1: Systems of Linear Equations

We begin with a linear (*algebraic*) equation in  $n$  variables  $x_1, x_2, \dots, x_n$  for some positive integer  $n$ .

A **linear equation** can be written in the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = b.$$

The numbers  $a_1, \dots, a_n$  are called the *coefficients*. These numbers and the right side  $b$  are real (or complex) constants that are **known**.

# Examples of Equations that are or are not Linear

Both  
linear

$$2x_1 = 4x_2 - 3x_3 + 5 \quad \text{and} \quad 12 - \sqrt{3}(x + y) = 0$$

$$2x_1 - 4x_2 + 3x_3 = 5$$

$$\sqrt{3}x + \sqrt{3}y = 12$$

Both  
non linear

$$x_1 + 3x_3 = \frac{1}{x_2} \quad \text{and} \quad xyz = \sqrt{w}$$

non  
linear  
term

non linear  
product of variables  
square root of variables

A *Linear System* is a collection of linear equations in the same variables

$$2x_1 + x_2 - 3x_3 + x_4 = -3$$

$$-x_1 + 3x_2 + 4x_3 - 2x_4 = 8$$

$$x + 2y + 3z = 4$$

$$3x + 12z = 0$$

$$2x + 2y - 5z = -6$$

# Some terms

- ▶ A **solution** is a list of numbers  $(s_1, s_2, \dots, s_n)$  that reduce each equation in the system to a true statement upon substitution.
- ▶ A **solutions set** is the set of all possible solutions of a linear system.
- ▶ Two systems are called **equivalent** if they have the same solution set.

# An Example

$$\begin{array}{rcl} 2x & - & y = -1 \\ -4x & + & 2y = 2 \end{array}$$

(a) Show that  $(1, 3)$  is a solution.

$x=1$  and  $y=3$  in the pair  $(1, 3)$ .

In equation 1 setting  $x=1$ ,  $y=3$

$$2(1) - 3 = 2 - 3 = -1 \Rightarrow -1 = -1 \quad \text{this is true}$$

In equation 2

$$-4(1) + 2(3) = -4 + 6 = 2 \Rightarrow 2 = 2$$

Both equations are satisfied  $\Rightarrow (1, 3)$  is a solution.

## An Example Continued

$$\begin{array}{rclcrcl} 2x & - & y & = & -1 \\ -4x & + & 2y & = & 2 \end{array}$$


(b) Note that  $\{(x, y) | y = 2x + 1\}$  is the solution set.

$$\text{From } 2x - y = -1 \Rightarrow 2x = y - 1 \Rightarrow 2x + 1 = y$$

$$\begin{aligned} \text{From } -4x + 2y &= 2 \Rightarrow 2y = 2 + 4x \\ 2y &= 2(1 + 2x) \\ y &= 1 + 2x \end{aligned}$$


The set condition is the same as each equation in the system.

# The Geometry of 2 Equations with 2 Variables



**Figure:** The system  $x - y = -1$  and  $2x + y = 3$  with solution set  $\{(2/3, 5/2)\}$ . These equations represent lines that intersect at one point.


# The Geometry of 2 Equations with 2 Variables



**Figure:** The system  $x - y = -1$  and  $2x - 2y = -2$  with solution set  $\{(x, y) | y = x + 1\}$ . Both equations represent the same line which share all common points as solutions.



# The Geometry of 2 Equations with 2 Variables



**Figure:** The system  $x - y = -1$  and  $2x - 2y = 2$  with solution set  $\emptyset$ . These equations represent parallel lines having no common points.

# Theorem

A linear system of equations has exactly one of the following:

- i No solution, or
- ii Exactly one solution, or
- iii Infinitely many solutions.

Terms: A system is

**consistent** if it has at least one solution (cases ii and iii), and  
**inconsistent** if it has no solutions (case i).

Two critical questions about any linear system are: (1) Does it have a solution? (existence), and (2) If it has a solution, is there only one? (uniqueness)

# Matrices

**Definition:** A matrix is a rectangular array of numbers. It's **size** (a.k.a. dimension/order) is  $m \times n$  (read " $m$  by  $n$ ") where  $m$  is the number of rows and  $n$  is the number of columns the matrix has.

Examples:

$$\begin{bmatrix} 2 & 0 & -1 & 3 \\ 1 & 1 & 13 & -4 \\ 12 & -3 & 2 & -2 \end{bmatrix},$$

$3 \times 4$

$$\begin{bmatrix} 2 & 0 \\ 4 & 4 \\ 3 & -5 \end{bmatrix}$$

$3 \times 2$

## Linear System: Coefficient Matrix

Given any linear system of equations, we can associate two matrices with the system. These are the **coefficient** matrix and the **augmented** matrix<sup>1</sup>.

Example:

$$\begin{array}{rrcrcl} x_1 & + & 2x_2 & - & x_3 & = & -4 \\ 2x_1 & & & + & x_3 & = & 7 \\ x_1 & + & x_2 & + & x_3 & = & 6 \end{array}$$

$$\begin{bmatrix} 1 & 2 & -1 \\ 2 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

The coefficient matrix contains the coefficients of the system.

$m = \# \text{ equations}$

$n = \# \text{ variables}$

---

<sup>1</sup>Note that like variables should be lined up vertically!

## Linear System: Augmented Matrix

Given any linear system of equations, we can associate two matrices with the system. These are the **coefficient** matrix and the **augmented** matrix.

Example:

$$\begin{array}{rrcrcl} x_1 & + & 2x_2 & - & x_3 & = & -4 \\ 2x_1 & & & + & x_3 & = & 7 \\ x_1 & + & x_2 & + & x_3 & = & 6 \end{array}$$

$$\left[ \begin{array}{cccc} 1 & 2 & -1 & -4 \\ 2 & 0 & 1 & 7 \\ 1 & 1 & 1 & 6 \end{array} \right]$$

The augmented matrix is the coefficient matrix with an extra column for the right side.

$m = \# \text{ equations}$

$n = \# \text{ variables} + 1$

# Legitimate Operations for Solving a System

We can perform three basic operation **without** changing the solution set of a system. These are

- ▶ swap the order of any two equations (**swap**),
- ▶ multiply an equation by any nonzero constant (**scale**), and
- ▶ replace an equation with the sum of itself and a nonzero multiple of any other equation (**replace**).

# Some Operation Notation

## Notation

- ▶ Swap equations  $i$  and  $j$ :

$$E_i \leftrightarrow E_j$$

- ▶ Scale equation  $i$  by  $k$ :

$$kE_i \rightarrow E_i$$

- ▶ Replace equation  $j$  with the sum of itself and  $k$  times equation  $i$ :

$$kE_i + E_j \rightarrow E_j$$

Solve the following system of equations by *elimination*. Keep tabs on the augmented matrix at each step.

$$\begin{array}{rrcr} x_1 & + & 2x_2 & - & x_3 & = & -4 \\ 2x_1 & & & + & x_3 & = & 7 \\ x_1 & + & x_2 & + & x_3 & = & 6 \end{array}$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & -1 & -4 \\ 2 & 0 & 1 & 7 \\ 1 & 1 & 1 & 6 \end{array} \right]$$

$$-2E_1 + E_2 \rightarrow E_2$$

$$-E_1 + E_3 \rightarrow E_3$$

$$\begin{array}{rrcr} x_1 & + & 2x_2 & - & x_3 & = & -4 \\ & & -4x_2 & + & 3x_3 & = & 15 \\ & & -x_2 & + & 2x_3 & = & 10 \end{array}$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & -1 & -4 \\ 0 & -4 & 3 & 15 \\ 0 & -1 & 2 & 10 \end{array} \right]$$

$$E_2 \leftrightarrow E_3$$

$$\text{then } -E_2 \rightarrow E_2$$



$$x_1 + 2x_2 - x_3 = -4$$

$$x_2 - 2x_3 = -10$$

$$-4x_2 + 3x_3 = 15$$

$$\begin{bmatrix} 1 & 2 & -1 & -4 \\ 0 & 1 & -2 & -10 \\ 0 & -4 & 3 & 15 \end{bmatrix}$$

$$4E_2 + E_3 \rightarrow E_3$$

$$x_1 + 2x_2 - x_3 = -4$$

$$x_2 - 2x_3 = -10$$

$$-5x_3 = -25$$

$$\begin{bmatrix} 1 & 2 & -1 & -4 \\ 0 & 1 & -2 & -10 \\ 0 & 0 & -5 & -25 \end{bmatrix}$$

$$-\frac{1}{5}E_3 \rightarrow E_3$$

$$x_1 + 2x_2 - x_3 = -4$$

$$x_2 - 2x_3 = -10$$

$$x_3 = 5$$

$$\begin{bmatrix} 1 & 2 & -1 & -4 \\ 0 & 1 & -2 & -10 \\ 0 & 0 & 1 & 5 \end{bmatrix}$$

From E2  $x_2 = -10 + 2x_3$  but  $x_3 = 5$   
 $= -10 + 10 = 0$

From E1  $x_1 = -4 - 2x_2 + x_3 = -4 - 0 + 5 = 1$

The solution can be expressed as  
a triple  $(1, 0, 5)$ .

# Elementary Row Operations

If any sequence of the following operations are performed on a matrix, the resulting matrix is **row equivalent**.

- i Replace a row with the sum of itself and a multiple of another row (**replacement**).
- ii Interchange any two rows (**row swap**).
- iii Multiply a row by any nonzero constant (**scaling**).

**Theorem:** If the augmented matrices of two linear systems are row equivalent, then the systems have the same solution set. (i.e. The systems are equivalent!)

## A key here is *structure*!

Consider the following augmented matrix. Determine if the associated system is consistent or inconsistent. If it is consistent, determine the solution set.

$$(a) \quad \left[ \begin{array}{cccc} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

$$\begin{aligned} x_1 &= 3 \\ x_2 &= 1 \\ x_3 &= -2 \end{aligned}$$

This is consistent with solution  $\begin{aligned} x_1 &= 3 \\ x_2 &= 1 \\ x_3 &= -2 \end{aligned}$

(b)

$$\begin{bmatrix} 1 & 2 & 0 & 3 \\ 0 & 1 & -1 & 4 \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

$$x_1 + 2x_2 = 3$$

$$x_2 - x_3 = 4$$

$$0 = 3 \leftarrow \text{always false}$$

The system is inconsistent.

$$(c) \begin{bmatrix} 1 & 1 & -1 & 0 \\ 0 & 1 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$x_1 + x_2 - x_3 = 0$$

$$x_2 + x_3 = 4$$

$$0 = 0$$

$$\text{From } E_2 \quad x_2 = 4 - x_3$$

$$\text{From } E_1 \quad x_1 = -x_2 + x_3 = -(4 - x_3) + x_3 = -4 + 2x_3$$

We can express this as

$$x_1 = -4 + 2x_3$$

$$x_2 = 4 - x_3$$

$x_3$  is free

This could be expressed  $x_3 \in \mathbb{R}$

Here  $x_1$  and  $x_2$  are called basic variables.