

Section 1.1: Definitions and Terminology

Suppose $y = \phi(x)$ is a differentiable function. We know that $dy/dx = \phi'(x)$ is another (related) function.

For example, if $y = \cos(2x)$, then y is differentiable on $(-\infty, \infty)$. In fact,

$$\frac{dy}{dx} = -2 \sin(2x).$$

Even dy/dx is differentiable with $d^2y/dx^2 = -4 \cos(2x)$. Note that

$$\frac{d^2y}{dx^2} + 4y = 0.$$

$$-4 \cos(2x) + 4 \cos(2x) = 0 \quad \checkmark$$

Differential Equation

The equation

$$\frac{d^2y}{dx^2} + 4y = 0.$$

is an example of a **differential equation**.

Questions: If we only started with the equation, how could we determine that $\cos(2x)$ satisfies it? Also, is $\cos(2x)$ the only possible function that y could be?

Definition

A **Differential Equation** is an equation containing the derivative(s) of one or more dependent variables, with respect to one or more independent variables.

Independent Variable: will appear as one that derivatives are taken **with respect to**.

Dependent Variable: will appear as one that derivatives are taken **of**.

$$\frac{dy}{dx}$$

Handwritten notes: $y \leftarrow$ depend., $x \uparrow$ independ.

$$\frac{du}{dt}$$

Handwritten notes: $u \leftarrow$ depend., $t \uparrow$ indep.

$$\frac{dx}{dr}$$

Handwritten notes: $x \leftarrow$ dep, $r \uparrow$ indep.

Classifications

Type: An **ordinary differential equation (ODE)** has exactly one independent variable¹. For example

$$\frac{dy}{dx} - y^2 = 3x, \quad \text{or} \quad \frac{dy}{dt} + 2\frac{dx}{dt} = t, \quad \text{or} \quad y'' + 4y = 0$$

A **partial differential equation (PDE)** has two or more independent variables. For example

$$\frac{\partial y}{\partial t} = \frac{\partial^2 y}{\partial x^2}, \quad \text{or} \quad \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$$

¹These are the subject of this course.

Classifications

Order: The order of a differential equation is the same as the highest order derivative appearing anywhere in the equation.

$$\frac{dy}{dx} - y^2 = 3x \quad 1^{\text{st}} \text{ order}$$

$$y''' + (y')^4 = x^3 \quad 3^{\text{rd}} \text{ order}$$

$$\frac{\partial y}{\partial t} = \frac{\partial^2 y}{\partial x^2} \quad 2^{\text{nd}} \text{ order}$$

Notations and Symbols

We'll use standard derivative notations:

Leibniz: $\frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^ny}{dx^n},$ or

Prime & superscripts: $y', y'', \dots y^{(n)}.$

Newton's **dot notation** may be used if the independent variable is time. For example if s is a position function, then

velocity is $\frac{ds}{dt} = \dot{s},$ and acceleration is $\frac{d^2s}{dt^2} = \ddot{s}$

Notations and Symbols

An n^{th} order ODE, with independent variable x and dependent variable y can always be expressed as an equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

where F is some real valued function of $n + 2$ variables.

Normal Form: If it is possible to isolate the highest derivative term, then we can write a **normal form** of the equation

$$\frac{d^n y}{dx^n} = f(x, y, y', \dots, y^{(n-1)}).$$

Ex: $\frac{dy}{dx} - y^2 = 3x$

$$\Rightarrow \underbrace{\frac{dy}{dx} - y^2 - 3x}_{F(x, y, y')} = 0$$

In normal form $\frac{dy}{dx} = \underbrace{y^2 + 3x}_{f(x, y)}$

Notations and Symbols

If $n = 1$ or $n = 2$, an equation in normal form would look like

$$\frac{dy}{dx} = f(x, y) \quad \text{or} \quad \frac{d^2y}{dx^2} = f(x, y, y').$$

Differential Form: A first order equation may appear in the form

$$M(x, y) dx + N(x, y) dy = 0$$