

Section 6.4 Trigonometric Substitution

AP_EX **CALCULUS II**
Version 2.0

Authors

Gregory Hartman, Ph.D.

*Department of Applied Mathematics
Virginia Military Institute*

Brian Heinold, Ph.D.

*Department of Mathematics and Computer Science
Mount Saint Mary's University*

Troy Siemers, Ph.D.

*Department of Applied Mathematics
Virginia Military Institute*


Dimplekumar Chalishajar, Ph.D.

*Department of Applied Mathematics
Virginia Military Institute*

Editor

Jennifer Bowen, Ph.D.

*Department of Mathematics and Computer Science
The College of Wooster*



Copyright © 2014 Gregory Hartman
Licensed to the public under Creative Commons
Attribution-Noncommercial 3.0 United States License

Contents

Preface	iii
Table of Contents	v
5 Integration	185
5.1 Antiderivatives and Indefinite Integration	185
5.2 The Definite Integral	194
5.3 Riemann Sums	204
5.4 The Fundamental Theorem of Calculus	221
5.5 Numerical Integration	233
6 Techniques of Antidifferentiation	247
6.1 Substitution	247
6.2 Integration by Parts	266
6.3 Trigonometric Integrals	276
6.4 Trigonometric Substitution	286
6.5 Partial Fraction Decomposition	295
6.6 Hyperbolic Functions	303
6.7 L'Hôpital's Rule	313
6.8 Improper Integration	321
7 Applications of Integration	333
7.1 Area Between Curves	334
7.2 Volume by Cross-Sectional Area; Disk and Washer Methods	341
7.3 The Shell Method	348
7.4 Arc Length and Surface Area	356
7.5 Work	365
7.6 Fluid Forces	375
8 Sequences and Series	383
8.1 Sequences	383
8.2 Infinite Series	395
8.3 Integral and Comparison Tests	410
8.4 Ratio and Root Tests	419

8.5	Alternating Series and Absolute Convergence	424
8.6	Power Series	434
8.7	Taylor Polynomials	446
8.8	Taylor Series	457

A Solutions To Selected Problems **A.1**

Index **A.11**

6.4 Trigonometric Substitution

In Section 5.2 we defined the definite integral as the “signed area under the curve.” In that section we had not yet learned the Fundamental Theorem of Calculus, so we evaluated special definite integrals which described nice, geometric shapes. For instance, we were able to evaluate

$$\int_{-3}^3 \sqrt{9 - x^2} dx = \frac{9\pi}{2} \quad (6.1)$$

as we recognized that $f(x) = \sqrt{9 - x^2}$ described the upper half of a circle with radius 3.

We have since learned a number of integration techniques, including Substitution and Integration by Parts, yet we are still unable to evaluate the above integral without resorting to a geometric interpretation. This section introduces Trigonometric Substitution, a method of integration that fills this gap in our integration skill. This technique works on the same principle as Substitution as found in Section 6.1, though it can feel “backward.” In Section 6.1, we set $u = f(x)$, for some function f , and replaced $f(x)$ with u . In this section, we will set $x = f(\theta)$, where f is a trigonometric function, then replace x with $f(\theta)$.

We start by demonstrating this method in evaluating the integral in (6.1). After the example, we will generalize the method and give more examples.

Example 172 Using Trigonometric Substitution

Evaluate $\int_{-3}^3 \sqrt{9 - x^2} dx$.

SOLUTION We begin by noting that $9 \sin^2 \theta + 9 \cos^2 \theta = 9$, and hence $9 \cos^2 \theta = 9 - 9 \sin^2 \theta$. If we let $x = 3 \sin \theta$, then $9 - x^2 = 9 - 9 \sin^2 \theta = 9 \cos^2 \theta$.

Setting $x = 3 \sin \theta$ gives $dx = 3 \cos \theta d\theta$. We are almost ready to substitute. We also wish to change our bounds of integration. The bound $x = -3$ corresponds to $\theta = -\pi/2$ (for when $\theta = -\pi/2$, $x = 3 \sin \theta = -3$). Likewise, the bound of $x = 3$ is replaced by the bound $\theta = \pi/2$. Thus

$$\begin{aligned} \int_{-3}^3 \sqrt{9 - x^2} dx &= \int_{-\pi/2}^{\pi/2} \sqrt{9 - 9 \sin^2 \theta} (3 \cos \theta) d\theta \\ &= \int_{-\pi/2}^{\pi/2} 3\sqrt{9 \cos^2 \theta} \cos \theta d\theta \\ &= \int_{-\pi/2}^{\pi/2} 3|3 \cos \theta| \cos \theta d\theta. \end{aligned}$$

On $[-\pi/2, \pi/2]$, $\cos \theta$ is always positive, so we can drop the absolute value bars, then employ a power-reducing formula:

Notes:

$$\begin{aligned}
 &= \int_{-\pi/2}^{\pi/2} 9 \cos^2 \theta \, d\theta \\
 &= \int_{-\pi/2}^{\pi/2} \frac{9}{2} (1 + \cos(2\theta)) \, d\theta \\
 &= \frac{9}{2} \left(\theta + \frac{1}{2} \sin(2\theta) \right) \Big|_{-\pi/2}^{\pi/2} = \frac{9}{2} \pi.
 \end{aligned}$$

This matches our answer from before.

We now describe in detail Trigonometric Substitution. This method excels when dealing with integrands that contain $\sqrt{a^2 - x^2}$, $\sqrt{x^2 - a^2}$ and $\sqrt{x^2 + a^2}$. The following Key Idea outlines the procedure for each case, followed by more examples.

Key Idea 13 Trigonometric Substitution


- (a) For integrands containing $\sqrt{a^2 - x^2}$:

Let $x = a \sin \theta$, $dx = a \cos \theta \, d\theta$

Thus $\theta = \sin^{-1}(x/a)$, for $-\pi/2 \leq \theta \leq \pi/2$.

On this interval, $\cos \theta \geq 0$, so

$$\sqrt{a^2 - x^2} = a \cos \theta$$




- (b) For integrands containing $\sqrt{x^2 + a^2}$:

Let $x = a \tan \theta$, $dx = a \sec^2 \theta \, d\theta$

Thus $\theta = \tan^{-1}(x/a)$, for $-\pi/2 < \theta < \pi/2$.

On this interval, $\sec \theta > 0$, so

$$\sqrt{x^2 + a^2} = a \sec \theta$$




- (c) For integrands containing $\sqrt{x^2 - a^2}$:

Let $x = a \sec \theta$, $dx = a \sec \theta \tan \theta \, d\theta$

Thus $\theta = \sec^{-1}(x/a)$. If $x/a \geq 1$, then $0 \leq \theta < \pi/2$; if $x/a \leq -1$, then $\pi/2 < \theta \leq \pi$.

We restrict our work to where $x \geq a$, so $x/a \geq 1$, and $0 \leq \theta < \pi/2$. On this interval, $\tan \theta \geq 0$, so

$$\sqrt{x^2 - a^2} = a \tan \theta$$



Notes:

Example 173 Using Trigonometric Substitution

Evaluate $\int \frac{1}{\sqrt{5+x^2}} dx$.

SOLUTION Using Key Idea 13(b), we recognize $a = \sqrt{5}$ and set $x = \sqrt{5} \tan \theta$. This makes $dx = \sqrt{5} \sec^2 \theta d\theta$. We will use the fact that $\sqrt{5+x^2} = \sqrt{5+5 \tan^2 \theta} = \sqrt{5 \sec^2 \theta} = \sqrt{5} \sec \theta$. Substituting, we have:

$$\begin{aligned}\int \frac{1}{\sqrt{5+x^2}} dx &= \int \frac{1}{\sqrt{5+5 \tan^2 \theta}} \sqrt{5} \sec^2 \theta d\theta \\ &= \int \frac{\sqrt{5} \sec^2 \theta}{\sqrt{5} \sec \theta} d\theta \\ &= \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C.\end{aligned}$$

While the integration steps are over, we are not yet done. The original problem was stated in terms of x , whereas our answer is given in terms of θ . We must convert back to x .

The reference triangle given in Key Idea 13(b) helps. With $x = \sqrt{5} \tan \theta$, we have

$$\tan \theta = \frac{x}{\sqrt{5}} \quad \text{and} \quad \sec \theta = \frac{\sqrt{x^2+5}}{\sqrt{5}}.$$

This gives

$$\begin{aligned}\int \frac{1}{\sqrt{5+x^2}} dx &= \ln |\sec \theta + \tan \theta| + C \\ &= \ln \left| \frac{\sqrt{x^2+5}}{\sqrt{5}} + \frac{x}{\sqrt{5}} \right| + C.\end{aligned}$$

We can leave this answer as is, or we can use a logarithmic identity to simplify it. Note:

$$\begin{aligned}\ln \left| \frac{\sqrt{x^2+5}}{\sqrt{5}} + \frac{x}{\sqrt{5}} \right| + C &= \ln \left| \frac{1}{\sqrt{5}} (\sqrt{x^2+5} + x) \right| + C \\ &= \ln \left| \frac{1}{\sqrt{5}} \right| + \ln |\sqrt{x^2+5} + x| + C \\ &= \ln |\sqrt{x^2+5} + x| + C,\end{aligned}$$

where the $\ln(1/\sqrt{5})$ term is absorbed into the constant C . (In Section 6.6 we will learn another way of approaching this problem.)

Notes:

Example 174 Using Trigonometric Substitution

Evaluate $\int \sqrt{4x^2 - 1} dx$.

SOLUTION We start by rewriting the integrand so that it looks like $\sqrt{x^2 - a^2}$ for some value of a :

$$\begin{aligned}\sqrt{4x^2 - 1} &= \sqrt{4\left(x^2 - \frac{1}{4}\right)} \\ &= 2\sqrt{x^2 - \left(\frac{1}{2}\right)^2}.\end{aligned}$$

So we have $a = 1/2$, and following Key Idea 13(c), we set $x = \frac{1}{2} \sec \theta$, and hence $dx = \frac{1}{2} \sec \theta \tan \theta d\theta$. We now rewrite the integral with these substitutions:

$$\begin{aligned}\int \sqrt{4x^2 - 1} dx &= \int 2\sqrt{x^2 - \left(\frac{1}{2}\right)^2} dx \\ &= \int 2\sqrt{\frac{1}{4} \sec^2 \theta - \frac{1}{4}} \left(\frac{1}{2} \sec \theta \tan \theta\right) d\theta \\ &= \int \sqrt{\frac{1}{4}(\sec^2 \theta - 1)} (\sec \theta \tan \theta) d\theta \\ &= \int \sqrt{\frac{1}{4} \tan^2 \theta} (\sec \theta \tan \theta) d\theta \\ &= \int \frac{1}{2} \tan^2 \theta \sec \theta d\theta \\ &= \frac{1}{2} \int (\sec^2 \theta - 1) \sec \theta d\theta \\ &= \frac{1}{2} \int (\sec^3 \theta - \sec \theta) d\theta.\end{aligned}$$

We integrated $\sec^3 \theta$ in Example 170, finding its antiderivatives to be

$$\int \sec^3 \theta d\theta = \frac{1}{2} (\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|) + C.$$

Notes:

Thus

$$\begin{aligned}\int \sqrt{4x^2 - 1} dx &= \frac{1}{2} \int (\sec^3 \theta - \sec \theta) d\theta \\ &= \frac{1}{2} \left(\frac{1}{2} \left(\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta| \right) - \ln |\sec \theta + \tan \theta| \right) + C \\ &= \frac{1}{4} (\sec \theta \tan \theta - \ln |\sec \theta + \tan \theta|) + C.\end{aligned}$$

We are not yet done. Our original integral is given in terms of x , whereas our final answer, as given, is in terms of θ . We need to rewrite our answer in terms of x . With $a = 1/2$, and $x = \frac{1}{2} \sec \theta$, the reference triangle in Key Idea 13(c) shows that

$$\tan \theta = \sqrt{x^2 - 1/4}/(1/2) = 2\sqrt{x^2 - 1/4} \quad \text{and} \quad \sec \theta = 2x.$$

Thus

$$\begin{aligned}\frac{1}{4} (\sec \theta \tan \theta - \ln |\sec \theta + \tan \theta|) + C &= \frac{1}{4} (2x \cdot 2\sqrt{x^2 - 1/4} - \ln |2x + 2\sqrt{x^2 - 1/4}|) + C \\ &= \frac{1}{4} (4x\sqrt{x^2 - 1/4} - \ln |2x + 2\sqrt{x^2 - 1/4}|) + C.\end{aligned}$$

The final answer is given in the last line above, repeated here:

$$\int \sqrt{4x^2 - 1} dx = \frac{1}{4} (4x\sqrt{x^2 - 1/4} - \ln |2x + 2\sqrt{x^2 - 1/4}|) + C.$$

Example 175 Using Trigonometric Substitution

Evaluate $\int \frac{\sqrt{4-x^2}}{x^2} dx$.

SOLUTION We use Key Idea 13(a) with $a = 2$, $x = 2 \sin \theta$, $dx = 2 \cos \theta$ and hence $\sqrt{4-x^2} = 2 \cos \theta$. This gives

$$\begin{aligned}\int \frac{\sqrt{4-x^2}}{x^2} dx &= \int \frac{2 \cos \theta}{4 \sin^2 \theta} (2 \cos \theta) d\theta \\ &= \int \cot^2 \theta d\theta \\ &= \int (\csc^2 \theta - 1) d\theta \\ &= -\cot \theta - \theta + C.\end{aligned}$$

Notes:

We need to rewrite our answer in terms of x . Using the reference triangle found in Key Idea 13(a), we have $\cot \theta = \sqrt{4 - x^2}/x$ and $\theta = \sin^{-1}(x/2)$. Thus

$$\int \frac{\sqrt{4 - x^2}}{x^2} dx = -\frac{\sqrt{4 - x^2}}{x} - \sin^{-1}\left(\frac{x}{2}\right) + C.$$

Trigonometric Substitution can be applied in many situations, even those not of the form $\sqrt{a^2 - x^2}$, $\sqrt{x^2 - a^2}$ or $\sqrt{x^2 + a^2}$. In the following example, we apply it to an integral we already know how to handle.

Example 176 Using Trigonometric Substitution

Evaluate $\int \frac{1}{x^2 + 1} dx$.

SOLUTION We know the answer already as $\tan^{-1} x + C$. We apply Trigonometric Substitution here to show that we get the same answer without inherently relying on knowledge of the derivative of the arctangent function.

Using Key Idea 13(b), let $x = \tan \theta$, $dx = \sec^2 \theta d\theta$ and note that $x^2 + 1 = \tan^2 \theta + 1 = \sec^2 \theta$. Thus

$$\begin{aligned} \int \frac{1}{x^2 + 1} dx &= \int \frac{1}{\sec^2 \theta} \sec^2 \theta d\theta \\ &= \int 1 d\theta \\ &= \theta + C. \end{aligned}$$

Since $x = \tan \theta$, $\theta = \tan^{-1} x$, and we conclude that $\int \frac{1}{x^2 + 1} dx = \tan^{-1} x + C$.

The next example is similar to the previous one in that it does not involve a square-root. It shows how several techniques and identities can be combined to obtain a solution.

Example 177 Using Trigonometric Substitution

Evaluate $\int \frac{1}{(x^2 + 6x + 10)^2} dx$.

SOLUTION We start by completing the square, then make the substitution $u = x + 3$, followed by the trigonometric substitution of $u = \tan \theta$:

$$\int \frac{1}{(x^2 + 6x + 10)^2} dx = \int \frac{1}{(u^2 + 1)^2} du.$$

Notes:

Now make the substitution $u = \tan \theta$, $du = \sec^2 \theta d\theta$:

$$\begin{aligned} &= \int \frac{1}{(\tan^2 \theta + 1)^2} \sec^2 \theta d\theta \\ &= \int \frac{1}{(\sec^2 \theta)^2} \sec^2 \theta d\theta \\ &= \int \cos^2 \theta d\theta. \end{aligned}$$

Applying a power reducing formula, we have

$$\begin{aligned} &= \int \left(\frac{1}{2} + \frac{1}{2} \cos(2\theta) \right) d\theta \\ &= \frac{1}{2}\theta + \frac{1}{4} \sin(2\theta) + C. \end{aligned} \quad (6.2)$$

We need to return to the variable x . As $u = \tan \theta$, $\theta = \tan^{-1} u$. Using the identity $\sin(2\theta) = 2 \sin \theta \cos \theta$ and using the reference triangle found in Key Idea 13(b), we have

$$\frac{1}{4} \sin(2\theta) = \frac{1}{2} \frac{u}{\sqrt{u^2 + 1}} \cdot \frac{1}{\sqrt{u^2 + 1}} = \frac{1}{2} \frac{u}{u^2 + 1}.$$

Finally, we return to x with the substitution $u = x + 3$. We start with the expression in Equation (6.2):

$$\begin{aligned} \frac{1}{2}\theta + \frac{1}{4} \sin(2\theta) + C &= \frac{1}{2} \tan^{-1} u + \frac{1}{2} \frac{u}{u^2 + 1} + C \\ &= \frac{1}{2} \tan^{-1}(x + 3) + \frac{x + 3}{2(x^2 + 6x + 10)} + C. \end{aligned}$$

Stating our final result in one line,

$$\int \frac{1}{(x^2 + 6x + 10)^2} dx = \frac{1}{2} \tan^{-1}(x + 3) + \frac{x + 3}{2(x^2 + 6x + 10)} + C.$$

Our last example returns us to definite integrals, as seen in our first example. Given a definite integral that can be evaluated using Trigonometric Substitution, we could first evaluate the corresponding indefinite integral (by changing from an integral in terms of x to one in terms of θ , then converting back to x) and then evaluate using the original bounds. It is much more straightforward, though, to change the bounds as we substitute.

Notes:

Example 178 Definite integration and Trigonometric Substitution

Evaluate $\int_0^5 \frac{x^2}{\sqrt{x^2 + 25}} dx$.

SOLUTION Using Key Idea 13(b), we set $x = 5 \tan \theta$, $dx = 5 \sec^2 \theta d\theta$, and note that $\sqrt{x^2 + 25} = 5 \sec \theta$. As we substitute, we can also change the bounds of integration.

The lower bound of the original integral is $x = 0$. As $x = 5 \tan \theta$, we solve for θ and find $\theta = \tan^{-1}(x/5)$. Thus the new lower bound is $\theta = \tan^{-1}(0) = 0$. The original upper bound is $x = 5$, thus the new upper bound is $\theta = \tan^{-1}(5/5) = \pi/4$.

Thus we have

$$\begin{aligned}\int_0^5 \frac{x^2}{\sqrt{x^2 + 25}} dx &= \int_0^{\pi/4} \frac{25 \tan^2 \theta}{5 \sec \theta} 5 \sec^2 \theta d\theta \\ &= 25 \int_0^{\pi/4} \tan^2 \theta \sec \theta d\theta.\end{aligned}$$

We encountered this indefinite integral in Example 174 where we found

$$\int \tan^2 \theta \sec \theta d\theta = \frac{1}{2} (\sec \theta \tan \theta - \ln |\sec \theta + \tan \theta|).$$

So

$$\begin{aligned}25 \int_0^{\pi/4} \tan^2 \theta \sec \theta d\theta &= \frac{25}{2} (\sec \theta \tan \theta - \ln |\sec \theta + \tan \theta|) \Big|_0^{\pi/4} \\ &= \frac{25}{2} (\sqrt{2} - \ln(\sqrt{2} + 1)) \\ &\approx 6.661.\end{aligned}$$

The following equalities are very useful when evaluating integrals using Trigonometric Substitution.

Key Idea 14 Useful Equalities with Trigonometric Substitution

1. $\sin(2\theta) = 2 \sin \theta \cos \theta$
2. $\cos(2\theta) = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$
3. $\int \sec^3 \theta d\theta = \frac{1}{2} (\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|) + C$
4. $\int \cos^2 \theta d\theta = \int \frac{1}{2} (1 + \cos(2\theta)) d\theta = \frac{1}{2} (\theta + \sin \theta \cos \theta) + C$

Notes:

Exercises 6.4

Terms and Concepts

1. Trigonometric Substitution works on the same principles as Integration by Substitution, though it can feel “_____”.
2. If one uses Trigonometric Substitution on an integrand containing $\sqrt{25 - x^2}$, then one should set $x = _____$.
3. Consider the Pythagorean Identity $\sin^2 \theta + \cos^2 \theta = 1$.
 - (a) What identity is obtained when both sides are divided by $\cos^2 \theta$?
 - (b) Use the new identity to simplify $9 \tan^2 \theta + 9$.
4. Why does Key Idea 13(a) state that $\sqrt{a^2 - x^2} = a \cos \theta$, and not $|a \cos \theta|$?

Problems

In Exercises 5 – 16, apply Trigonometric Substitution to evaluate the indefinite integrals.

5. $\int \sqrt{x^2 + 1} dx$
6. $\int \sqrt{x^2 + 4} dx$
7. $\int \sqrt{1 - x^2} dx$
8. $\int \sqrt{9 - x^2} dx$
9. $\int \sqrt{x^2 - 1} dx$
10. $\int \sqrt{x^2 - 16} dx$
11. $\int \sqrt{4x^2 + 1} dx$
12. $\int \sqrt{1 - 9x^2} dx$
13. $\int \sqrt{16x^2 - 1} dx$
14. $\int \frac{8}{\sqrt{x^2 + 2}} dx$
15. $\int \frac{3}{\sqrt{7 - x^2}} dx$
16. $\int \frac{5}{\sqrt{x^2 - 8}} dx$

In Exercises 17 – 26, evaluate the indefinite integrals. Some may be evaluated without Trigonometric Substitution.

17. $\int \frac{\sqrt{x^2 - 11}}{x} dx$
18. $\int \frac{1}{(x^2 + 1)^2} dx$
19. $\int \frac{x}{\sqrt{x^2 - 3}} dx$
20. $\int x^2 \sqrt{1 - x^2} dx$
21. $\int \frac{x}{(x^2 + 9)^{3/2}} dx$
22. $\int \frac{5x^2}{\sqrt{x^2 - 10}} dx$
23. $\int \frac{1}{(x^2 + 4x + 13)^2} dx$
24. $\int x^2 (1 - x^2)^{-3/2} dx$
25. $\int \frac{\sqrt{5 - x^2}}{7x^2} dx$
26. $\int \frac{x^2}{\sqrt{x^2 + 3}} dx$

In Exercises 27 – 32, evaluate the definite integrals by making the proper trigonometric substitution and changing the bounds of integration. (Note: each of the corresponding indefinite integrals has appeared previously in this Exercise set.)

27. $\int_{-1}^1 \sqrt{1 - x^2} dx$
28. $\int_4^8 \sqrt{x^2 - 16} dx$
29. $\int_0^2 \sqrt{x^2 + 4} dx$
30. $\int_{-1}^1 \frac{1}{(x^2 + 1)^2} dx$
31. $\int_{-1}^1 \sqrt{9 - x^2} dx$
32. $\int_{-1}^1 x^2 \sqrt{1 - x^2} dx$

Solutions to Odd Exercises

33. $\frac{x^2}{2} + 3x + \ln|x| + C$

35. $\frac{x^3}{3} - \frac{x^2}{2} + x - 2\ln|x+1| + C$

37. $\frac{3}{2}x^2 - 8x + 15\ln|x+1| + C$

39. $\sqrt{7}\tan^{-1}\left(\frac{x}{\sqrt{7}}\right) + C$

41. $14\sin^{-1}\left(\frac{x}{\sqrt{5}}\right) + C$

43. $\frac{5}{4}\sec^{-1}(|x|/4) + C$

45. $\frac{\tan^{-1}\left(\frac{x-1}{\sqrt{7}}\right)}{\sqrt{7}} + C$

47. $-3\sin^{-1}\left(\frac{4-x}{5}\right) + C$

49. $-\frac{1}{3(x^3+3)} + C$

51. $-\sqrt{1-x^2} + C$

53. $-\frac{2}{3}\cos^{\frac{3}{2}}(x) + C$

55. $\frac{7}{3}\ln|3x+2| + C$

57. $\ln|x^2+7x+3| + C$

59. $-\frac{x^2}{2} + 2\ln|x^2-7x+1| + 7x + C$

61. $\tan^{-1}(2x) + C$

63. $\frac{1}{3}\sin^{-1}\left(\frac{3x}{4}\right) + C$

65. $\frac{19}{5}\tan^{-1}\left(\frac{x+6}{5}\right) - \ln|x^2+12x+61| + C$

67. $\frac{x^2}{2} - \frac{9}{2}\ln|x^2+9| + C$

69. $-\tan^{-1}(\cos(x)) + C$

71. $\ln|\sec x + \tan x| + C$ (integrand simplifies to $\sec x$)

73. $\sqrt{x^2-6x+8} + C$

75. $352/15$

77. $1/5$

79. $\pi/2$

81. $\pi/6$

Section 6.2

1. T

3. Determining which functions in the integrand to set equal to “ u ” and which to set equal to “ dv ”.

5. $-e^{-x} - xe^{-x} + C$

7. $-x^3\cos x + 3x^2\sin x + 6x\cos x - 6\sin x + C$

9. $x^3e^x - 3x^2e^x + 6xe^x - 6e^x + C$

11. $1/2e^x(\sin x - \cos x) + C$

13. $1/13e^{2x}(2\sin(3x) - 3\cos(3x)) + C$

15. $-1/2\cos^2 x + C$

17. $x\tan^{-1}(2x) - \frac{1}{4}\ln|4x^2+1| + C$

19. $\sqrt{1-x^2} + x\sin^{-1}x + C$

21. $-\frac{x^2}{4} + \frac{1}{2}x^2\ln|x| + 2x - 2x\ln|x| + C$

23. $\frac{1}{2}x^2\ln(x^2) - \frac{x^2}{2} + C$

25. $2x + x(\ln|x|)^2 - 2x\ln|x| + C$

27. $x\tan(x) + \ln|\cos(x)| + C$

29. $\frac{2}{5}(x-2)^{5/2} + \frac{4}{3}(x-2)^{3/2} + C$

31. $\sec x + C$

33. $-x\csc x - \ln|\csc x + \cot x| + C$

35. $2\sin(\sqrt{x}) - 2\sqrt{x}\cos(\sqrt{x}) + C$

37. $2\sqrt{x}e^{\sqrt{x}} - 2e^{\sqrt{x}} + C$

39. π

41. 0

43. $1/2$

45. $\frac{3}{4e^2} - \frac{5}{4e^4}$

47. $1/5(e^\pi + e^{-\pi})$

Section 6.3

1. F

3. F

5. $\frac{1}{4}\sin^4(x) + C$

7. $\frac{1}{6}\cos^6 x - \frac{1}{4}\cos^4 x + C$

9. $-\frac{1}{9}\sin^9(x) + \frac{3\sin^7(x)}{7} - \frac{3\sin^5(x)}{5} + \frac{\sin^3(x)}{3} + C$

11. $\frac{1}{2}\left(-\frac{1}{8}\cos(8x) - \frac{1}{2}\cos(2x)\right) + C$

13. $\frac{1}{2}\left(\frac{1}{4}\sin(4x) - \frac{1}{10}\sin(10x)\right) + C$

15. $\frac{1}{2}(\sin(x) + \frac{1}{3}\sin(3x)) + C$

17. $\frac{\tan^5(x)}{5} + C$

19. $\frac{\tan^6(x)}{6} + \frac{\tan^4(x)}{4} + C$

21. $\frac{\sec^5(x)}{5} - \frac{\sec^3(x)}{3} + C$

23. $\frac{1}{3}\tan^3 x - \tan x + x + C$

25. $\frac{1}{2}(\sec x \tan x - \ln|\sec x + \tan x|) + C$

27. $\frac{2}{5}$

29. $32/315$

31. $2/3$

33. $16/15$

Section 6.4

1. backwards

3. (a) $\tan^2 \theta + 1 = \sec^2 \theta$

(b) $9\sec^2 \theta$.

5. $\frac{1}{2}\left(x\sqrt{x^2+1} + \ln|\sqrt{x^2+1}+x|\right) + C$

7. $\frac{1}{2}\left(\sin^{-1}x + x\sqrt{1-x^2}\right) + C$

9. $\frac{1}{2}x\sqrt{x^2-1} - \frac{1}{2}\ln|x+\sqrt{x^2-1}| + C$

11. $x\sqrt{x^2+1/4} + \frac{1}{4}\ln|2\sqrt{x^2+1/4}+2x| + C = \frac{1}{2}x\sqrt{4x^2+1} + \frac{1}{4}\ln|\sqrt{4x^2+1}+2x| + C$

13. $4\left(\frac{1}{2}x\sqrt{x^2-1/16} - \frac{1}{32}\ln|4x+4\sqrt{x^2-1/16}|\right) + C = \frac{1}{2}x\sqrt{16x^2-1} - \frac{1}{8}\ln|4x+\sqrt{16x^2-1}| + C$

15. $3\sin^{-1}\left(\frac{x}{\sqrt{7}}\right) + C$ (Trig. Subst. is not needed)

17. $\sqrt{x^2-11} - \sqrt{11}\sec^{-1}(x/\sqrt{11}) + C$

19. $\sqrt{x^2-3} + C$ (Trig. Subst. is not needed)

21. $-\frac{1}{\sqrt{x^2+9}} + C$ (Trig. Subst. is not needed)

23. $\frac{1}{18} \frac{x+2}{x^2+4x+13} + \frac{1}{54} \tan^{-1}\left(\frac{x+2}{2}\right) + C$

25. $\frac{1}{7} \left(-\frac{\sqrt{5-x^2}}{x} - \sin^{-1}(x/\sqrt{5}) \right) + C$

27. $\pi/2$

29. $2\sqrt{2} + 2\ln(1+\sqrt{2})$

31. $9\sin^{-1}(1/3) + \sqrt{8}$ Note: the new lower bound is $\theta = \sin^{-1}(-1/3)$ and the new upper bound is $\theta = \sin^{-1}(1/3)$.
The final answer comes with recognizing that $\sin^{-1}(-1/3) = -\sin^{-1}(1/3)$ and that $\cos(\sin^{-1}(1/3)) = \cos(\sin^{-1}(-1/3)) = \sqrt{8}/3$.

Section 6.5

1. rational

3. $\frac{A}{x} + \frac{B}{x-3}$

5. $\frac{A}{x-\sqrt{7}} + \frac{B}{x+\sqrt{7}}$

7. $3\ln|x-2| + 4\ln|x+5| + C$

9. $\frac{1}{3}(\ln|x+2| - \ln|x-2|) + C$

11. $-\frac{4}{x+8} - 3\ln|x+8| + C$

13. $-\ln|2x-3| + 5\ln|x-1| + 2\ln|x+3| + C$

15. $x + \ln|x-1| - \ln|x+2| + C$

17. $2x + C$

19. $-\frac{3}{2}\ln|x^2+4x+10| + x + \frac{\tan^{-1}\left(\frac{x+2}{\sqrt{6}}\right)}{\sqrt{6}} + C$

21. $2\ln|x-3| + 2\ln|x^2+6x+10| - 4\tan^{-1}(x+3) + C$

23. $\frac{1}{2}(3\ln|x^2+2x+17| - 4\ln|x-7| + \tan^{-1}\left(\frac{x+1}{4}\right)) + C$

25. $\frac{1}{2}\ln|x^2+10x+27| + 5\ln|x+2| - 6\sqrt{2}\tan^{-1}\left(\frac{x+5}{\sqrt{2}}\right) + C$

27. $5\ln(9/4) - \frac{1}{3}\ln(17/2) \approx 3.3413$

29. $1/8$

Section 6.6

1. Because $\cosh x$ is always positive.

3. $\coth^2 x - \operatorname{csch}^2 x = \left(\frac{e^x + e^{-x}}{e^x - e^{-x}}\right)^2 - \left(\frac{2}{e^x - e^{-x}}\right)^2$
 $= \frac{(e^{2x} + 2 + e^{-2x}) - (4)}{e^{2x} - 2 + e^{-2x}}$
 $= \frac{e^{2x} - 2 + e^{-2x}}{e^{2x} - 2 + e^{-2x}}$
 $= 1$

5. $\cosh^2 x = \left(\frac{e^x + e^{-x}}{2}\right)^2$
 $= \frac{e^{2x} + 2 + e^{-2x}}{4}$
 $= \frac{1}{2} \frac{(e^{2x} + e^{-2x}) + 2}{2}$
 $= \frac{1}{2} \left(\frac{e^{2x} + e^{-2x}}{2} + 1\right)$
 $= \frac{\cosh 2x + 1}{2}$

7. $\frac{d}{dx} [\operatorname{sech} x] = \frac{d}{dx} \left[\frac{2}{e^x + e^{-x}} \right]$
 $= \frac{-2(e^x - e^{-x})}{(e^x + e^{-x})^2}$
 $= -\frac{2(e^x - e^{-x})}{(e^x + e^{-x})(e^x + e^{-x})}$
 $= -\frac{2}{e^x + e^{-x}} \cdot \frac{e^x - e^{-x}}{e^x + e^{-x}}$
 $= -\operatorname{sech} x \tanh x$

9. $\int \tanh x \, dx = \int \frac{\sinh x}{\cosh x} \, dx$
Let $u = \cosh x$; $du = (\sinh x)dx$
 $= \int \frac{1}{u} \, du$
 $= \ln|u| + C$
 $= \ln(\cosh x) + C$.

11. $2\sinh 2x$

13. $\coth x$

15. $x \cosh x$

17. $\frac{3}{\sqrt{9x^2+1}}$

19. $\frac{1}{1-(x+5)^2}$

21. $\sec x$

23. $y = 3/4(x - \ln 2) + 5/4$

25. $y = x$

27. $1/2 \ln(\cosh(2x)) + C$

29. $1/2 \sinh^2 x + C$ or $1/2 \cosh^2 x + C$

31. $x \cosh(x) - \sinh(x) + C$

33. $\cosh^{-1}(x^2/2) + C = \ln(x^2 + \sqrt{x^4 - 4}) + C$

35. $\frac{1}{16}\tan^{-1}(x/2) + \frac{1}{32}\ln|x-2| + \frac{1}{32}\ln|x+2| + C$

37. $\tan^{-1}(e^x) + C$

39. $x \tanh^{-1} x + 1/2 \ln|x^2 - 1| + C$

41. 0

43. 2

Section 6.7

1. $0/0, \infty/\infty, 0 \cdot \infty, \infty - \infty, 0^0, 1^\infty, \infty^0$

3. F

5. derivatives; limits

7. Answers will vary.

9. $-5/3$

11. $-\sqrt{2}/2$

13. 0

15. a/b

17. $1/2$

19. 0

21. ∞

23. 0

25. -2

27. 0

29. 0