

Section 6: Linear Equations Theory and Terminology

Recall that an n^{th} order linear IVP consists of an equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

to solve subject to conditions

$$y(x_0) = y_0, \quad y'(x_0) = y_1, \quad \dots, \quad y^{(n-1)}(x_0) = y_{n-1}.$$

The problem is called **homogeneous** if $g(x) \equiv 0$. Otherwise it is called **nonhomogeneous**.

Theorem: Existence & Uniqueness

Theorem: If a_0, \dots, a_n and g are continuous on an interval I , $a_n(x) \neq 0$ for each x in I , and x_0 is any point in I , then for any choice of constants y_0, \dots, y_{n-1} , the IVP has a unique solution $y(x)$ on I .

Put differently, we're guaranteed to have a solution exist, and it is the only one there is!

Homogeneous Equations

We'll consider the equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0$$

and assume that each a_i is continuous and a_n is never zero on the interval of interest.

Theorem: If y_1, y_2, \dots, y_k are all solutions of this homogeneous equation on an interval I , then the *linear combination*

$$y(x) = c_1 y_1(x) + c_2 y_2(x) + \cdots + c_k y_k(x)$$

is also a solution on I for any choice of constants c_1, \dots, c_k .

This is called the **principle of superposition**.

Corollaries

- (i) If y_1 solves the homogeneous equation, then any constant multiple $y = cy_1$ is also a solution.
- (ii) The solution $y = 0$ (called the trivial solution) is always a solution to a homogeneous equation.

Big Questions:

- ▶ Does an equation have any **nontrivial** solution(s), and
- ▶ since y_1 and cy_1 aren't truly *different* solutions, what criteria will be used to call solutions distinct?

Linear Dependence

Definition: A set of functions $f_1(x), f_2(x), \dots, f_n(x)$ are said to be **linearly dependent** on an interval I if there exists a set of constants c_1, c_2, \dots, c_n with at least one of them being nonzero such that

$$c_1 f_1(x) + c_2 f_2(x) + \dots + c_n f_n(x) = 0 \quad \text{for all } x \text{ in } I.$$

A set of functions that is not linearly dependent on I is said to be **linearly independent** on I .

$c_1 f_1(x) + c_2 f_2(x) + \dots + c_n f_n(x) = 0$ can always be made true by taking all c 's to be zero. If the f_1, f_2, \dots, f_n are linearly independent, then this is the only way to make it true.

Example: A linearly Dependent Set

The functions $f_1(x) = x^2$, $f_2(x) = 4x$, and $f_3(x) = x - x^2$ are linearly dependent on $I = (-\infty, \infty)$.

We need to show that $c_1 f_1(x) + c_2 f_2(x) + c_3 f_3(x) = 0$ for all real x for some coefficients c_1, c_2, c_3 not all zero.

Consider $c_1 = 1$, $c_2 = \frac{-1}{4}$ and $c_3 = 1$. Note

$$c_1 f_1(x) + c_2 f_2(x) + c_3 f_3(x) =$$

$$1 \cdot x^2 + \left(\frac{-1}{4}\right)(4x) + 1 \cdot (x - x^2) =$$

$$x^2 - x + x - x^2 = 0 \quad \text{for all } x$$

The functions are linearly dependent - c_1, c_2, c_3 were not all zero (none was zero!).

To see where the c 's come from, consider

$$c_1 f_1(x) + c_2 f_2(x) + c_3 f_3(x) = 0$$

$$c_1 x^2 + c_2 (4x) + c_3 (x - x^2) = 0$$

This will hold as long as $c_1 = c_3$ and $c_2 = \frac{1}{4} c_3$

Example: A linearly Independent Set

The functions $f_1(x) = \sin x$ and $f_2(x) = \cos x$ are linearly independent on $I = (-\infty, \infty)$.

We'll show that $c_1 f_1(x) + c_2 f_2(x) = 0$ for all x
is only true if $c_1 = 0$ and $c_2 = 0$.

Suppose $c_1 f_1(x) + c_2 f_2(x) = 0$ for all real x .

So
$$c_1 \sin x + c_2 \cos x = 0$$

This has to hold when $x=0$. When $x=0$, the equation is

$$c_1 \sin(0) + c_2 \cos(0) = 0$$

$$c_1(0) + c_2(1) = 0 \quad \Rightarrow \quad c_2 = 0$$

The equation has to hold when $x = \pi/2$. When $x = \pi/2$, the equation is

$$c_1 \sin\left(\frac{\pi}{2}\right) + 0 \cdot \overset{c_2}{\cos\left(\frac{\pi}{2}\right)} = 0$$

$$c_1(1) = 0 \Rightarrow c_1 = 0$$

Both c 's must be zero. The functions are linearly independent.

Definition of Wronskian

Let f_1, f_2, \dots, f_n possess at least $n - 1$ continuous derivatives on an interval I . The **Wronskian** of this set of functions is the determinant

$$W(f_1, f_2, \dots, f_n)(x) = \begin{vmatrix} f_1 & f_2 & \cdots & f_n \\ f_1' & f_2' & \cdots & f_n' \\ \vdots & \vdots & \vdots & \vdots \\ f_1^{(n-1)} & f_2^{(n-1)} & \cdots & f_n^{(n-1)} \end{vmatrix}.$$

(Note that, in general, this Wronskian is a function of the independent variable x .)

Determinants

If A is a 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then its determinant

$$\det(A) = ad - bc.$$

If A is a 3×3 matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$, then its determinant

$$\det(A) = a_{11} \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$

Determine the Wronskian of the Functions

$$f_1(x) = \sin x, \quad f_2(x) = \cos x$$

2 functions 2×2 matrix

$$W(f_1, f_2)(x) = \begin{vmatrix} f_1 & f_2 \\ f_1' & f_2' \end{vmatrix} = \begin{vmatrix} \sin x & \cos x \\ \cos x & -\sin x \end{vmatrix}$$

$$= \sin x (-\sin x) - \cos x (\cos x)$$

$$= -\sin^2 x - \cos^2 x$$

$$= -(\sin^2 x + \cos^2 x)$$

$$= -1$$

$$W(f_1, f_2)(x) = -1$$

Determine the Wronskian of the Functions

$$f_1(x) = x^2, \quad f_2(x) = 4x, \quad f_3(x) = x - x^2$$

3 functions \Rightarrow 3×3 matrix

$$W(f_1, f_2, f_3)(x) = \begin{vmatrix} f_1 & f_2 & f_3 \\ f_1' & f_2' & f_3' \\ f_1'' & f_2'' & f_3'' \end{vmatrix} = \begin{vmatrix} x^2 & 4x & x-x^2 \\ 2x & 4 & 1-2x \\ 2 & 0 & -2 \end{vmatrix}$$

$$= x^2 \begin{vmatrix} 4 & 1-2x \\ 0 & -2 \end{vmatrix} - 4x \begin{vmatrix} 2x & 1-2x \\ 2 & -2 \end{vmatrix} + (x-x^2) \begin{vmatrix} 2x & 4 \\ 2 & 0 \end{vmatrix}$$

$$= x^2 (4(-2) - 0(1-2x)) - 4x (2x(-2) - 2(1-2x)) + (x-x^2) (2x(0) - 2(4))$$

$$= x^2(-8) - 4x(-4x - 2 + 4x) + (x-x^2)(-8)$$

$$= -8x^2 - 4x(-2) - 8x + 8x^2$$

$$= 8x - 8x = 0$$

$$W(f_1, f_2, f_3)(x) = 0$$

Theorem (a test for linear independence)

Let f_1, f_2, \dots, f_n be $n - 1$ times continuously differentiable on an interval I . If there exists x_0 in I such that $W(f_1, f_2, \dots, f_n)(x_0) \neq 0$, then the functions are **linearly independent** on I .

If y_1, y_2, \dots, y_n are n solutions of the linear homogeneous n^{th} order equation on an interval I , then the solutions are **linearly independent** on I if and only if $W(y_1, y_2, \dots, y_n)(x) \neq 0$ for¹ each x in I .

¹For solutions of one linear homogeneous ODE, the Wronskian is either always zero or is never zero.

Determine if the functions are linearly dependent or independent:

$$y_1 = x^2, \quad y_2 = x^3 \quad I = (0, \infty)$$

We'll use the Wronskian.

$$W(y_1, y_2)(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} x^2 & x^3 \\ 2x & 3x^2 \end{vmatrix}$$

$$= x^2(3x^2) - 2x(x^3) = 3x^4 - 2x^4 = x^4$$

$$W(y_1, y_2)(x) = x^4$$

$W \neq 0$, the functions are linearly independent.

Fundamental Solution Set

We're still considering this equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0$$

with the assumptions $a_n(x) \neq 0$ and $a_i(x)$ are continuous on I .

Definition: A set of functions y_1, y_2, \dots, y_n is a **fundamental solution set** of the n^{th} order homogeneous equation provided they

- (i) are solutions of the equation,
- (ii) there are n of them, and
- (iii) they are linearly independent.

Theorem: Under the assumed conditions, the equation has a fundamental solution set.