

Section 6: Linear Equations Theory and Terminology

We'll consider the equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0$$

and assume that each a_i is continuous and a_n is never zero on the interval of interest.

Theorem: If y_1, y_2, \dots, y_k are all solutions of this homogeneous equation on an interval I , then the *linear combination*

$$y(x) = c_1 y_1(x) + c_2 y_2(x) + \cdots + c_k y_k(x)$$

is also a solution on I for any choice of constants c_1, \dots, c_k .

This is called the **principle of superposition.**

Linear Dependence

Definition: A set of functions $f_1(x), f_2(x), \dots, f_n(x)$ are said to be **linearly dependent** on an interval I if there exists a set of constants c_1, c_2, \dots, c_n with at least one of them being nonzero such that

$$c_1 f_1(x) + c_2 f_2(x) + \dots + c_n f_n(x) = 0 \quad \text{for all } x \text{ in } I.$$

A set of functions that is not linearly dependent on I is said to be **linearly independent** on I .

Examples

We saw that $f_1(x) = x^2$, $f_2(x) = 4x$, $f_3(x) = x - x^2$ were linearly dependent on $(-\infty, \infty)$ because

$$4x^2 - 1(4x) + 4(x - x^2) = 0 \quad \text{for all real } x$$

At least one (in fact all) of the coefficients are nonzero. By comparison, $f_1(x) = \sin(x)$, $f_2(x) = \cos(x)$ are linearly independent on $(-\infty, \infty)$ because the equation

$$c_1 \sin(x) + c_2 \cos(x) = 0 \quad \text{for all real } x$$

is ONLY true if $c_1 = 0$ and $c_2 = 0$.

We would like a general tool to determine linear dependence/independence at least in the case that we have sufficiently differentiable functions (like the ones that solve differential equations).

Definition of Wronskian

Let f_1, f_2, \dots, f_n posses at least $n - 1$ continuous derivatives on an interval I . The **Wronskian** of this set of functions is the determinant

$$W(f_1, f_2, \dots, f_n)(x) = \begin{vmatrix} f_1 & f_2 & \cdots & f_n \\ f_1' & f_2' & \cdots & f_n' \\ \vdots & \vdots & \vdots & \vdots \\ f_1^{(n-1)} & f_2^{(n-1)} & \cdots & f_n^{(n-1)} \end{vmatrix}.$$

(Note that, in general, this Wronskian is a function of the independent variable x .)

Determinants

If A is a 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then its determinant

$$\det(A) = ad - bc.$$

If A is a 3×3 matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$, then its determinant

$$\det(A) = a_{11}\det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12}\det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13}\det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$

Determine the Wronskian of the Functions

$$f_1(x) = \sin x, \quad f_2(x) = \cos x$$

2 functions \rightarrow 2x2 matrix

$$W(f_1, f_2)(x) = \begin{vmatrix} f_1(x) & f_2(x) \\ f_1'(x) & f_2'(x) \end{vmatrix}$$

$$= \begin{vmatrix} \sin x & \cos x \\ \cos x & -\sin x \end{vmatrix}$$

$$= \sin x (-\sin x) - \cos x (\cos x)$$

$$= -\sin^2 x - \cos^2 x$$

$$= -(\sin^2 x + \cos^2 x)$$

$$= -1$$

$$W(f_1, f_2)(x) = -1.$$

Determine the Wronskian of the Functions

$$f_1(x) = x^2, \quad f_2(x) = 4x, \quad f_3(x) = x - x^2$$

3 functions \rightarrow 3×3 matrix

$$W(f_1, f_2, f_3)(x) = \begin{vmatrix} f_1(x) & f_2(x) & f_3(x) \\ f_1'(x) & f_2'(x) & f_3'(x) \\ f_1''(x) & f_2''(x) & f_3''(x) \end{vmatrix}$$

$$= \begin{vmatrix} x^2 & 4x & x - x^2 \\ 2x & 4 & 1 - 2x \\ 2 & 0 & -2 \end{vmatrix}$$

$$= x^2 \begin{vmatrix} 4 & 1-2x \\ 0 & -2 \end{vmatrix} - 4x \begin{vmatrix} 2x & 1-2x \\ 2 & -2 \end{vmatrix} + (x-x^2) \begin{vmatrix} 2x & 4 \\ 2 & 0 \end{vmatrix}$$

$$= x^2(4(-2) - 0(1-2x)) - 4x(2x(-2) - 2(1-2x)) + (x-x^2)(2x(0) - 2(4))$$

$$= x^2(-8) - 4x(-4x - 2 + 4x) + (x-x^2)(-8)$$

$$= -8x^2 + 8x - 8x + 8x^2$$

$$= 0$$

$$W(f_1, f_2, f_3)(x) = 0$$

Theorem (a test for linear independence)

Let f_1, f_2, \dots, f_n be $n - 1$ times continuously differentiable on an interval I . If there exists x_0 in I such that $W(f_1, f_2, \dots, f_n)(x_0) \neq 0$, then the functions are **linearly independent** on I .

If y_1, y_2, \dots, y_n are n solutions of the linear homogeneous n^{th} order equation on an interval I , then the solutions are **linearly independent** on I if and only if $W(y_1, y_2, \dots, y_n)(x) \neq 0$ for¹ each x in I .

¹For solutions of one linear homogeneous ODE, the Wronskian is either always zero or is never zero.

Determine if the functions are linearly dependent or independent:

$$y_1 = e^x, \quad y_2 = e^{-2x} \quad I = (-\infty, \infty)$$

We can use the Wronskian.

$$\begin{aligned} W(y_1, y_2)(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \\ &= \begin{vmatrix} e^x & e^{-2x} \\ e^x & -2e^{-2x} \end{vmatrix} \end{aligned}$$

$$= e^x (-2e^{-2x}) - e^x (e^{-2x})$$

$$= -2e^{-x} - e^{-x} = -3e^{-x}$$

$$W(y_1, y_2)(x) = -3e^{-x} \quad \text{nonzero}$$

The functions are linearly independent.

Fundamental Solution Set

We're still considering this equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0$$

with the assumptions $a_n(x) \neq 0$ and $a_i(x)$ are continuous on I .

Definition: A set of functions y_1, y_2, \dots, y_n is a **fundamental solution set** of the n^{th} order homogeneous equation provided they

- (i) are solutions of the equation,
- (ii) there are n of them, and
- (iii) they are linearly independent.

Theorem: Under the assumed conditions, the equation has a fundamental solution set.

General Solution of n^{th} order Linear Homogeneous Equation

Let y_1, y_2, \dots, y_n be a fundamental solution set of the n^{th} order linear homogeneous equation. Then the **general solution** of the equation is

$$y(x) = c_1 y_1(x) + c_2 y_2(x) + \cdots + c_n y_n(x),$$

where c_1, c_2, \dots, c_n are arbitrary constants.

Example

Verify that $y_1 = x^2$ and $y_2 = x^3$ form a fundamental solution set of the ODE

$$x^2 y'' - 4xy' + 6y = 0 \quad \text{on } (0, \infty),$$

and determine the general solution.

We need to show that y_1, y_2 are solutions and that they are linearly independent.

Show y_1 is a solution:

$$y_1 = x^2, \quad y_1' = 2x, \quad y_1'' = 2$$

$$\begin{aligned} x^2 y_1'' - 4x y_1' + 6y_1 &= x^2(2) - 4x(2x) + 6(x^2) \\ &= 2x^2 - 8x^2 + 6x^2 = 0 \end{aligned}$$

y_1 is a solution.

Show y_2 solves the ODE.

$$y_2 = x^3, \quad y_2' = 3x^2, \quad y_2'' = 6x$$

$$\begin{aligned} x^2 y_2'' - 4x y_2' + 6y_2 &= x^2(6x) - 4x(3x^2) + 6(x^3) \\ &= 6x^3 - 12x^3 + 6x^3 \\ &= 0 \end{aligned}$$

So y_2 solves it as well.

We can use the Wronskian to show they

are linearly independent,

$$\begin{aligned} W(y_1, y_2)(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \\ &= \begin{vmatrix} x^2 & x^3 \\ 2x & 3x^2 \end{vmatrix} = x^2(3x^2) - 2x(x^3) \\ &= 3x^4 - 2x^4 = x^4 \end{aligned}$$

$$W(y_1, y_2) = x^4 \neq 0$$

They are lin. independent. We have a fundamental solution set. The general solution

$$y = C_1 x^2 + C_2 x^3.$$

Nonhomogeneous Equations

Now we will consider the equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

where g is not the zero function. We'll continue to assume that a_n doesn't vanish and that a_i and g are continuous.

The **associated homogeneous equation** is

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0.$$

Theorem: General Solution of Nonhomogeneous Equation

Let y_p be any solution of the nonhomogeneous equation, and let y_1, y_2, \dots, y_n be any fundamental solution set of the associated homogeneous equation.

Then the general solution of the nonhomogeneous equation is

$$y = \underbrace{c_1 y_1(x) + c_2 y_2(x) + \cdots + c_n y_n(x)}_{y_c} + y_p(x)$$

where c_1, c_2, \dots, c_n are arbitrary constants.

Note the form of the solution $y_c + y_p$!
(complementary plus particular)

Another Superposition Principle (for nonhomogeneous eqns.)

Let $y_{p_1}, y_{p_2}, \dots, y_{p_k}$ be k particular solutions to the nonhomogeneous linear equations

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g_i(x)$$

for $i = 1, \dots, k$. Assume the domain of definition for all k equations is a common interval I .

Then

$$y_p = y_{p_1} + y_{p_2} + \cdots + y_{p_k}$$

is a particular solution of the nonhomogeneous equation

$$a_n(x) \frac{d^n y}{dx^n} + \cdots + a_0(x)y = g_1(x) + g_2(x) + \cdots + g_k(x).$$

Example $x^2 y'' - 4xy' + 6y = 36 - 14x$

We will construct the general solution by considering sub-problems.

(a) **Part 1** Verify that

$$y_{p_1} = 6 \quad \text{solves} \quad x^2 y'' - 4xy' + 6y = 36.$$

$$y_{p_1}' = 0$$

$$y_{p_1}'' = 0$$

$$x^2 y_{p_1}'' - 4x y_{p_1}' + 6y_{p_1} =$$

$$x^2(0) - 4x(0) + 6(6) = 36$$

So y_{p_1} is a particular solution.

Example $x^2y'' - 4xy' + 6y = 36 - 14x$

(b) **Part 2** Verify that

$$y_{p2} = -7x \quad \text{solves} \quad x^2y'' - 4xy' + 6y = -14x.$$

$$y_{p2}' = -7$$

$$x^2y_{p2}'' - 4xy_{p2}' + 6y_{p2} =$$

$$y_{p2}'' = 0$$

$$x^2(0) - 4x(-7) + 6(-7x) =$$

$$28x - 42x = -14x$$

y_{p2} is a particular solution.

Example $x^2y'' - 4xy' + 6y = 36 - 14x$

(c) **Part 3** We already know that $y_1 = x^2$ and $y_2 = x^3$ is a fundamental solution set of

$$x^2y'' - 4xy' + 6y = 0. \quad \leftarrow \text{associated homogeneous eqn.}$$

Use this along with results (a) and (b) to write the general solution of $x^2y'' - 4xy' + 6y = 36 - 14x$.

$$y_c = C_1y_1 + C_2y_2 = C_1x^2 + C_2x^3$$

By the principle of superposition $y_p = y_{p1} + y_{p2} = 6 - 7x$

The general solution to the nonhomogeneous equation is $y = C_1x^2 + C_2x^3 + 6 - 7x$.

Solve the IVP

$$x^2 y'' - 4xy' + 6y = 36 - 14x, \quad y(1) = 0, \quad y'(1) = 5$$

The general solution to the ODE is

$$y = C_1 x^2 + C_2 x^3 + 6 - 7x$$

So $y' = 2C_1 x + 3C_2 x^2 - 7$

Applying the I.C.

$$y(1) = C_1(1)^2 + C_2(1)^3 + 6 - 7 \cdot 1 = 0$$

$$C_1 + C_2 - 1 = 0$$

$$C_1 + C_2 = 1$$

$$y'(1) = 2C_1 \cdot 1 + 3C_2(1)^2 - 7 = 5$$

$$2C_1 + 3C_2 - 7 = 5$$

$$2C_1 + 3C_2 = 12$$

$$\begin{array}{rcl} C_1 + C_2 = 1 & \Rightarrow & 2C_1 + 2C_2 = 2 \\ 2C_1 + 3C_2 = 12 & & \underline{-(2C_1 + 3C_2 = 12)} \end{array}$$

$$-C_2 = -10$$

$$C_2 = 10$$

$$C_1 = 1 - C_2 = -9$$

The solution to the IVP is

$$y = -9x^2 + 10x^3 + 6 - 7x$$