

Section 2.2: Inverse of a Matrix

If A is an $n \times n$ matrix, a matrix A^{-1} that satisfies

$$A^{-1}A = AA^{-1} = I_n.$$

is called the inverse of A .

If such matrix A^{-1} exists, we'll say that A is **nonsingular** (a.k.a. *invertible*). Otherwise, we'll say that A is **singular**.

Theorem (2×2 case)

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$, then A is invertible and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

If $ad - bc = 0$, then A is singular.

The number $ad - bc$ is called the determinant of the matrix A .

Theorem

If A is an invertible $n \times n$ matrix, then for each \mathbf{b} in \mathbb{R}^n , the equation $A\mathbf{x} = \mathbf{b}$ has unique solution $\mathbf{x} = A^{-1}\mathbf{b}$.

- ▶ This provides a solution technique for $n \times n$ systems, and in terms of coefficient matrices
- ▶ suggests that being singular or nonsingular is related to consistency for a linear system,
- ▶ suggests a connection between a *determinant*¹ and consistency,
- ▶ suggests a connection between singular/nonsingular and the linear dependence/independence of matrix columns.

¹We will have to define this, and we will.

Theorem

(i) If A is invertible, then A^{-1} is also invertible and

$$(A^{-1})^{-1} = A.$$

(ii) If A and B are invertible $n \times n$ matrices, then the product AB is also invertible² with

$$(AB)^{-1} = B^{-1}A^{-1}.$$

also for A, B, C invertible $(ABC)^{-1} = C^{-1}B^{-1}A^{-1}$

(iii) If A is invertible, then so is A^T . Moreover

$$(A^T)^{-1} = (A^{-1})^T.$$

²This can generalize to the product of k invertible matrices. 

Elementary Matrices

Definition: An **elementary** matrix is a square matrix obtained from the identity by performing one elementary row operation.

Examples:

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}, \quad E_3 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$3R_2 \rightarrow R_2$$

$$2R_1 + R_3 \rightarrow R_3$$

$$R_1 \leftrightarrow R_2$$

Action of Elementary Matrices

Let $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$, and compute the following products

E_1A , E_2A , and E_3A .

$$E_1A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$= \begin{bmatrix} a & b & c \\ 3d & 3e & 3f \\ g & h & i \end{bmatrix}$$

↪ scales
row 2
by 3

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$= \begin{bmatrix} a & b & c \\ d & e & f \\ 2a+g & 2b+h & 2c+i \end{bmatrix}$$

$$E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

performers:
 $2R_1 + R_3 \rightarrow R_3$

$$A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$E_3 A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$= \begin{bmatrix} d & e & f \\ a & b & c \\ g & h & i \end{bmatrix}$$

swaps

$R_1 \leftrightarrow R_2$

$$E_3 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Remarks

- ▶ Elementary row operations can be equated with matrix multiplication (multiply on the left by an elementary matrix),
- ▶ Each elementary matrix is invertible where the inverse *undoes* the row operation,
- ▶ Reduction to rref is a sequence of row operations, so it is a sequence of matrix multiplications

$$\text{rref}(A) = E_k \cdots E_2 E_1 A.$$

Theorem

An $n \times n$ matrix A is invertible if and only if it is row equivalent to the identity matrix I_n . Moreover, if

$$\text{rref}(A) = E_k \cdots E_2 E_1 A = I_n, \quad \text{then} \quad A = (E_k \cdots E_2 E_1)^{-1} I_n.$$

That is,

$$A^{-1} = \left[(E_k \cdots E_2 E_1)^{-1} \right]^{-1} = E_k \cdots E_2 E_1.$$

The sequence of operations that reduces A to I_n , transforms I_n into A^{-1} .

This last observation—operations that take A to I_n also take I_n to A^{-1} —gives us a method for computing an inverse!

Algorithm for finding A^{-1}

To find the inverse of a given matrix A :

- ▶ Form the $n \times 2n$ augmented matrix $[A \quad I]$.
- ▶ Perform whatever row operations are needed to get the first n columns (the A part) to rref.
- ▶ If $\text{rref}(A)$ is I , then $[A \quad I]$ is row equivalent to $[I \quad A^{-1}]$, and the inverse A^{-1} will be the last n columns of the reduced matrix.
- ▶ If $\text{rref}(A)$ is NOT I , then A is not invertible.

Remarks: We don't need to know ahead of time if A is invertible to use this algorithm.

If A is singular, we can stop as soon as it's clear that $\text{rref}(A) \neq I$.

Examples: Find the Inverse if Possible

$$(a) \begin{bmatrix} 1 & 2 & -1 \\ -4 & -7 & 3 \\ -2 & -6 & 4 \end{bmatrix}$$

Form an augmented matrix

$$[A \ I]$$

$A =$

$$[A \ I] = \begin{bmatrix} 1 & 2 & -1 & 1 & 0 & 0 \\ -4 & -7 & 3 & 0 & 1 & 0 \\ -2 & -6 & 4 & 0 & 0 & 1 \end{bmatrix}$$

Do row ops

$$4R_1 + R_2 \rightarrow R_2$$

$$2R_1 + R_3 \rightarrow R_3$$

$$\begin{bmatrix} 1 & 2 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 4 & 1 & 0 \\ 0 & -2 & 2 & 2 & 0 & 1 \end{bmatrix}$$

$$2R_2 + R_3 \rightarrow R_3$$

$$\begin{bmatrix} 1 & 2 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 4 & 1 & 0 \\ 0 & 0 & 0 & 10 & 2 & 1 \end{bmatrix}$$



need
a non zero
entry here,
but can't get one

$\text{ref}(A)$ is not I , A^{-1} doesn't
exist; A is singular.