

## Section 2.2: Inverse of a Matrix

If  $A$  is an  $n \times n$  matrix, a matrix  $A^{-1}$  that satisfies

$$A^{-1}A = AA^{-1} = I_n.$$

is called the inverse of  $A$ .

If such matrix  $A^{-1}$  exists, we'll say that  $A$  is **nonsingular** (a.k.a. *invertible*). Otherwise, we'll say that  $A$  is **singular**.

## Theorem ( $2 \times 2$ case)

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . If  $ad - bc \neq 0$ , then  $A$  is invertible and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

If  $ad - bc = 0$ , then  $A$  is singular.

$ad - bc$  is called the determinant  
of  $A$ .

# Theorem

If  $A$  is an invertible  $n \times n$  matrix, then for each  $\mathbf{b}$  in  $\mathbb{R}^n$ , the equation  $A\mathbf{x} = \mathbf{b}$  has unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .

- ▶ This provides a solution technique for  $n \times n$  systems, and in terms of coefficient matrices
- ▶ suggests that being singular or nonsingular is related to consistency for a linear system,
- ▶ suggests a connection between a *determinant*<sup>1</sup> and consistency,
- ▶ suggests a connection between singular/nonsingular and the linear dependence/independence of matrix columns.

---

<sup>1</sup>We will have to define this, and we will.

# Theorem

(i) If  $A$  is invertible, then  $A^{-1}$  is also invertible and

$$(A^{-1})^{-1} = A.$$

(ii) If  $A$  and  $B$  are invertible  $n \times n$  matrices, then the product  $AB$  is also invertible<sup>2</sup> with

$$(AB)^{-1} = B^{-1}A^{-1}.$$

If  $A, B, C$  are nonsingular  $(ABC)^{-1} = C^{-1}B^{-1}A^{-1}$

(iii) If  $A$  is invertible, then so is  $A^T$ . Moreover

$$(A^T)^{-1} = (A^{-1})^T.$$

---

<sup>2</sup>This can generalize to the product of  $k$  invertible matrices. 

# Elementary Matrices

**Definition:** An **elementary** matrix is a square matrix obtained from the identity by performing one elementary row operation.

Examples:

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}, \quad E_3 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$3R_2 \rightarrow R_2$$

$$2R_1 + R_3 \rightarrow R_3$$

$$R_2 \leftrightarrow R_1$$

## Action of Elementary Matrices

Let  $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$ , and compute the following products

$E_1A$ ,  $E_2A$ , and  $E_3A$ .

$$E_1A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ 3d & 3e & 3f \\ g & h & i \end{bmatrix} \quad 3R_2 \rightarrow R_2$$

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$E_2 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$2R_1 + R_3 \rightarrow R_3$

$$= \begin{bmatrix} a & b & c \\ d & e & f \\ 2a+g & 2b+h & 2c+i \end{bmatrix}$$

$$E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$E_3 A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$= \begin{bmatrix} d & e & f \\ a & b & c \\ g & h & i \end{bmatrix}$$

$R_1 \leftrightarrow R_2$

$$E_3 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



## Remarks

- ▶ Elementary row operations can be equated with matrix multiplication (multiply on the left by an elementary matrix),
- ▶ Each elementary matrix is invertible where the inverse *undoes* the row operation,
- ▶ Reduction to rref is a sequence of row operations, so it is a sequence of matrix multiplications

$$\text{rref}(A) = E_k \cdots E_2 E_1 A.$$

## Theorem

An  $n \times n$  matrix  $A$  is invertible if and only if it is row equivalent to the identity matrix  $I_n$ . Moreover, if

$$\text{rref}(A) = E_k \cdots E_2 E_1 A = I_n, \quad \text{then} \quad A = (E_k \cdots E_2 E_1)^{-1} I_n.$$

That is,

$$A^{-1} = \left[ (E_k \cdots E_2 E_1)^{-1} \right]^{-1} = E_k \cdots E_2 E_1.$$

The sequence of operations that reduces  $A$  to  $I_n$ , transforms  $I_n$  into  $A^{-1}$ .

**This last observation—operations that take  $A$  to  $I_n$  also take  $I_n$  to  $A^{-1}$ —gives us a method for computing an inverse!**

## Algorithm for finding $A^{-1}$

To find the inverse of a given matrix  $A$ :

- ▶ Form the  $n \times 2n$  augmented matrix  $[A \quad I]$ .
- ▶ Perform whatever row operations are needed to get the first  $n$  columns (the  $A$  part) to rref.
- ▶ If  $\text{rref}(A)$  is  $I$ , then  $[A \quad I]$  is row equivalent to  $[I \quad A^{-1}]$ , and the inverse  $A^{-1}$  will be the last  $n$  columns of the reduced matrix.
- ▶ If  $\text{rref}(A)$  is NOT  $I$ , then  $A$  is not invertible.

**Remarks:** We don't need to know ahead of time if  $A$  is invertible to use this algorithm.

If  $A$  is singular, we can stop as soon as it's clear that  $\text{rref}(A) \neq I$ .

## Examples: Find the Inverse if Possible

$$(a) \begin{bmatrix} 1 & 2 & -1 \\ -4 & -7 & 3 \\ -2 & -6 & 4 \end{bmatrix}$$

$A =$

$$\left[ \begin{array}{ccc|ccc} 1 & 2 & -1 & 1 & 0 & 0 \\ -4 & -7 & 3 & 0 & 1 & 0 \\ -2 & -6 & 4 & 0 & 0 & 1 \end{array} \right]$$

Form an augmented matrix

$$[A \ I]$$

$$4R_1 + R_2 \rightarrow R_2$$

$$2R_1 + R_3 \rightarrow R_3$$

$$\left[ \begin{array}{ccc|ccc} 1 & 2 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 4 & 1 & 0 \\ 0 & -2 & 2 & 2 & 0 & 1 \end{array} \right]$$

$$2R_2 + R_3 \rightarrow R_3$$

$$\begin{bmatrix} 1 & 2 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 4 & 1 & 0 \\ 0 & 0 & 0 & 10 & 2 & 1 \end{bmatrix}$$



need a nonzero.  
entry here which  
isn't possible.

$$\text{rref}(A) \neq I_3$$

So  $A$  is singular, it doesn't have  
an inverse.