

Section 8: Homogeneous Equations with Constant Coefficients

We consider a second order, linear, homogeneous equation with constant coefficients

$$a \frac{d^2 y}{dx^2} + b \frac{dy}{dx} + cy = 0.$$

We found that $y = e^{mx}$ is a solution provided m is a solution to the equation

$$am^2 + bm + c = 0$$

called the characteristic (or auxiliary) equation.

Auxiliary a.k.a. Characteristic Equation

$$am^2 + bm + c = 0$$

There are three cases:

- I $b^2 - 4ac > 0$ and there are two distinct real roots $m_1 \neq m_2$
- II $b^2 - 4ac = 0$ and there is one repeated real root $m_1 = m_2 = m$
- III $b^2 - 4ac < 0$ and there are two roots that are complex conjugates
 $m_{1,2} = \alpha \pm i\beta$

Case I: Two distinct real roots

$$ay'' + by' + cy = 0, \quad \text{where } b^2 - 4ac > 0$$

The values of m are

$$m_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

The general solution is

$$y = c_1 e^{m_1 x} + c_2 e^{m_2 x}$$

Example

Solve the IVP

$$y'' + y' - 12y = 0, \quad y(0) = 1, \quad y'(0) = 10$$

2nd order, linear, homogeneous, constant coefficient

Characteristic equation: $m^2 + m - 12 = 0$

$$(m+4)(m-3) = 0$$

$$m_1 = -4, \quad m_2 = 3$$

Solutions $y_1 = e^{-4x}$, $y_2 = e^{3x}$

The general solution to the ODE is $y = c_1 e^{-4x} + c_2 e^{3x}$

$$\text{Apply } y(0)=1, y'(0)=10$$

$$y' = -4c_1 e^{-4x} + 3c_2 e^{3x}$$

$$y(0) = c_1 e^0 + c_2 e^0 = 1 \Rightarrow c_1 + c_2 = 1$$

$$y'(0) = -4c_1 e^0 + 3c_2 e^0 = 10 \Rightarrow -4c_1 + 3c_2 = 10$$

$$\begin{aligned} 4c_1 + 4c_2 &= 4 \Rightarrow 7c_2 = 14 & c_2 &= 2 \\ -4c_1 + 3c_2 &= 10 \end{aligned}$$

$$c_1 = 1 - c_2 = 1 - 2 = -1$$

The solution to the IVP

is

$$y = -e^{-4x} + 2e^{3x}$$

Case II: One repeated real root

$$ay'' + by' + cy = 0, \quad \text{where } b^2 - 4ac = 0$$

$$y = c_1 e^{mx} + c_2 x e^{mx} \quad \text{where } m = \frac{-b}{2a}$$

Use reduction of order to show that if $y_1 = e^{\frac{-bx}{2a}}$, then $y_2 = x e^{\frac{-bx}{2a}}$.

$$y_1 = e^{\frac{-b}{2a}x} \text{ is known. } y_2 = u y_1, \text{ where } u = \int \frac{-\int P(x) dx}{(y_1)^2} dx$$

Standard form

$$y'' + \frac{b}{a}y' + \frac{c}{a}y = 0 \quad P(x) = \frac{b}{a}$$

$$e^{-\int P(x) dx} = e^{-\int \frac{b}{a} dx} = e^{-\frac{b}{a}x}$$

$$(y_1)^2 = \left(e^{\frac{-b}{2a}x} \right)^2 = e^{2\left(\frac{-b}{2a}x\right)} = e^{\frac{-b}{a}x}$$

$$u = \int \frac{-f(x)dx}{(y_1)^2} = \int \frac{e^{\frac{-b}{a}x}}{e^{\frac{-b}{a}x}} dx = \int dx = x$$

$$y_2 = uy_1 = x e^{\frac{-b}{2a}x}$$

You can show that $y_1 = e^{\frac{-b}{2a}x}$ and $y_2 = x e^{\frac{-b}{2a}x}$
are linearly independent

Example

Solve the ODE

2nd order, linear, constant coefficient, homogeneous

$$4y'' - 4y' + y = 0$$

Characteristic eqn: $4m^2 - 4m + 1 = 0$

$$(2m - 1)^2 = 0 \Rightarrow m = \frac{1}{2} \text{ repeated}$$

$$y_1 = e^{\frac{1}{2}x} \quad \text{and} \quad y_2 = x e^{\frac{1}{2}x}$$

The General solution is

$$y = C_1 e^{\frac{1}{2}x} + C_2 x e^{\frac{1}{2}x}$$

Case III: Complex conjugate roots

$$ay'' + by' + cy = 0, \quad \text{where } b^2 - 4ac < 0$$

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x)), \quad \text{where the roots}$$

$$m = \alpha \pm i\beta, \quad \alpha = \frac{-b}{2a} \quad \text{and} \quad \beta = \frac{\sqrt{4ac - b^2}}{2a}$$

The solutions can be written as

$$Y_1 = e^{(\alpha+i\beta)x} = e^{\alpha x} e^{i\beta x}, \quad \text{and} \quad Y_2 = e^{(\alpha-i\beta)x} = e^{\alpha x} e^{-i\beta x}.$$

$$e^{dx+i\beta x}$$

$$e^{dx-i\beta x}$$

Deriving the solutions Case III

Recall Euler's Formula:

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$\begin{aligned} Y_1 &= e^{\alpha x} e^{i\beta x} = e^{\alpha x} (\cos(\beta x) + i \sin(\beta x)) \\ &= e^{\alpha x} \cos(\beta x) + i e^{\alpha x} \sin(\beta x) \end{aligned}$$

$$\cos(-\beta x) = \cos(\beta x)$$

$$\sin(-\beta x) = -\sin(\beta x)$$

$$\begin{aligned} Y_2 &= e^{\alpha x} e^{-i\beta x} = e^{\alpha x} (\cos(\beta x) - i \sin(\beta x)) \\ &= e^{\alpha x} \cos(\beta x) - i e^{\alpha x} \sin(\beta x) \end{aligned}$$

By superposition we can take $y_1, y_2 = k_1 Y_1 + k_2 Y_2$

Let

$$y_1 = \frac{1}{2} (Y_1 + Y_2) = \frac{1}{2} (2 e^{\alpha x} \cos(\beta x) + i \cdot 0) = e^{\alpha x} \cos(\beta x)$$

and

$$y_2 = \frac{1}{2i} (Y_1 - Y_2) = \frac{1}{2i} (0 + i 2 e^{\alpha x} \sin(\beta x))$$

$$= e^{\alpha x} \sin(\beta x)$$

Our fundamental solution set is

$$y_1 = e^{\alpha x} \cos(\beta x), \quad y_2 = e^{\alpha x} \sin(\beta x)$$