

Section 1.8: Intro to Linear Transformations

Recall that the product $A\mathbf{x}$ is a linear combination of the columns of A —turns out to be a vector. If the columns of A are vectors in \mathbb{R}^m , and there are n of them, then

- ▶ A is an $m \times n$ matrix,
- ▶ the product $A\mathbf{x}$ is defined for \mathbf{x} in \mathbb{R}^n , and
- ▶ the vector $\mathbf{b} = A\mathbf{x}$ is a vector in \mathbb{R}^m .

So we can think of A as an **object that acts** on vectors \mathbf{x} in \mathbb{R}^n (via the product $A\mathbf{x}$) to produce vectors \mathbf{b} in \mathbb{R}^m .

Transformation from \mathbb{R}^n to \mathbb{R}^m

Definition: A transformation T (a.k.a. **function** or **mapping**) from \mathbb{R}^n to \mathbb{R}^m is a rule that assigns to each vector \mathbf{x} in \mathbb{R}^n a vector $T(\mathbf{x})$ in \mathbb{R}^m .

Some relevant terms and notation include

- ▶ \mathbb{R}^n is the **domain** and \mathbb{R}^m is called the **codomain**.
- ▶ For \mathbf{x} in the domain, $T(\mathbf{x})$ is called the **image** of \mathbf{x} under T .
- ▶ The collection of all images is called the **range**.
- ▶ The notation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ may be used to indicate that \mathbb{R}^n is the domain and \mathbb{R}^m is the codomain.
- ▶ If $T(\mathbf{x})$ is defined by multiplication by the $m \times n$ matrix A , we may denote this by $\mathbf{x} \mapsto A\mathbf{x}$.

"T of x"

"x maps to Ax"

Matrix Transformation Example

Let $A = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 0 & -2 \end{bmatrix}$. Define the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ by the mapping $T(\mathbf{x}) = A\mathbf{x}$.

(a) Find the image of the vector $\mathbf{u} = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$ under T .

$$\begin{aligned} T(\vec{u}) &= A\vec{u} \\ &= \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 1 \\ -3 \end{bmatrix} = \begin{bmatrix} -8 \\ -10 \\ 6 \end{bmatrix} \end{aligned}$$

Note $\vec{u} = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$
is in \mathbb{R}^2

The output
 $\begin{bmatrix} -8 \\ -10 \\ 6 \end{bmatrix}$ is in \mathbb{R}^3

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 0 & -2 \end{bmatrix}$$

(b) Determine a vector \mathbf{x} in \mathbb{R}^2 whose image under T is $\begin{bmatrix} -4 \\ -4 \\ 4 \end{bmatrix}$.

We want to solve $T(\vec{x}) = \begin{bmatrix} -4 \\ -4 \\ 4 \end{bmatrix}$.

$T(\vec{x}) = A\vec{x}$, so our equation is

$$A\vec{x} = \begin{bmatrix} -4 \\ -4 \\ 4 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -4 \\ -4 \\ 4 \end{bmatrix}$$

Using an augmented matrix

$$\begin{bmatrix} 1 & 3 & -4 \\ 2 & 4 & -4 \\ 0 & -2 & 4 \end{bmatrix} \xrightarrow{\text{rref}} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{bmatrix}$$

$$x_1 = 2, \quad x_2 = -2$$

The vector $\vec{x} = \begin{bmatrix} 2 \\ -2 \end{bmatrix}$ has

image $\begin{bmatrix} -4 \\ -4 \\ 4 \end{bmatrix}$.

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 4 \\ 0 & -2 \end{bmatrix}$$

(c) Determine if $\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$ is in the range of T .

Is $\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$ the image under T of some \vec{x} in \mathbb{R}^2 ? Is $T(\vec{x}) = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$ solvable? This is the same as asking if $A\vec{x} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$ is consistent.

Using an augmented matrix

$$\begin{bmatrix} 1 & 3 & 1 \\ 2 & 4 & 0 \\ 0 & -2 & 1 \end{bmatrix} \xrightarrow{\text{rref}} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

↑
pivot
column

$A\vec{x} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$ is inconsistent.

$\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$ is not in the range of T .

Linear Transformations

Definition: A transformation T is **linear** provided

- (i) $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for every \mathbf{u}, \mathbf{v} in the domain of T , and
- (ii) $T(c\mathbf{u}) = cT(\mathbf{u})$ for every scalar c and vector \mathbf{u} in the domain of T .

Every matrix transformation (e.g. $\mathbf{x} \mapsto A\mathbf{x}$) is a linear transformation. And it turns out that every linear transformation from \mathbb{R}^n to \mathbb{R}^m can be expressed in terms of matrix multiplication.

A Theorem About Linear Transformations:

If T is a linear transformation, then

$$T(\mathbf{0}) = \mathbf{0},$$

$$T(c\mathbf{u} + d\mathbf{v}) = cT(\mathbf{u}) + dT(\mathbf{v})$$

for scalars c, d and vectors \mathbf{u}, \mathbf{v} .

And in fact

$$T(c_1\mathbf{u}_1 + c_2\mathbf{u}_2 + \cdots + c_k\mathbf{u}_k) = c_1T(\mathbf{u}_1) + c_2T(\mathbf{u}_2) + \cdots + c_kT(\mathbf{u}_k).$$

Example

Let r be a nonzero scalar. The transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\mathbf{x}) = r\mathbf{x}$$

is a linear transformation¹.

Show that T is a linear transformation.

We need to show that for each \vec{u}, \vec{v} in \mathbb{R}^2 and scalar c ,

$$T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v}) \quad \text{and}$$

$$T(c\vec{u}) = cT(\vec{u}).$$

$$T(\vec{u} + \vec{v}) = r(\vec{u} + \vec{v}) = r\vec{u} + r\vec{v} = T(\vec{u}) + T(\vec{v})$$

¹It's called a **contraction** if $0 < r < 1$ and a **dilation** when $r \geq 1$

The 1st property holds. Similarly

$$\begin{aligned} T(c\vec{u}) &= r(c\vec{u}) = rc\vec{u} = cr\vec{u} = c(r\vec{u}) \\ &= cT(\vec{u}). \end{aligned}$$

Hence both properties hold. T is
a linear transformation.




Figure: Geometry of dilation $\mathbf{x} \mapsto 2\mathbf{x}$. The 4 by 4 square maps to an 8 by 8 square.

Section 1.9: The Matrix for a Linear Transformation

Elementary Vectors: We'll use the notation \mathbf{e}_i to denote the vector in \mathbb{R}^n having a 1 in the i^{th} position and zero everywhere else.

e.g. in \mathbb{R}^2 the elementary vectors are

$$\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix},$$

in \mathbb{R}^3 they would be

$$\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{e}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

and so forth.

Note that in \mathbb{R}^n , the elementary vectors are the columns of the identity I_n .

Matrix of Linear Transformation

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^4$ be a linear transformation, and suppose

$$T(\mathbf{e}_1) = \begin{bmatrix} 0 \\ 1 \\ -2 \\ 4 \end{bmatrix}, \quad \text{and} \quad T(\mathbf{e}_2) = \begin{bmatrix} 1 \\ 1 \\ -1 \\ 6 \end{bmatrix}.$$

Use the fact that T is linear, and the fact that for each \mathbf{x} in \mathbb{R}^2 we have

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2$$

to find a matrix A such that

$$T(\mathbf{x}) = A\mathbf{x} \quad \text{for every} \quad \mathbf{x} \in \mathbb{R}^2.$$

$$T(\mathbf{e}_1) = \begin{bmatrix} 0 \\ 1 \\ -2 \\ 4 \end{bmatrix}, \quad \text{and} \quad T(\mathbf{e}_2) = \begin{bmatrix} 1 \\ 1 \\ -1 \\ 6 \end{bmatrix}$$

We know that for any \vec{x} in \mathbb{R}^2 , $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$

$\vec{x} = x_1 \vec{e}_1 + x_2 \vec{e}_2$. Then

$$T(\vec{x}) = T(x_1 \vec{e}_1 + x_2 \vec{e}_2)$$

$$= x_1 T(\vec{e}_1) + x_2 T(\vec{e}_2)$$

$$= x_1 \begin{bmatrix} 0 \\ 1 \\ -2 \\ 4 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 1 \\ -1 \\ 6 \end{bmatrix}$$

Since T is a
lin. Transformation

given

$$= \begin{bmatrix} 0 & 1 \\ 1 & 1 \\ -2 & -1 \\ 4 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

By definition of
matrix product

$$= A\vec{x} \quad \text{if} \quad A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \\ -2 & -1 \\ 4 & 6 \end{bmatrix}$$

This holds for all \vec{x} in \mathbb{R}^2 . That is

$$T(\vec{x}) = A\vec{x} \quad \text{for this matrix } A.$$

Theorem

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. There exists a unique $m \times n$ matrix A such that

$$T(\mathbf{x}) = A\mathbf{x} \quad \text{for every } \mathbf{x} \in \mathbb{R}^n.$$

Moreover, the j^{th} column of the matrix A is the vector $T(\mathbf{e}_j)$, where \mathbf{e}_j is the j^{th} column of the $n \times n$ identity matrix I_n . That is,

$$A = [T(\mathbf{e}_1) \quad T(\mathbf{e}_2) \quad \cdots \quad T(\mathbf{e}_n)].$$

The matrix A is called the **standard matrix** for the linear transformation T .

Example

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the scaling transformation (contraction or dilation for $r > 0$) defined by

$$T(\mathbf{x}) = r\mathbf{x}, \quad \text{for positive scalar } r.$$


Find the standard matrix for T .

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad \text{so}$$
$$T(\vec{e}_1) = r \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} r \\ 0 \end{bmatrix}, \quad \text{and} \quad T(\vec{e}_2) = r \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ r \end{bmatrix}.$$

$$A = [T(\vec{e}_1) \quad T(\vec{e}_2)] = \begin{bmatrix} r & 0 \\ 0 & r \end{bmatrix}.$$

Example

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the rotation transformation that rotates each point in \mathbb{R}^2 counter clockwise about the origin through an angle ϕ . Find the standard matrix for T .



point $(\cos(\frac{\pi}{2} + \phi), \sin(\frac{\pi}{2} + \phi))$

point $(\cos \phi, \sin \phi)$

$$T(\vec{e}_1) = \begin{bmatrix} \cos \phi \\ \sin \phi \end{bmatrix}$$

$$T(\vec{e}_2) = \begin{bmatrix} \cos(\frac{\pi}{2} + \phi) \\ \sin(\frac{\pi}{2} + \phi) \end{bmatrix}$$

$$\begin{aligned}\text{Nota } \cos\left(\frac{\pi}{2} + \varphi\right) &= \cos\frac{\pi}{2} \cos \varphi - \sin\frac{\pi}{2} \sin \varphi \\ &= -\sin \varphi\end{aligned}$$

$$\begin{aligned}\sin\left(\frac{\pi}{2} + \varphi\right) &= \sin\frac{\pi}{2} \cos \varphi + \cos\frac{\pi}{2} \sin \varphi \\ &= \cos \varphi\end{aligned}$$

$$\text{so } T(\vec{e}_2) = \begin{bmatrix} -\sin \varphi \\ \cos \varphi \end{bmatrix}$$

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix} = \begin{bmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{bmatrix}$$