

January 12 Math 2306 sec 58 Spring 2016

Section 1: Concepts and Terminology

Suppose $y = \phi(x)$ is a differentiable function. We know that $dy/dx = \phi'(x)$ is another (related) function.

For example, if $y = \cos(2x)$, then y is differentiable on $(-\infty, \infty)$. In fact,

$$\frac{dy}{dx} = -2 \sin(2x).$$

Even dy/dx is differentiable with $d^2y/dx^2 = -4 \cos(2x)$. Note that

$$\frac{d^2y}{dx^2} + 4y = 0.$$

The equation

$$\frac{d^2y}{dx^2} + 4y = 0.$$

is an example of a **differential equation**.

Questions: If we only started with the equation, how could we determine that $\cos(2x)$ satisfies it? Also, is $\cos(2x)$ the only possible function that y could be?

Definition

A **Differential Equation** is an equation containing the derivative(s) of one or more dependent variables, with respect to one or more independent variables.

Independent Variable: will appear as one that derivatives are taken with respect to.

Dependent Variable: will appear as one that derivatives are taken of.

y is dependent ↘

$$\frac{dy}{dx}$$

↑ *x is independent*

u is dependent ↘

$$\frac{du}{dt}$$

↑ *t is independent*

x is dependent ↘

$$\frac{dx}{dr}$$

↑ *r is independent*

Classifications

Type: An **ordinary differential equation (ODE)** has exactly one independent variable¹. For example

$$\frac{dy}{dx} - y^2 = 3x, \quad \text{or} \quad \frac{dy}{dt} + 2\frac{dx}{dt} = t, \quad \text{or} \quad y'' + 4y = 0$$

A **partial differential equation (PDE)** has two or more independent variables. For example

$$\frac{\partial y}{\partial t} = \frac{\partial^2 y}{\partial x^2}, \quad \text{or} \quad \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$$

∂ -partial symbol

¹These are the subject of this course.

Classifications

Order: The order of a differential equation is the same as the highest order derivative appearing anywhere in the equation.

$$\frac{dy}{dx} - y^2 = 3x \quad 1^{\text{st}} \text{ order eqn.}$$

$$y''' + (y')^4 = x^3 \quad 3^{\text{rd}} \text{ order eqn.}$$

$$\frac{\partial y}{\partial t} = \frac{\partial^2 y}{\partial x^2} \quad 2^{\text{nd}} \text{ order eqn.}$$

Notations and Symbols

We'll use standard derivative notations:

$$\text{Leibniz: } \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^ny}{dx^n}, \text{ or}$$

$$\text{Prime \& superscripts: } y', y'', \dots, y^{(n)}.$$

Newton's **dot notation** may be used if the independent variable is time. For example if s is a position function, then

$$\text{velocity is } \frac{ds}{dt} = \dot{s}, \quad \text{and acceleration is } \frac{d^2s}{dt^2} = \ddot{s}$$

Notations and Symbols

An n^{th} order ODE, with independent variable x and dependent variable y can always be expressed as an equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

where F is some real valued function of $n + 2$ variables.

Normal Form: If it is possible to isolate the highest derivative term, then we can write a **normal form** of the equation

$$\frac{d^n y}{dx^n} = f(x, y, y', \dots, y^{(n-1)}).$$

Example : $y''' + (y')^4 = x^3$

$\Rightarrow y''' + (y')^4 - x^3 = 0$
this is $F(x, y, y', y'', y''')$

Normal form for this is

$y''' = -(y')^4 + x^3$ here $f(x, y, y', y'') = -(y')^4 + x^3$

Notations and Symbols

If $n = 1$, an equation in normal form would look like

$$\frac{dy}{dx} = f(x, y).$$

If $n = 2$, an equation in normal form would look like

$$\frac{d^2y}{dx^2} = f(x, y, y').$$

Differential Form

A first order equation may appear in the form

$$M(x, y) dx + N(x, y) dy = 0$$

We can write 2 different Normal forms:

$$N(x, y) dy = -M(x, y) dx$$

OR

$$M(x, y) dx = -N(x, y) dy$$

$$\frac{dy}{dx} = \frac{-M(x, y)}{N(x, y)}$$

if $N(x, y) \neq 0$

$$\frac{dx}{dy} = \frac{-N(x, y)}{M(x, y)}$$

if $M(x, y) \neq 0$

Classifications

Linearity: An n^{th} order differential equation is said to be **linear** if it can be written in the form

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x).$$

Note that each of the coefficients a_0, \dots, a_n and the right hand side g may depend on the independent variable but not on the dependent variable or any of its derivatives.

Examples (Linear -vs- Nonlinear)

$$y'' + 4y = 0 \quad \text{Linear}$$

$$a_2(x)y'' + a_1(x)y' + a_0(x)y = f(x)$$

$$a_2(x) = 1 \quad a_0(x) = 4$$

$$a_1(x) = 0 \quad f(x) = 0$$

$$t^2 \frac{d^2x}{dt^2} + 2t \frac{dx}{dt} - x = e^t$$

$$a_2(t) = t^2 \quad a_0(t) = -1$$

$$a_1(t) = 2t \quad f(t) = e^t$$

$$\frac{d^3y}{dx^3} + \left(\frac{dy}{dx}\right)^4 = x^3$$

non linear

$$u'' + u' = \cos u$$

$$\frac{d^3y}{dx^3} + \left(\frac{dy}{dx}\right)^3 \frac{dy}{dx} = x^3$$

non linear term

non linear term

Examples

Identify the independent and dependent variables. Determine the order of the equation. State whether it is linear or nonlinear.

(a) $y'' + 2ty' = \cos t + y$

Independent - t
dependent - y
2nd order

$$y'' + 2ty' - y = \cos t$$

this is linear

Examples

$$(b) \frac{d^3y}{dx^3} + 2y \frac{dy}{dx} = \frac{d^2y}{dx^2} + \tan(x)$$

Independent x

dependent y

3rd order

$$\frac{d^3y}{dx^3} - \frac{d^2y}{dx^2} + 2y \frac{dy}{dx} = \tan x$$

non linear term

The eqn. is non linear.

Examples

(c) $\ddot{\theta} + \frac{g}{l} \sin \theta = 0$ g and l are constant

Independent is t
(note the dot)

dependent θ

2nd order


Not linear eqn.

$\sin \theta$ is the non linear term.

Solution of $F(x, y, y', \dots, y^{(n)}) = 0$ (*)

Definition: A function ϕ defined on an interval I^2 and possessing at least n continuous derivatives on I is a **solution** of (*) on I if upon substitution (i.e. setting $y = \phi(x)$) the equation reduces to an identity.

Definition: An **implicit solution** of (*) is a relation $G(x, y) = 0$ provided there exists at least one function $y = \phi$ that satisfies both the differential equation (*) and this relation.

²The interval is called the *domain of the solution* or the *interval of definition*. 

Examples:

Verify that the given function is an solution of the ODE on the indicated interval.

$$\phi(t) = 3e^{2t}, \quad I = (-\infty, \infty), \quad \frac{d^2y}{dt^2} - \frac{dy}{dt} - 2y = 0$$

$\phi(t) = 3e^{2t}$ has derivatives of all orders on \mathbb{I}

$$\text{Set } y = 3e^{2t} \quad \text{and} \quad y'' = 12e^{2t}$$
$$y' = 6e^{2t}$$