

Section 1.1: Limits of Functions Using Numerical and Graphical Techniques



Recall: For a line $y = mx + b$, the slope tells us how a change in x (Δx) causes a change in y (Δy). In fact, the slope $m = \frac{\Delta y}{\Delta x}$.

For a non-line curve, $y = f(x)$ we wanted to define *slope*. We still want slope to say something about how y changes if x changes. But we don't expect slope to be the same number for every x .

We want to define the slope of a **tangent line**.

Slope of the Tangent Line

We consider a sequence of points $Q_1 = (x_1, f(x_1))$, $Q_2 = (x_2, f(x_2))$, and so forth in such a way that the x -values are getting closer to c . Note that the resulting secant lines tend to have slopes closer to that of the tangent line.



Slope of the Tangent Line

$$m_{\text{sec}} = \frac{f(x) - f(c)}{x - c}$$

$$m_{\text{sec}} \quad m_{\text{tan}}$$

This is a limit process. We can write this as

$$m_{\text{tan}} = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$$

(read “the limit as x approaches c of ...”).

This required us to define what a limit is.

A Working Definition of a Limit

Definition: Let f be defined on an open interval containing the number c except possibly at c . Then

$$\lim_{x \rightarrow c} f(x) = L$$

provided the value of $f(x)$ can be made arbitrarily close to the number L by taking x sufficiently close to c but not equal to c .

We considered the example (using a calculator)

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x}$$

$$\lim_{x \rightarrow c} f(x) = L$$

here $c = 0$
and
 $L = 1$

x	$f(x) = \frac{e^x - 1}{x}$
-0.1	0.9516
-0.01	0.9950
-0.001	0.9995
0	undefined
0.001	1.0005
0.01	1.0050
0.1	1.0517

x close to c but less than c

x close to c but greater than c

From the values, we conclude that $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$.

Left and Right Hand Limits

In our examples, we considered x -values to the left (less than) and to the right (greater than) c . This illustrates the notion of **one sided limits**. We have a special notation for this.

Left Hand Limit: We write

$$\lim_{x \rightarrow c^-} f(x) = L_L$$

and say *the limit as x approaches c from the left of $f(x)$ equals L_L provided we can make $f(x)$ arbitrarily close to the number L_L by taking x sufficiently close to, but less than c .*

Left and Right Hand Limits

Right Hand Limit: We write

$$\lim_{x \rightarrow c^+} f(x) = L_R$$

and say *the limit as x approaches c from the right of $f(x)$ equals L_R provided we can make $f(x)$ arbitrarily close to the number L_R by taking x sufficiently close to, but greater than c .*

Some other common phrases:

”from the left” is the same as ”from below”

”from the right” is the same as ”from above.”


Example

Plot the function $f(x) = \begin{cases} x^2, & x < 1 \\ 2, & x = 1 \\ 1, & x > 1 \end{cases}$

graph.

Investigate $\lim_{x \rightarrow 1} f(x)$ using the

$c=1$



looks like $\lim_{x \rightarrow 1^-} f(x) = 1$

looks like $\lim_{x \rightarrow 1^+} f(x) = 1$

X	f(X)
0.9	$(0.9)^2 = 0.81$
0.99	$(0.99)^2 = 0.9801$
0.999	$(0.999)^2 = 0.998001$
1	2
1.001	1
1.01	1
1.1	1

In fact

$$\lim_{x \rightarrow 1} f(x) = 1$$

Observations

Observation 1: The limit L of a function $f(x)$ as x approaches c does not depend on whether $f(c)$ exists or what its value may be.

Observation 2: If $\lim_{x \rightarrow c} f(x) = L$, then the number L is unique. That is, a function can not have two different limits as x approaches a single number c .

Observation 3: A function need not have a limit as x approaches c . If $f(x)$ can not be made arbitrarily close to any one number L as x approaches c , then we say that $\lim_{x \rightarrow c} f(x)$ **does not exist** (shorthand **DNE**).

Questions

(1) **True** or **False** It is possible that both $\lim_{x \rightarrow 3} f(x) = 5$ AND $f(3) = 7$.

f(c) doesn't have to affect the limit.

(2) **True** or **False** It is possible that both $\lim_{x \rightarrow 3} f(x) = 5$ AND $\lim_{x \rightarrow 3} f(x) = 7$.

Limits are unique.