

## Section 1.1: Limits of Functions Using Numerical and Graphical Techniques

**Definition:** Let  $f$  be defined on an open interval containing the number  $c$  except possibly at  $c$ . Then

$$\lim_{x \rightarrow c} f(x) = L$$

provided the value of  $f(x)$  can be made arbitrarily close to the number  $L$  by taking  $x$  sufficiently close to  $c$  but not equal to  $c$ .

# One Sided Limits

**Left Hand Limit:** We write

$$\lim_{x \rightarrow c^-} f(x) = L_L$$

and say *the limit as  $x$  approaches  $c$  from the left of  $f(x)$  equals  $L_L$  provided we can make  $f(x)$  arbitrarily close to the number  $L_L$  by taking  $x$  sufficiently close to, but less than  $c$ .*

**Right Hand Limit:** We write

$$\lim_{x \rightarrow c^+} f(x) = L_R$$

and say *the limit as  $x$  approaches  $c$  from the right of  $f(x)$  equals  $L_R$  provided we can make  $f(x)$  arbitrarily close to the number  $L_R$  by taking  $x$  sufficiently close to, but greater than  $c$ .*

## Observations


**Observation 1:** The limit  $L$  of a function  $f(x)$  as  $x$  approaches  $c$  does not depend on whether  $f(c)$  exists or what its value may be.

**Observation 2:** If  $\lim_{x \rightarrow c} f(x) = L$ , then the number  $L$  is unique. That is, a function can not have two different limits as  $x$  approaches a single number  $c$ .

**Observation 3:** A function need not have a limit as  $x$  approaches  $c$ . If  $f(x)$  can not be made arbitrarily close to any one number  $L$  as  $x$  approaches  $c$ , then we say that  $\lim_{x \rightarrow c} f(x)$  **does not exist** (shorthand **DNE**).

## A Limit Failing to Exist

$$\text{Consider } H(x) = \begin{cases} 0, & x < 0 \\ 1, & x \geq 0 \end{cases}.$$



We determined using a graph that

$$\lim_{x \rightarrow 0^-} H(x) = 0, \quad \text{and} \quad \lim_{x \rightarrow 0^+} H(x) = 1.$$

Since there is no unique number  $L$  such that  $H(x)$  gets *arbitrarily close* to  $L$  when  $x$  is sufficiently close to zero, it turns out that

$$\lim_{x \rightarrow 0} H(x) \quad \text{Does not exist.}$$

## Weakness of Technology

Suppose we wish to investigate

$$\lim_{x \rightarrow 0} \sin\left(\frac{\pi}{x^2}\right).$$

We consider values of  $x$  closer to zero, and plug them into a calculator. Let's look at two attempts.

$x$	$\sin\left(\frac{\pi}{x^2}\right)$
-0.1	0
-0.01	0
-0.001	0
0	undefined
0.001	0
0.01	0
0.1	0

*Limit  
looks  
like  
zero*

$x$	$\sin\left(\frac{\pi}{x^2}\right)$
$-\frac{2}{3}$	0.707
$-\frac{2}{13}$	0.707
$-\frac{2}{23}$	0.707
0	undefined
$\frac{2}{23}$	0.707
$\frac{2}{13}$	0.707
$\frac{2}{3}$	0.707

*Limit  
looks  
like  
0.707*

## Weakness of Technology

In every interval containing zero, the graph of  $\sin(\pi/x^2)$  passes through every  $y$ -value between  $-1$  and  $1$  infinitely many times.




Figure:  $y = \sin\left(\frac{\pi}{x^2}\right)$

## Evaluating Limits

As this example illustrates, we would like to avoid too much reliance on technology for evaluating limits. The next section will be devoted to techniques for doing this for reasonably well behaved functions. We close with one theorem.


**Theorem:** Let  $f$  be defined on an open interval containing  $c$  except possibly at  $c$ . Then

$$\lim_{x \rightarrow c} f(x) = L$$

if and only if

$$\lim_{x \rightarrow c^-} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow c^+} f(x) = L.$$


## Limits from a graph



Evaluate  $\lim_{x \rightarrow 1^-} f(x)$




# Limits from a graph



$$\lim_{x \rightarrow 1^-} f(x) = 4$$

from  
the graph

# Question



aside  
lim f(x)  
x → 1 DNE

$$\lim_{x \rightarrow 1^+} f(x) =$$

(a) 4

(b) -2

(c) DNE

(d) 1

## Section 1.2: Limits of Functions Using Properties of Limits

We begin with two of the simplest limits we may encounter.


**Theorem:** If  $f(x) = A$  where  $A$  is a constant, then for any real number  $c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} A = A$$

*limit of a constant is that constant*

**Theorem:** If  $f(x) = x$ , then for any real number  $c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x = c$$



# Examples

(a)  $\lim_{x \rightarrow 0} 7 = 7$       limit of a constant

(b)  $\lim_{x \rightarrow \pi^+} 3\pi = 3\pi$

\* we know  $\lim_{x \rightarrow \pi} 3\pi = 3\pi$

(c)  $\lim_{x \rightarrow -\sqrt{5}} x = -\sqrt{5}$

since  $\lim_{x \rightarrow c} x = c$

## Question

$$\lim_{x \rightarrow 4^-} x =$$

(a)  $x$

(b)  $-4$

(c)  $4$

Since  $\lim_{x \rightarrow 4} x = 4$

(d) the one sided limit can't be determined

## Additional Limit Law Theorems

Suppose

$$\lim_{x \rightarrow c} f(x) = L, \quad \lim_{x \rightarrow c} g(x) = M, \quad \text{and } k \text{ is constant.}$$

**Theorem: (Sums)**  $\lim_{x \rightarrow c} (f(x) + g(x)) = L + M$

**Theorem: (Differences)**  $\lim_{x \rightarrow c} (f(x) - g(x)) = L - M$

**Theorem: (Constant Multiples)**  $\lim_{x \rightarrow c} kf(x) = kL$

**Theorem: (Products)**  $\lim_{x \rightarrow c} f(x)g(x) = LM$

## Examples

Use the limit law theorems to evaluate if possible

$$\begin{aligned} \text{(a)} \quad \lim_{x \rightarrow 2} (3x+2) &= \lim_{x \rightarrow 2} 3x + \lim_{x \rightarrow 2} 2 && \text{Sum} \\ &= 3 \lim_{x \rightarrow 2} x + \lim_{x \rightarrow 2} 2 && \text{Constant} \\ & && \text{multiply} \\ &= 3(2) + 2 \\ &= 6 + 2 = 8 \end{aligned}$$

## Examples

Use the limit law theorems to evaluate if possible

$$(b) \lim_{x \rightarrow -3} (x+1)^2$$

Consider  $\lim_{x \rightarrow -3} (x+1) = \lim_{x \rightarrow -3} x + \lim_{x \rightarrow -3} 1$

$$= -3 + 1 = -2$$

S.  $\lim_{x \rightarrow -3} (x+1)^2 = \left( \lim_{x \rightarrow -3} (x+1) \right) \cdot \left( \lim_{x \rightarrow -3} (x+1) \right)$  product property

$$= (-2) \cdot (-2) = 4$$



## Examples

Use the limit law theorems to evaluate if possible

$$(c) \quad \lim_{x \rightarrow 0} f(x) \quad \text{where} \quad f(x) = \begin{cases} x + 2, & x < 0 \\ 1, & x = 0 \\ 2x - 3, & x > 0 \end{cases}$$

we know  $\lim_{x \rightarrow 0} f(x) = L$  if and only if

$$\lim_{x \rightarrow 0^-} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow 0^+} f(x) = L$$

$$\begin{aligned} \lim_{x \rightarrow 0^-} f(x) &= \lim_{x \rightarrow 0^-} (x + 2) = \lim_{x \rightarrow 0^-} x + \lim_{x \rightarrow 0^-} 2 \\ &= 0 + 2 = 2 \end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow 0^+} f(x) &= \lim_{x \rightarrow 0^+} (2x - 3) = \lim_{x \rightarrow 0^+} 2x - \lim_{x \rightarrow 0^+} 3 \\ &= 2 \lim_{x \rightarrow 0^+} x - \lim_{x \rightarrow 0^+} 3 \\ &= 2 \cdot 0 - 3 = -3\end{aligned}$$

We have

$$\lim_{x \rightarrow 0^-} f(x) = 2 \quad \text{and} \quad \lim_{x \rightarrow 0^+} f(x) = -3$$

$$\lim_{x \rightarrow 0} f(x) \quad \text{DNE}$$

## Question

(1)  $\lim_{x \rightarrow 1} f(x)$  where  $f(x) = \begin{cases} x^2 + 1, & x \leq 1 \\ 3 - x, & x > 1 \end{cases}$

(a) 4

It is the case that

(b) 2

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x^2 + 1) = 2$$

(c) 1

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (3 - x) = 2$$

(d) DNE

## Additional Limit Law Theorems

Suppose  $\lim_{x \rightarrow c} f(x) = L$  and  $n$  is a positive integer.

**Theorem: (Power)**  $\lim_{x \rightarrow c} (f(x))^n = L^n$

Note in particular that this tells us that  $\lim_{x \rightarrow c} x^n = c^n$ .

**Theorem: (Root)**  $\lim_{x \rightarrow c} \sqrt[n]{f(x)} = \sqrt[n]{L}$  (if this is defined)

Combining the sum, difference, constant multiple and power laws:

**Theorem:** If  $P(x)$  is a polynomial, then

$$\lim_{x \rightarrow c} P(x) = P(c).$$

## Question

Hint: let  $P(x) = \underline{3x^2 - 4x + 7}$

(1)  $\lim_{x \rightarrow 2} (3x^2 - 4x + 7) =$

(a) 7

(b) DNE

(c) -11

(d) 11

$$\begin{aligned} P(2) &= 3(2)^2 - 4 \cdot 2 + 7 \\ &= 12 - 8 + 7 = 11 \end{aligned}$$

## Notation Reminder

The notation "lim" is **always** followed by a function expression and never immediately by an equal sign.

$\lim_{x \rightarrow c}$

## Question

(2) Suppose that we have determined that  $\lim_{x \rightarrow 7} f(x) = 13$ .

**True or False:** It is acceptable to write this as

$$\text{" } \lim_{x \rightarrow 7} = 13 \text{"}$$

This is like writing " $\sqrt{\quad} = 4$ "

## Additional Limit Law Theorems

Suppose  $\lim_{x \rightarrow c} f(x) = L$ ,  $\lim_{x \rightarrow c} g(x) = M$  and  $M \neq 0$

**Theorem: (Quotient)**  $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{L}{M}$

Combined with our result for polynomials:

**Theorem:** If  $R(x) = \frac{p(x)}{q(x)}$  is a rational function, and  $c$  is in the domain of  $R$ , then

$$\lim_{x \rightarrow c} R(x) = R(c).$$



## Examples

Evaluate  $\lim_{x \rightarrow 2} \frac{x^2 + 5}{x^2 + x - 1}$

$R(x) = \frac{x^2 + 5}{x^2 + x - 1}$  is rational

is 2 in the domain of  $R$ ?

$2^2 + 2 - 1 = 4 + 2 - 1 = 5 \neq 0$   
Yes, 2 is in the domain!

So  $\lim_{x \rightarrow 2} \frac{x^2 + 5}{x^2 + x - 1} = \frac{2^2 + 5}{2^2 + 2 - 1} = \frac{9}{5}$

# Examples

Evaluate  $\lim_{x \rightarrow 1} \frac{\sqrt{x+1}}{x+5}$

Note  $\lim_{x \rightarrow 1} (x+5) = 1+5 = 6$  ( $x+5$  is a polynomial)

also  $\lim_{x \rightarrow 1} (x+1) = 1+1 = 2$  so  $\lim_{x \rightarrow 1} \sqrt{x+1} = \sqrt{2}$

Hence  $\lim_{x \rightarrow 1} \frac{\sqrt{x+1}}{x+5} = \frac{\sqrt{2}}{6}$

## Additional Techniques: When direct laws fail

Evaluate if possible  $\lim_{x \rightarrow 2} \frac{x^2 - x - 2}{x^2 - 4}$

Let  $R(x) = \frac{x^2 - x - 2}{x^2 - 4}$  which is rational

Unfortunately, 2 is not in the domain since

$$2^2 - 4 = 0.$$

Note:  $2^2 - 2 - 2 = 0$

Since 2 is a root of both polynomials,  $x - 2$  is a factor!

This motivates trying to cancel this common factor.

$$\lim_{x \rightarrow 2} \frac{x^2 - x - 2}{x^2 - 4} = \lim_{x \rightarrow 2} \frac{(x-2)(x+1)}{(x-2)(x+2)}$$

then cancel  
common  
factor (5)

$$= \lim_{x \rightarrow 2} \frac{x+1}{x+2}$$

$$= \frac{2+1}{2+2} = \frac{3}{4}$$

## Additional Techniques: When direct laws fail

Evaluate if possible  $\lim_{x \rightarrow 1} \frac{\sqrt{x+3} - 2}{x-1}$

Note  $\lim_{x \rightarrow 1} (x-1) = 0$  and  $\lim_{x \rightarrow 1} (\sqrt{x+3} - 2) = \sqrt{1+3} - 2 = 0$

Since the top is zero @  $x=1$ , we can think of  $x-1$  as a "factor" in some sense.

we'll coax it out by rationalizing.

we'll use the conjugate  $\sqrt{x+3} + 2$

$$\lim_{x \rightarrow 1} \frac{\sqrt{x+3} - 2}{x-1} = \lim_{x \rightarrow 1} \left( \frac{\sqrt{x+3} - 2}{x-1} \right) \cdot \left( \frac{\sqrt{x+3} + 2}{\sqrt{x+3} + 2} \right)$$

↓ mult. by 1

$$= \lim_{x \rightarrow 1} \frac{(\sqrt{x+3})^2 - 2\sqrt{x+3} + 2\sqrt{x+3} - 4}{(x-1)(\sqrt{x+3} + 2)}$$

$$= \lim_{x \rightarrow 1} \frac{x+3-4}{(x-1)(\sqrt{x+3} + 2)}$$

$$= \lim_{x \rightarrow 1} \frac{\cancel{x-1}}{(\cancel{x-1})(\sqrt{x+3} + 2)} = \lim_{x \rightarrow 1} \frac{1}{\sqrt{x+3} + 2} = \frac{1}{\sqrt{1+3} + 2} = \frac{1}{4}$$

## Question

Evaluate if possible  $\lim_{x \rightarrow 2} \frac{x-2}{\sqrt{x}-\sqrt{2}} = \lim_{x \rightarrow 2} \left( \frac{x-2}{\sqrt{x}-\sqrt{2}} \right) \cdot \left( \frac{\sqrt{x}+\sqrt{2}}{\sqrt{x}+\sqrt{2}} \right)$

(a)  $\frac{1}{\sqrt{2}}$

$$= \lim_{x \rightarrow 2} \frac{(x-2)(\sqrt{x}+\sqrt{2})}{x-2}$$

(b)  $\sqrt{2}$

(c) DNE

$$= \lim_{x \rightarrow 2} (\sqrt{x}+\sqrt{2}) = \sqrt{2} + \sqrt{2} = 2\sqrt{2}$$

(d)  $2\sqrt{2}$

## Observations

In limit taking, the form " $\frac{0}{0}$ " sometimes appears. This is called an **indeterminate form**. Standard strategies are

- (1) Try to factor the numerator and denominator to see if a common factor  $-(x - c)$  can be cancelled.
- (2) If dealing with roots, try rationalizing to reveal a common factor.

The form

$$\frac{\text{„ nonzero constant „}}{0}$$

is not indeterminate. It is undefined. When it appears, the limit doesn't exist.