

## Section 1.1: Limits of Functions Using Numerical and Graphical Techniques

**Definition:** Let  $f$  be defined on an open interval containing the number  $c$  except possibly at  $c$ . Then

$$\lim_{x \rightarrow c} f(x) = L$$

provided the value of  $f(x)$  can be made arbitrarily close to the number  $L$  by taking  $x$  sufficiently close to  $c$  but not equal to  $c$ .

# One Sided Limits

**Left Hand Limit:** We write

$$\lim_{x \rightarrow c^-} f(x) = L_L$$

and say *the limit as  $x$  approaches  $c$  from the left of  $f(x)$  equals  $L_L$  provided we can make  $f(x)$  arbitrarily close to the number  $L_L$  by taking  $x$  sufficiently close to, but less than  $c$ .*

**Right Hand Limit:** We write

$$\lim_{x \rightarrow c^+} f(x) = L_R$$

and say *the limit as  $x$  approaches  $c$  from the right of  $f(x)$  equals  $L_R$  provided we can make  $f(x)$  arbitrarily close to the number  $L_R$  by taking  $x$  sufficiently close to, but greater than  $c$ .*

## Observations

**Observation 1:** The limit  $L$  of a function  $f(x)$  as  $x$  approaches  $c$  does not depend on whether  $f(c)$  exists or what its value may be.

**Observation 2:** If  $\lim_{x \rightarrow c} f(x) = L$ , then the number  $L$  is unique. That is, a function can not have two different limits as  $x$  approaches a single number  $c$ .


**Observation 3:** A function need not have a limit as  $x$  approaches  $c$ . If  $f(x)$  can not be made arbitrarily close to any one number  $L$  as  $x$  approaches  $c$ , then we say that  $\lim_{x \rightarrow c} f(x)$  **does not exist** (shorthand **DNE**).

## A Limit Failing to Exist

*H is the Heaviside step function*

Consider  $H(x) = \begin{cases} 0, & x < 0 \\ 1, & x \geq 0 \end{cases}$ . Evaluate if possible

$$\lim_{x \rightarrow 0^-} H(x), \quad \lim_{x \rightarrow 0^+} H(x), \quad \text{and} \quad \lim_{x \rightarrow 0} H(x)$$



$$\lim_{x \rightarrow 0^-} H(x) = 0 \quad \text{from the graph}$$

$$\lim_{x \rightarrow 0^+} H(x) = 1 \quad \text{from the graph}$$

$\lim_{x \rightarrow 0} H(x)$  Does not exist (DNE)

There is no one number  $L$  that  $H(x)$  values get arbitrarily close to.

This is called a jump where

$$\lim_{x \rightarrow c^-} f(x) = L_L \quad \lim_{x \rightarrow c^+} f(x) = L_R$$

and  $L_L \neq L_R$

## Weakness of Technology

Suppose we wish to investigate

$$\lim_{x \rightarrow 0} \sin\left(\frac{\pi}{x^2}\right).$$

We consider values of  $x$  closer to zero, and plug them into a calculator. Let's look at two attempts.

$x$	$\sin\left(\frac{\pi}{x^2}\right)$
-0.1	0
-0.01	0
-0.001	0
0	undefined
0.001	0
0.01	0
0.1	0


*limit  
looks  
like zero*

$x$	$\sin\left(\frac{\pi}{x^2}\right)$
$-\frac{2}{3}$	0.707
$-\frac{2}{13}$	0.707
$-\frac{2}{23}$	0.707
0	undefined
$\frac{2}{23}$	0.707
$\frac{2}{13}$	0.707
$\frac{2}{3}$	0.707

*limit  
looks  
like  
0.707...*

## Weakness of Technology

In every interval containing zero, the graph of  $\sin(\pi/x^2)$  passes through every  $y$ -value between  $-1$  and  $1$  infinitely many times.



$\lim_{x \rightarrow 0} \sin\left(\frac{\pi}{x^2}\right) \text{ DNE}$   
"by oscillation"

Figure:  $y = \sin\left(\frac{\pi}{x^2}\right)$

## Evaluating Limits

As this example illustrates, we would like to avoid too much reliance on technology for evaluating limits. The next section will be devoted to techniques for doing this for reasonably well behaved functions. We close with one theorem.

**Theorem:** Let  $f$  be defined on an open interval containing  $c$  except possibly at  $c$ . Then


$$\lim_{x \rightarrow c} f(x) = L$$

if and only if

$$\lim_{x \rightarrow c^-} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow c^+} f(x) = L.$$




## Limits from a graph




Evaluate  $\lim_{x \rightarrow 1^-} f(x)$

## Limits from a graph



# Question



aside:  
 $\lim_{x \rightarrow 1} f(x)$  DNE

$$\lim_{x \rightarrow 1^+} f(x) =$$

(a) 4

(b) -2

(c) DNE

(d) 1

## Section 1.2: Limits of Functions Using Properties of Limits

We begin with two of the simplest limits we may encounter.


**Theorem:** If  $f(x) = A$  where  $A$  is a constant, then for any real number  $c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} A = A$$

*the limit of a constant is that constant*

**Theorem:** If  $f(x) = x$ , then for any real number  $c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x = c$$



# Examples

$$(a) \lim_{x \rightarrow 0} 7 = 7$$

limit of a constant is that constant

$$(b) \lim_{x \rightarrow \pi^+} 3\pi = 3\pi$$

$$\lim_{x \rightarrow c} f(x) = L$$

if and only if

We know that

$$\lim_{x \rightarrow \pi} 3\pi = 3\pi$$

$$\lim_{x \rightarrow c^+} f(x) = L \text{ and } \lim_{x \rightarrow c^-} f(x) = L$$

$$(c) \lim_{x \rightarrow -\sqrt{5}} x = -\sqrt{5}$$

since  $\lim_{x \rightarrow c} x = c$

## Question

$$\lim_{x \rightarrow 4^-} x =$$

$$\lim_{x \rightarrow 4} x = 4$$

(a)  $x$

(b)  $-4$

(c)  $4$

(d) the one sided limit can't be determined

## Additional Limit Law Theorems

Suppose

$$\lim_{x \rightarrow c} f(x) = L, \quad \lim_{x \rightarrow c} g(x) = M, \quad \text{and } k \text{ is constant.}$$

**Theorem: (Sums)**  $\lim_{x \rightarrow c} (f(x) + g(x)) = L + M$

**Theorem: (Differences)**  $\lim_{x \rightarrow c} (f(x) - g(x)) = L - M$

**Theorem: (Constant Multiples)**  $\lim_{x \rightarrow c} kf(x) = kL$

**Theorem: (Products)**  $\lim_{x \rightarrow c} f(x)g(x) = LM$

## Examples

Use the limit law theorems to evaluate if possible

$$(a) \quad \lim_{x \rightarrow 2} (3x+2) : \quad \lim_{x \rightarrow 2} 3x \quad + \quad \lim_{x \rightarrow 2} 2 \quad \text{(Sum)}$$

$$= 3 \lim_{x \rightarrow 2} x \quad + \quad \lim_{x \rightarrow 2} 2 \quad \text{(constant multiple)}$$

$$= 3 \cdot 2 + 2 = 8$$



## Examples

Use the limit law theorems to evaluate if possible

$$(b) \quad \lim_{x \rightarrow -3} (x+1)^2$$

$$\begin{aligned} \text{Consider } \lim_{x \rightarrow -3} (x+1) &= \lim_{x \rightarrow -3} x + \lim_{x \rightarrow -3} 1 \\ &= -3 + 1 = -2 \end{aligned}$$

$$\begin{aligned} \lim_{x \rightarrow -3} (x+1)^2 &= \left( \lim_{x \rightarrow -3} (x+1) \right) \cdot \left( \lim_{x \rightarrow -3} (x+1) \right) \quad (\text{product}) \\ &= (-2) \cdot (-2) = 4 \end{aligned}$$

## Examples

Use the limit law theorems to evaluate if possible

$$(c) \lim_{x \rightarrow 0} f(x) \quad \text{where} \quad f(x) = \begin{cases} x + 2, & x < 0 \\ 1, & x = 0 \\ 2x - 3, & x > 0 \end{cases}$$

$$\lim_{x \rightarrow 0} f(x) = L \quad \text{if and only if} \quad \lim_{x \rightarrow 0^-} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow 0^+} f(x) = L.$$

We can compute the 1-sided limits.

$$\begin{aligned} \lim_{x \rightarrow 0^-} f(x) &= \lim_{x \rightarrow 0^-} (x + 2) = \lim_{x \rightarrow 0^-} x + \lim_{x \rightarrow 0^-} 2 \\ &= 0 + 2 = 2 \end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow 0^+} f(x) &= \lim_{x \rightarrow 0^+} (2x - 3) \\ &= \lim_{x \rightarrow 0^+} 2x - \lim_{x \rightarrow 0^+} 3 \\ &= 2 \lim_{x \rightarrow 0^+} x - \lim_{x \rightarrow 0^+} 3 = 2 \cdot 0 - 3 = -3\end{aligned}$$

$$\lim_{x \rightarrow 0^-} f(x) = 2 \quad \text{and} \quad \lim_{x \rightarrow 0^+} f(x) = -3$$

$$\lim_{x \rightarrow 0} f(x) \text{ DNE}$$

## Question

(1)  $\lim_{x \rightarrow 1} f(x)$  where  $f(x) = \begin{cases} x^2 + 1, & x \leq 1 \\ 3 - x, & x > 1 \end{cases}$

(a) 4

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x^2 + 1) = 2$$

(b) 2

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (3 - x) = 2$$

(c) 1

(d) DNE

## Additional Limit Law Theorems

Suppose  $\lim_{x \rightarrow c} f(x) = L$  and  $n$  is a positive integer.

**Theorem: (Power)**  $\lim_{x \rightarrow c} (f(x))^n = L^n$

Note in particular that this tells us that  $\lim_{x \rightarrow c} x^n = c^n$ .

**Theorem: (Root)**  $\lim_{x \rightarrow c} \sqrt[n]{f(x)} = \sqrt[n]{L}$  (if this is defined)

Combining the sum, difference, constant multiple and power laws:

**Theorem:** If  $P(x)$  is a polynomial, then

$$\lim_{x \rightarrow c} P(x) = P(c).$$

*To take  
the limit,  
plug in  $c$ .*

## Question

$$(1) \lim_{x \rightarrow 2} (3x^2 - 4x + 7) =$$

(a) 7

(b) DNE

(c) -11

(d) 11

Let  $P(x) = 3x^2 - 4x + 7$   
this is a polynomial.

$$\begin{aligned} P(2) &= 3 \cdot 2^2 - 4 \cdot 2 + 7 \\ &= 12 - 8 + 7 = 11 \end{aligned}$$

## Notation Reminder

The notation "lim" is **always** followed by a function expression and never immediately by an equal sign.

$$\lim_{x \rightarrow c}$$

## Question

(2) Suppose that we have determined that  $\lim_{x \rightarrow 7} f(x) = 13$ .

**True or False:** It is acceptable to write this as

$$\text{" } \lim_{x \rightarrow 7} = 13 \text{"}$$

This is similar to writing  $\sqrt{\quad} = 13$



## Additional Limit Law Theorems

Suppose  $\lim_{x \rightarrow c} f(x) = L$ ,  $\lim_{x \rightarrow c} g(x) = M$  and  $M \neq 0$

**Theorem: (Quotient)**  $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{L}{M}$

Combined with our result for polynomials:

**Theorem:** If  $R(x) = \frac{p(x)}{q(x)}$  is a rational function, and  $c$  is in the domain of  $R$ , then

$$\lim_{x \rightarrow c} R(x) = R(c).$$

# Examples

Evaluate  $\lim_{x \rightarrow 2} \frac{x^2 + 5}{x^2 + x - 1}$

$R(x) = \frac{x^2 + 5}{x^2 + x - 1}$  is a rational function. Is 2 in its domain?

$2^2 + 2 - 1 = 4 + 2 - 1 = 5 \neq 0$  so yes, 2 is in its domain.

So  $\lim_{x \rightarrow 2} \frac{x^2 + 5}{x^2 + x - 1} = \frac{2^2 + 5}{2^2 + 2 - 1} = \frac{9}{5}$

# Examples

Evaluate  $\lim_{x \rightarrow 1} \frac{\sqrt{x+1}}{x+5}$

Note  $\lim_{x \rightarrow 1} (x+5) = 1+5 = 6$   *$x+5$  is polynomial*

and  $\lim_{x \rightarrow 1} (x+1) = 1+1 = 2$  so  $\lim_{x \rightarrow 1} \sqrt{x+1} = \sqrt{2}$

so  $\lim_{x \rightarrow 1} \frac{\sqrt{x+1}}{x+5} = \frac{\sqrt{2}}{6}$

## Additional Techniques: When direct laws fail

Evaluate if possible  $\lim_{x \rightarrow 2} \frac{x^2 - x - 2}{x^2 - 4}$

$\frac{x^2 - x - 2}{x^2 - 4}$  is rational, but 2 is not in its domain.

Note  $2^2 - 2 - 2 = 4 - 2 - 2 = 0$

For  $p(x) = x^2 - x - 2$  and  $q(x) = x^2 - 4$ , since  $p(2) = 0$

we know that  $x - 2$  is a factor of  $p$ .

Similarly  $q(2) = 0$  so  $x - 2$  is a factor of  $q$ .

We can try to cancel this common factor.

$$\begin{aligned}\lim_{x \rightarrow 2} \frac{x^2 - x - 2}{x^2 - 4} &= \lim_{x \rightarrow 2} \frac{\cancel{(x-2)}(x+1)}{\cancel{(x-2)}(x+2)} \\ &= \lim_{x \rightarrow 2} \frac{x+1}{x+2} = \frac{2+1}{2+2} = \frac{3}{4}\end{aligned}$$

## Additional Techniques: When direct laws fail

Evaluate if possible  $\lim_{x \rightarrow 1} \frac{\sqrt{x+3} - 2}{x-1}$

$$\lim_{x \rightarrow 1} (x-1) = 0 \quad \text{but also} \quad \lim_{x \rightarrow 1} (\sqrt{x+3} - 2) = 0$$

The latter suggests that  $x-1$  is a "factor" of  $\sqrt{x+3} - 2$ . We'll use the conjugate of  $\sqrt{x+3} - 2$ , namely  $\sqrt{x+3} + 2$ .

$$\lim_{x \rightarrow 1} \frac{\sqrt{x+3} - 2}{x-1} = \lim_{x \rightarrow 1} \left( \frac{\sqrt{x+3} - 2}{x-1} \right) \cdot \left( \frac{\sqrt{x+3} + 2}{\sqrt{x+3} + 2} \right)$$

multiply  
by 1

$$= \lim_{x \rightarrow 1} \frac{(\sqrt{x+3})^2 - 2\sqrt{x+3} + 2\sqrt{x+3} - 4}{(x-1)(\sqrt{x+3} + 2)}$$

$$= \lim_{x \rightarrow 1} \frac{x+3 - 4}{(x-1)(\sqrt{x+3} + 2)}$$

$$= \lim_{x \rightarrow 1} \frac{\cancel{x-1}}{(\cancel{x-1})(\sqrt{x+3} + 2)} = \lim_{x \rightarrow 1} \frac{1}{\sqrt{x+3} + 2} = \frac{1}{\sqrt{1+3} + 2} = \frac{1}{4}$$

## Observations

In limit taking, the form " $\frac{0}{0}$ " sometimes appears. This is called an **indeterminate form**. Standard strategies are

- (1) Try to factor the numerator and denominator to see if a common factor— $(x - c)$ —can be cancelled.
- (2) If dealing with roots, try rationalizing to reveal a common factor.

The form

$$\frac{\text{„ nonzero constant „}}{0}$$

is not indeterminate. It is undefined. When it appears, the limit doesn't exist.