

Section 4: Bernoulli Equations

Suppose $P(x)$ and $f(x)$ are continuous on some interval (a, b) and n is a real number different from 0 or 1 (not necessarily an integer). An equation of the form

$$\frac{dy}{dx} + P(x)y = f(x)y^n$$

is called a **Bernoulli** equation.

Observation: This equation has the flavor of a linear ODE, but since $n \neq 0, 1$ it is necessarily nonlinear. So our previous approach involving an integrating factor does not apply directly. Fortunately, we can use a change of variables to obtain a related linear equation.

Solving the Bernoulli Equation

$$\frac{dy}{dx} + P(x)y = f(x)y^n$$

Let u be a new dependent variable defined by

$$u = y^{1-n} \quad \text{Then} \quad \frac{du}{dx} = (1-n)y^{-n-1} \frac{dy}{dx} = (1-n)y^{-n} \frac{dy}{dx}$$

Divide the ODE by y^n

$$\underbrace{y^{-n} \frac{dy}{dx}}_{\frac{1}{1-n} \frac{du}{dx}} + P(x) \underbrace{y^{-n} y}_{y^{1-n}} = f(x) \underbrace{y^n y^{-n}}_1$$

The equation for u is

$$\frac{1}{1-n} \frac{du}{dx} + P(x)u = f(x)$$

* $n \neq 1$ so
 $1-n \neq 0$

u solves the 1st order linear equation

$$\frac{du}{dx} + (1-n)P(x)u = (1-n)f(x).$$

Note the form

$$\frac{du}{dx} + P_1(x)u = f_1(x)$$

Where $P_1(x) = (1-n)P(x)$ and $f_1(x) = (1-n)f(x)$

Solve for u using an integrating factor.

Since $u = y^{1-n}$, $y = u^{\frac{1}{1-n}}$.

Example

Solve the initial value problem $y' - y = -e^{2x}y^3$, subject to $y(0) = 1$.

The equation is a Bernoulli ODE $\frac{dy}{dx} - y = -e^{2x}y^3$, $n=3$.

Let $u = y^{1-3} = y^{-2}$. Then $\frac{du}{dx} = -2y^{-3} \frac{dy}{dx}$

Divide by y^3

$$y^{-3} \frac{dy}{dx} - y^{-3} y = -e^{2x} \underbrace{y^{-3} y^{-3}}_1$$

$\frac{-1}{2} \frac{du}{dx} \quad u$

So u solves the ODE $\frac{-1}{2} \frac{du}{dx} - u = -e^{2x}$.

In standard form

$$\frac{du}{dx} + zu = 2e^{2x} \quad P_1(x) = 2$$

The integrating factor $\mu = e^{\int P_1(x) dx} = e^{\int 2 dx} = e^{2x}$

Using μ

$$\frac{d}{dx} (e^{2x} u) = 2e^{2x} e^{2x} = 2e^{4x}$$

$$\int \frac{d}{dx} (e^{2x} u) dx = \int 2e^{4x} dx$$

$$e^{2x} u = \frac{1}{2} e^{4x} + C$$

$$u = \frac{\frac{1}{2} e^{4x} + C}{e^{2x}} = \frac{1}{2} e^{2x} + C e^{-2x}$$

$$u = y^{-2} \quad \text{so} \quad y = u^{-\frac{1}{2}} = \frac{1}{\sqrt{u}}$$

$$y = \frac{1}{\sqrt{\frac{1}{2}e^{2x} + Ce^{-2x}}}$$

Now impose $y(0) = 1$.

$$1 = \frac{1}{\sqrt{\frac{1}{2}e^0 + Ce^0}} = \frac{1}{\sqrt{\frac{1}{2} + C}}$$

$$\sqrt{\frac{1}{2} + C} = 1$$

$$\frac{1}{2} + C = 1^2 = 1 \Rightarrow C = 1 - \frac{1}{2} = \frac{1}{2}$$

The solution to the IVP is

$$y = \frac{1}{\sqrt{\frac{1}{2}e^{2x} + \frac{1}{2}e^{-2x}}} = \frac{1}{\frac{1}{\sqrt{2}} \sqrt{e^{2x} + e^{-2x}}}$$
$$= \frac{\sqrt{2}}{\sqrt{e^{2x} + e^{-2x}}}$$

The solution to the IVP is

$$y = \frac{\sqrt{2}}{\sqrt{e^{2x} + e^{-2x}}}$$

Exact Equations

We considered first order equations of the form

$$M(x, y) dx + N(x, y) dy = 0. \quad (1)$$

The left side is called a *differential form*. We will assume here that M and N are continuous on some (shared) region in the plane.

Definition: The equation (1) is called an **exact equation** on some rectangle R if there exists a function $F(x, y)$ such that

$$\frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y)$$

for every (x, y) in R .

↑ Derivative of F with respect to x holding y constant

↑ Derivative of F wrt y holding x constant.

Exact Equation Solution

If $M(x, y) dx + N(x, y) dy = 0$ happens to be exact, then it is equivalent to

$$\frac{\partial F}{\partial x} dx + \frac{\partial F}{\partial y} dy = 0$$

This implies that the function F is constant on R and solutions to the DE are given by the relation

$$F(x, y) = C$$

The relation defines solutions to the ODE implicitly.

Recognizing Exactness

There is a theorem from calculus that ensures that if a function F has first partials on a domain, and if those partials are continuous, then the second mixed partials are equal. That is,

$$\frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y}.$$

If it is true that

$$\frac{\partial F}{\partial x} = M \quad \text{and} \quad \frac{\partial F}{\partial y} = N$$

this provides a condition for exactness, namely

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

$$\text{IF } \frac{\partial F}{\partial x} = M, \text{ then}$$

$$\frac{\partial^2 F}{\partial y \partial x} = \frac{\partial M}{\partial y}$$

and

$$\text{IF } \frac{\partial F}{\partial y} = N, \text{ then}$$

$$\frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x}$$

Exact Equations

Theorem: Let M and N be continuous on some rectangle R in the plane. Then the equation

$$M(x, y) dx + N(x, y) dy = 0$$

is exact if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}.$$

Example

Show that the equation is exact and obtain a family of solutions.

$$(2xy - \sec^2 x) dx + (x^2 + 2y) dy = 0$$

$$M(x,y) = 2xy - \sec^2 x \quad \text{and} \quad N(x,y) = x^2 + 2y$$

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y} (2xy - \sec^2 x) = 2x \cdot 1 - 0 = 2x$$

Treat x as a constant

$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x} (x^2 + 2y) = 2x + 0 = 2x$$

Treat y as a constant

$$\frac{\partial M}{\partial y} = 2x \frac{\partial N}{\partial x} \quad \text{the equation is exact.}$$

The solutions are given by the relation

$$F(x, y) = C,$$

for some function F . We know that

$$\frac{\partial F}{\partial x} = M(x, y) = 2xy - \sec^2 x \quad \text{and}$$

$$\frac{\partial F}{\partial y} = N(x, y) = x^2 + 2y$$

This is enough information to find F
up to an added constant.