

## Section 1.3: Continuity

**Definition: Continuity at a Point** A function  $f$  is continuous at a number  $c$  if

$$\lim_{x \rightarrow c} f(x) = f(c).$$

This definition is equivalent to the three statements

- (1)  $f(c)$  is defined (i.e.  $c$  is in the domain of  $f$ ),
- (2)  $\lim_{x \rightarrow c} f(x)$  exists, and
- (3) the limit actually equals the function value.

If a function  $f$  is not continuous at  $c$ , we may say that  $f$  is **discontinuous** at  $c$

## Question

Suppose  $f$  is continuous at  $-4$  and  $f(-4) = 2\pi$ . Then

$$\lim_{x \rightarrow -4} f(x) = f(-4) = 2\pi$$

(a)  $-4$

(b)  $-8\pi$

(c)  $2\pi$

(d) can't be determined without more information

## A Theorem on Continuous Functions

**Theorem** If  $f$  and  $g$  are continuous at  $c$  and for any constant  $k$ , the following are also continuous at  $c$ :

$$(i) f + g, \quad (ii) f - g, \quad (iii) kf, \quad (iv) fg, \quad \text{and} \quad (v) \frac{f}{g}, \text{ if } g(c) \neq 0.$$

In other words, if we combine continuous functions using addition, subtraction, multiplication, division, and using constant factors, the result is also continuous—provided of course that we don't introduce division by zero.

# Continuity on an Interval

**Definition** A function is continuous on an interval  $(a, b)$  if it is continuous at each point in  $(a, b)$ . A function is continuous on an interval such as  $(a, b]$  or  $[a, b)$  or  $[a, b]$  provided it is continuous on  $(a, b)$  and has one sided continuity at each included end point.

Graphically speaking, if  $f(x)$  is continuous on an interval  $(a, b)$ , then the curve  $y = f(x)$  will have no holes or gaps.

Find all values of  $A$  such that  $f$  is continuous on  $(-\infty, \infty)$ .

$$f(x) = \begin{cases} x + A, & x < 2 \\ Ax^2 - 3, & 2 \leq x \end{cases}$$

The pieces are continuous.  
we need to know what happens @ 2.

We require  $\lim_{x \rightarrow 2} f(x) = f(2)$ .

$$f(2) = A(2^2) - 3 = 4A - 3$$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (x+A) = 2+A$$

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (Ax^2-3) = 4A-3$$

For existence of the limit, it must be that

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x)$$

$$2+A = 4A-3$$

$$5 = 3A \Rightarrow A = \frac{5}{3}$$

So  $f$  is continuous @ 2 if  $A = \frac{5}{3}$ .

$$\text{Then } \lim_{x \rightarrow 2} f(x) = 2 + \frac{5}{3} = \frac{11}{3}$$

$$\text{and } f(2) = 4\left(\frac{5}{3}\right) - 3 = \frac{20}{3} - \frac{9}{3} = \frac{11}{3}$$

$$f(x) = \begin{cases} x + A, & x < 2 \\ Ax^2 - 3, & 2 \leq x \end{cases}$$





Figure: On the left,  $A \neq \frac{5}{3}$ ; on the right  $A = \frac{5}{3}$ .



# Compositions

Suppose  $\lim_{x \rightarrow c} g(x) = L$ , and  $f$  is continuous at  $L$ , then

$$\lim_{x \rightarrow c} f(g(x)) = f(L) \quad \text{i.e.} \quad \lim_{x \rightarrow c} f(g(x)) = f\left(\lim_{x \rightarrow c} g(x)\right).$$

**Theorem:** If  $g$  is continuous at  $c$  and  $f$  is continuous at  $g(c)$ , then  $(f \circ g)(x)$  is continuous at  $c$ .

Essentially, this says that "compositions of continuous functions are continuous."

## Example

Suppose we know that  $f(x) = e^x$  is continuous on  $(-\infty, \infty)$ <sup>1</sup>. Evaluate

$$\lim_{x \rightarrow \sqrt{\ln(3)}} e^{x^2 + \ln(2)}$$

If  $g(x) = x^2 + \ln 2$ , then  $g$  is continuous for all real  $x$ .

$$\text{Note } e^{x^2 + \ln 2} = f(g(x))$$

$$\begin{aligned} \text{So } \lim_{x \rightarrow \sqrt{\ln 3}} e^{x^2 + \ln 2} &= e^{(\sqrt{\ln 3})^2 + \ln 2} \\ &= e^{\ln 3 + \ln 2} \\ &= e^{\ln 3} \cdot e^{\ln 2} = 3 \cdot 2 = 6 \end{aligned}$$

---

<sup>1</sup>This is true.

# Inverse Functions



**Theorem:** If  $f$  is a one to one function that is continuous on its domain, then its inverse function  $f^{-1}$  is continuous on its domain.

Continuous functions (with inverses) have continuous inverses.

## Theorem:

**Intermediate Value Theorem (IVT)** Suppose  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number between  $f(a)$  and  $f(b)$ . Then there exists  $c$  in the interval  $(a, b)$  such that  $f(c) = N$ .

The continuous function passes through every number between  $f(a)$  and  $f(b)$ .



## Application of the IVT

Show that the equation has at least one solution in the interval.

$$x^3 + x^2 - 4 = 0 \quad 1 \leq x \leq 2$$

Let  $f(x) = x^3 + x^2 - 4$ . As a polynomial,  $f$  is continuous @ all reals, so it's continuous on  $[1, 2]$ .

$$f(1) = 1^3 + 1^2 - 4 = 2 - 4 = -2$$

$$f(2) = 2^3 + 2^2 - 4 = 8 + 4 - 4 = 8$$

Note that if  $N=0$  then


$$f(1) < N < f(2)$$

i.e. 0 is between -2 and 8.

By the IVT, there must be a number  $c$  in  $(1, 2)$  such that  $f(c) = N$

$$\text{That is, } f(c) = 0 \Rightarrow$$

$$c^3 + c^2 - 4 = 0$$



## Section 1.4: Limits and Continuity of Trigonometric, Exponential and Logarithmic Functions

Here we list without proof<sup>2</sup> the continuity properties of several well known functions.

**sin  $x$ :** The sine function  $y = \sin x$  is continuous on its domain  $(-\infty, \infty)$ .

**cos  $x$ :** The cosine function  $y = \cos x$  is continuous on its domain  $(-\infty, \infty)$ .

**$e^x$ :** The exponential function  $y = e^x$  is continuous on its domain  $(-\infty, \infty)$ .

**ln( $x$ ):** The natural log function  $y = \ln(x)$  is continuous on its domain  $(0, \infty)$ .

---

<sup>2</sup>You are already familiar with their graphs.



## Additional Functions

- ▶ By the quotient property, each of  $\tan x$ ,  $\cot x$ ,  $\sec x$  and  $\csc x$  are continuous on each of their respective domains.
- ▶ For  $a > 0$  with  $a \neq 1$ , the function

$$a^x = e^{x \ln a}.$$

By the composition property, each exponential function  $y = a^x$  is continuous on  $(-\infty, \infty)$ .

- ▶ For  $a > 0$  with  $a \neq 1$ , the function

$$\log_a(x) = \frac{\ln x}{\ln a}.$$

By the constant multiple property, each logarithm function  $y = \log_a(x)$  is continuous on  $(0, \infty)$ .

## What does all this mean?

The common functions we use, polynomial and rational functions, trigonometric functions, and logs and exponentials are continuous **everywhere on their respective domains.**

So, if  $f$  is anyone of these functions and  $c$  is a number in its domain, then  $\lim_{x \rightarrow c} f(x) = f(c)$ .

## Example

Evaluate each limit.

(a)  $\lim_{x \rightarrow \pi} \cos(x + \sin x)$

$g(x) = x + \sin x$  is  
continuous @  $\pi$

$f(x) = \cos(x)$  is continuous  
at  $g(\pi)$

So

$$\begin{aligned} \lim_{x \rightarrow \pi} \cos(x + \sin x) &= \cos(\pi + \sin \pi) \\ &= \cos(\pi + 0) = \cos(\pi) = -1 \end{aligned}$$

## Example

$$(b) \lim_{t \rightarrow \frac{\pi}{4}} e^{\tan t}$$

$g(t) = \tan t$  is continuous  
@  $\frac{\pi}{4}$

$f(t) = e^t$  is continuous  
everywhere.

$$\lim_{t \rightarrow \frac{\pi}{4}} e^{\tan t} = e^{\tan \frac{\pi}{4}} = e^1 = e$$

## Question

Evaluate the limit  $\lim_{x \rightarrow \pi} \ln(\cos^2 x)$ .

$$(\cos \pi)^2 = (-1)^2 = 1$$

(a)  $e$


$$\lim_{x \rightarrow \pi} \ln(\cos^2 x) = \ln(\cos^2 \pi)$$

(b) 1

$$= \ln 1 = 0$$

(c) DNE

(d) 0



## Squeeze Theorem:

**Theorem:** Suppose  $f(x) \leq g(x) \leq h(x)$  for all  $x$  in an interval containing  $c$  except possibly at  $c$ . If

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x) = L$$

then

$$\lim_{x \rightarrow c} g(x) = L.$$

Squeeze Theorem:

$$f(x) \leq g(x) \leq h(x)$$




Figure: Graphical Representation of the Squeeze Theorem.

# Example: Evaluate

$$\lim_{\theta \rightarrow 0} \theta^2 \sin \frac{1}{\theta}$$

Direct substitution doesn't work  
since " $\sin \frac{1}{0}$ " doesn't make sense.

We do know the sine is bounded.

$$-1 \leq \sin \frac{1}{\theta} \leq 1$$

$$-1 \cdot \theta^2 \leq \theta^2 \sin \frac{1}{\theta} \leq 1 \cdot \theta^2$$

$f(\theta)$                    $g(\theta)$                    $h(\theta)$


mult. by  $\theta^2$   
which is  
never negative

$$\lim_{\theta \rightarrow 0} -\theta^2 = 0 \quad \lim_{\theta \rightarrow 0} \theta^2 = 0 \Rightarrow \lim_{\theta \rightarrow 0} \theta^2 \sin \frac{1}{\theta} = 0 \text{ by the Squeeze Thm.}$$



# A Couple of Important Limits

**Theorem:**  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$  and  $\lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\theta} = 0$



-  $\sin \theta$

-  $\theta$

-  $\tan \theta$

$\theta \geq 0$

Note  $\sin \theta \leq \theta \leq \tan \theta$

We'll work with

①  $\sin \theta \leq \theta$  and

②  $\theta \leq \tan \theta$

① for  $\theta > 0$   $\sin \theta \leq \theta$  Divide by  $\theta$

$$\frac{\sin \theta}{\theta} \leq 1$$

Note  $\cdot$   $\frac{\sin(-\theta)}{-\theta} = \frac{-\sin \theta}{-\theta} = \frac{\sin \theta}{\theta}$

so  $\frac{\sin \theta}{\theta} \leq 1$  for all  $\theta \neq 0$

② for  $\theta > 0$   $\theta \leq \tan \theta \Rightarrow \theta \leq \frac{\sin \theta}{\cos \theta}$

for  $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$ ,  $\cos \theta > 0$

Multiply by  $\cos \theta$  and divide by  $\theta$

$$\cos \theta \leq \frac{\sin \theta}{\theta}$$

Since  $\cos(-\theta) = \cos \theta$  and  $\frac{\sin(-\theta)}{-\theta} = \frac{\sin \theta}{\theta}$

So for all  $\theta$  "close to zero"

$$\cos \theta \leq \frac{\sin \theta}{\theta} \leq 1$$

Note  $\lim_{\theta \rightarrow 0} \cos \theta = \cos 0 = 1$  and  $\lim_{\theta \rightarrow 0} 1 = 1$

By the squeeze theorem  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$ .

## Notational Note

The name of the variable used is irrelevant. That is

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1, \quad \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1, \quad \lim_{\heartsuit \rightarrow 0} \frac{\sin \heartsuit}{\heartsuit} = 1$$

In fact, this only requires the argument of the sine to match the denominator (exactly) and that this term is tending to zero. For example,

$$\lim_{\theta \rightarrow 0} \frac{\sin(6\theta)}{6\theta} = 1, \quad \text{and} \quad \lim_{\heartsuit \rightarrow 0} \frac{\sin(\pi\heartsuit)}{\pi\heartsuit} = 1$$

## Examples

Use  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$  to evaluate each limit.

(a)  $\lim_{x \rightarrow 0} \frac{\sin(4x)}{x}$

If we had  $\lim_{x \rightarrow 0} \frac{\sin(4x)}{4x}$  the limit would be 1.

this is 1  
↓

$$\lim_{x \rightarrow 0} \frac{\sin(4x)}{x} = \lim_{x \rightarrow 0} \frac{\sin(4x)}{x} \cdot \left(\frac{4}{4}\right)$$

$$= \lim_{x \rightarrow 0} 4 \left( \frac{\sin(4x)}{4x} \right) = 4 \cdot 1 = 4$$

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

$$(b) \quad \lim_{t \rightarrow 0} t \csc(3t)$$

$$= \lim_{t \rightarrow 0} \frac{t}{\sin(3t)}$$

$$= \lim_{t \rightarrow 0} \frac{t}{\sin(3t)} \cdot \frac{3}{3}$$

$$= \lim_{t \rightarrow 0} \frac{1}{3} \left( \frac{3t}{\sin(3t)} \right)$$

$$= \lim_{t \rightarrow 0} \frac{1}{3} \left( \frac{\sin(3t)}{3t} \right)^{-1}$$

$$= \frac{1}{3} (1)^{-1} = \frac{1}{3} \cdot \frac{1}{1} = \frac{1}{3}$$

Note

$$\lim_{\theta \rightarrow 0} \frac{\theta}{\sin \theta} = 1 \text{ as well}$$

# Questions

(1) Evaluate if possible  $\lim_{x \rightarrow 0} \frac{\tan(2x)}{4x}$

(a)  $\frac{1}{4}$

(b)  $\frac{1}{2}$

(c) 2

(d) DNE

$$= \lim_{x \rightarrow 0} \frac{\sin(2x)}{\cos(2x) \cdot 4x}$$

$$= \lim_{x \rightarrow 0} \frac{1}{2 \cos(2x)} \cdot \frac{\sin(2x)}{2x}$$



## A couple of important observations

$$\lim_{x \rightarrow 0} \cos x = 1, \quad \text{so for example} \quad \lim_{x \rightarrow 0} \frac{\cos x}{x} \text{ DNE}$$

While it is true that  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ , the statement

$$\frac{\sin x}{x} = 1$$

is **always false!** Don't be tempted to write this.

Also remember that  $\sin(kx) \neq k\sin(x)$ . Don't be tempted to try to *factor* out of a trig function.