

## Section 4.2: Maximum and Minimum Values; Critical Numbers

Let's start with some review questions inspired by last Thursday's quiz!

## Question

Let  $y = x^{e^x}$ . Which of the following is true?

- (a) The power rule gives  $y' = (e^x)x^{e^x-1}$ .
- (b) The exponential rule gives  $y' = x^{e^x} \ln x$ .
- (c)  $y$  is the product of  $x$  and  $e^x$ .
- (d)** There is no derivative rule that applies directly to  $y$ .

power function  
 $x^n$  ← constant power  
↑  
variable base

exponential function  
 $a^x$  ← variable power  
→  
constant base

## Question

Let  $y = x^{e^x}$ . Which of the following is true?

(a)  $y = e^x \ln x$

(b)  $\ln y = e^x \ln x$

(c)  $\frac{1}{y} = e^x \frac{1}{x}$

(d)  $\ln y = e \ln x^x$

$$\begin{aligned}\ln y &= \ln x^{e^x} \\ &= e^x \ln x\end{aligned}$$

## Question

Given that  $\ln y = e^x \ln x$ ,

(a)  $\frac{dy}{dx} = e^x \ln x + \frac{e^x}{x}$

(b)  $\frac{dy}{dx} = x^{e^x} \left( \frac{e^x}{x} \right)$

(c)  $\frac{dy}{dx} = \frac{e^x}{x}$

(d)  $\frac{dy}{dx} = x^{e^x} \left( e^x \ln x + \frac{e^x}{x} \right)$

*implicit*

*product rule*

$$\frac{d}{dx} \ln y = \frac{d}{dx} e^x \ln x$$

$$\frac{1}{y} \frac{dy}{dx} = e^x \ln x + e^x \cdot \frac{1}{x}$$


$$\frac{dy}{dx} = y \left( e^x \ln x + \frac{e^x}{x} \right)$$

$$= x^{e^x} \left( e^x \ln x + \frac{e^x}{x} \right)$$

## Question

True or **False**:  $\ln x = \frac{1}{x}$

But  $\frac{d}{dx} \ln x = \frac{1}{x}$



## Question

Given  $y = \ln(\ln x)$ , find  $y'$ .

(a)  $y' = \frac{1}{x \ln x}$

(b)  $y' = \frac{1}{\ln(\ln x)}$

(c)  $y' = \frac{1}{(\ln x)^2}$

(d)  $y' = \frac{1}{\ln x}$

If  $u = \ln x$  and  $f(u) = \ln u$

$$y = f(u)$$

$$y' = f'(u) \cdot u'(x)$$

$$= \frac{1}{u} \cdot \frac{1}{x} = \frac{1}{x \ln x}$$

$$y' = \frac{\frac{1}{x}}{\ln x} = \frac{1}{x \ln x}$$

## Section 4.3: The Mean Value Theorem

### Recall

**Rolle's Theorem:** Let  $f$  be a function that is

- i continuous on the closed interval  $[a, b]$ ,
- ii differentiable on the open interval  $(a, b)$ , and
- iii such that  $f(a) = f(b)$ .

Then there exists a number  $c$  in  $(a, b)$  such that  $f'(c) = 0$ .

If  $f$  takes the same  $y$ -value at both ends, and is continuous and *smooth* in between them, then it's either horizontal, or it turns around. Somewhere there must be a horizontal tangent!

# The Mean Value Theorem

**Theorem:** Suppose  $f$  is a function that satisfies

- i  $f$  is continuous on the closed interval  $[a, b]$ , and
- ii  $f$  is differentiable on the open interval  $(a, b)$ .

Then there exists a number  $c$  in  $(a, b)$  such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}, \quad \text{equivalently} \quad f(b) - f(a) = f'(c)(b - a).$$

If  $f$  is continuous and *smooth* between the point  $(a, f(a))$  and  $(b, f(b))$ , then somewhere between them is a point where the tangent is parallel to the secant line connecting the ends.



## Rate of Change and Slope

**From day 1 of class:** Given a line  $y = mx + b$ , and a change in the independent variable  $\Delta x$ ,

the change in the dependent variable is

$$\Delta y = m\Delta x.$$

**Now look at the conclusion of the MVT!**

$$f(b) - f(a) = f'(c) (b - a)$$

change in  $y$  = a slope change in  $x$

$$\Delta y = m \Delta x$$

## A couple of questions we can ask...

**An Interesting Question:** We know that  $f'(x) = 0$  if  $f$  is a constant function. Is the converse true? That is, if  $f'(x) = 0$  does that mean that  $f$  is a constant function?

**Another Interesting Question:** If we have two functions  $f$  and  $g$  such that on some interval

$$f'(x) = g'(x),$$

does this mean that  $f$  and  $g$  have to be the same function?

# Important Consequence of the MVT

**An Interesting Question:** We know that  $f'(x) = 0$  if  $f$  is a constant function. Is the converse true? That is, if  $f'(x) = 0$  does that mean that  $f$  is a constant function?

yes!

**Theorem:** If  $f'(x) = 0$  for all  $x$  in an interval  $(a, b)$ , then  $f$  is constant on  $(a, b)$ .

only constant functions have derivatives that are zero on an interval.

## Important Consequence of the MVT

**Another Interesting Question:** If we have two functions  $f$  and  $g$  such that on some interval

$$f'(x) = g'(x),$$

does this mean that  $f$  and  $g$  have to be the same function? **No**

*But they can only differ by an added constant.*

**Corollary:** If  $f'(x) = g'(x)$  for all  $x$  in an interval  $(a, b)$ , then  $f - g$  is constant on  $(a, b)$ . In other words,

$$f(x) = g(x) + C \quad \text{where } C \text{ is some constant.}$$

## Examples

Find all possible functions  $f(x)$  that satisfy the condition

(a)  $f'(x) = \cos x$  on  $(-\infty, \infty)$

Since  $\frac{d}{dx} \sin x = \cos x$

all functions can only differ by an 'added constant. So

$f(x) = \sin x + C$  for any constant  $C$ .

(b)  $f'(x) = 2x$  on  $(-\infty, \infty)$

$$\frac{d}{dx} x^2 = 2x$$

So all such functions are

$$f(x) = x^2 + C \text{ for arbitrary constant } C.$$

## Question

Find all possible functions  $h(t)$  that satisfy the condition

(c)  $h'(t) = \sec^2 t$  on  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

(a)  $h(t) = \sec^2 t + C$

(b)  $h(t) = \tan t + 1$

(c)  $h(t) = \tan t + C$

## Another Consequence of the MVT

Another significant consequence of the MVT is that it provides a test for the increasing and decreasing behavior of a differentiable function.

**Theorem:** Let  $f$  be differentiable on an open interval  $(a, b)$ . If

- ▶  $f'(x) > 0$  on  $(a, b)$ , the  $f$  is increasing on  $(a, b)$ , and if
- ▶  $f'(x) < 0$  on  $(a, b)$ , the  $f$  is decreasing on  $(a, b)$ .



## Example

Determine the intervals over which  $f$  is increasing and the intervals over which it is decreasing where

$$f(x) = 6x^4 - 32x^3 - 7 \quad \text{Domain is } (-\infty, \infty)$$

- find  $f'$
- find where  $f'(x) = 0$  or where  $f'(x)$  is undefined.
- Consider the intervals determined by the numbers found.
- Do a sign analysis by checking the sign of  $f'$  in each interval.
- State our conclusion

$$f(x) = 6x^4 - 32x^3 - 7$$

$$f'(x) = 24x^3 - 96x^2 = 24x^2(x-4)$$

$f'$  is always defined,  $f'(x) = 0 \Rightarrow$

$$24x^2(x-4) = 0$$

$$\Rightarrow x^2 = 0 \quad \text{or} \quad x-4 = 0$$

$$x = 0 \quad \text{or} \quad x = 4$$

This breaks the domain into 3 intervals,  
 $(-\infty, 0)$ ,  $(0, 4)$ , and  $(4, \infty)$ .

$$f'(x) = 24x^2(x-4)$$

## Sign analysis



Pick test points in each interval

test	-1	$f'(-1) = 24(-1)^2(-1-4) = -120$	negative
	+1	$f'(1) = 24(1)^2(1-4) = -72$	negative
	5	$f'(5) = 24(5)^2(5-4) = 600$	positive.

Based on the analysis

$f$  is decreasing on  $(-\infty, 0) \cup (0, 4)$ .

$f$  is increasing on  $(4, \infty)$ .

\* The conclusion is about the behavior of  
 $f$  on intervals.


## Question

Suppose that we compute the derivative of some function  $g$  and find

$$g'(x) = (2 + x)e^{x/2}.$$

Determine the intervals over which  $g$  is increasing and over which it is decreasing.

(a)  $g$  is increasing on  $(-1/2, \infty)$  and decreasing on  $(-\infty, -1/2)$ .

 (b)  $g$  is increasing on  $(-2, \infty)$  and decreasing on  $(-\infty, -2)$ .

(c)  $g$  is increasing on  $(2, \infty)$  and decreasing on  $(-\infty, 2)$ .

(d)  $g$  is increasing on  $(-\infty, -2)$  and decreasing on  $(-2, \infty)$ .

## Section 4.4: Local Extrema and Concavity

We have already seen that the first derivative  $f'$  can tell us about the behaviour of the function  $f$ —in particular, it gives information about where it is increasing or decreasing, and where it may take a local extreme value.

In this section, we'll expand on that as well as introduce information about a function that can be deduced from the nature of its second derivative.

## Theorem: First derivative test for local extrema

Let  $f$  be continuous and suppose that  $c$  is a critical number of  $f$ .

- ▶ If  $f'$  changes from negative to positive at  $c$ , then  $f$  has a local minimum at  $c$ .
- ▶ If  $f'$  changes from positive to negative at  $c$ , then  $f$  has a local maximum at  $c$ .
- ▶ If  $f'$  does not change signs at  $c$ , then  $f$  does not have a local extremum at  $c$ .

Note: we read from left to right as usual when looking for a sign change.




Figure: First derivative test



## Example

Find all the critical points of the function and classify each one as a local maximum, a local minimum, or neither.

$$f(x) = 16\sqrt[3]{x} - x\sqrt[3]{x} = 16x^{1/3} - x^{4/3}$$

first we'll do the same sign analysis.

$$\begin{aligned} f'(x) &= 16\left(\frac{1}{3}x^{-2/3}\right) - \frac{4}{3}x^{1/3} \\ &= \frac{16}{3}x^{-2/3} - \frac{4}{3}x^{1/3} = \frac{4}{3}x^{-2/3}(4-x) \\ &= \frac{4(4-x)}{3x^{2/3}} \end{aligned}$$

$f'(x) = 0$  if the numerator is zero


$$4(4-x) = 0 \Rightarrow x = 4$$

$f'(x)$  is undefined if the denominator is zero

$$3x^{2/3} = 0 \Rightarrow x = 0$$

The domain is all reals, and we have two critical numbers.

$$f'(x) = \frac{4(4-x)}{3x^{2/3}}$$



Test points:

$$-1 \quad f'(-1) = \frac{4(4 - (-1))}{3(-1)^{2/3}} \quad \begin{matrix} (+) \\ (+) \end{matrix} \quad \text{positive}$$

$$1 \quad f'(1) = \frac{4(4 - 1)}{3(1)^{2/3}} \quad \begin{matrix} (+) \\ (+) \end{matrix} \quad \text{positive}$$

$$5 \quad f'(5) = \frac{4(4 - 5)}{3(5)^{2/3}} \quad \begin{matrix} (-) \\ (+) \end{matrix}$$

Critical number zero is neither a local  
max nor min.

Critical number 4 corresponds to  
a local max.

## Question

Consider the function  $f(t) = t^4 + 4t^3$ . Which of the following is true about this function?

- (a)  $f$  has no critical numbers.
- (b)  $f$  has critical numbers 0 and  $-4$ .
- (c)  $f$  has critical numbers 4 and 12.
- (d)**  $f$  has critical numbers 0 and  $-3$ .

$$\begin{aligned}f'(t) &= 4t^3 + 12t^2 \\ &= 4t^2(t + 3)\end{aligned}$$



## Question

Consider the function  $f(t) = t^4 + 4t^3$ . Which of the following is true about this function?

- (a)  $f$  has a local minimum at  $t = 0$  and a local maximum at  $t = -3$ .
- (b)  $f$  has a local minimum at  $t = -3$  and a local maximum at  $t = 0$ .
- (c)  $f$  has a local minimum at  $t = -3$ .
- (d)  $f$  has a local minimum at  $t = 0$ .




$$f'(t) = 4t^2(t+3)$$

# Concavity and The Second Derivative


**Concavity:** refers to the *bending* nature of a graph. In particular, a curve is **concave down** if it's cupped side is down, and it is **concave up** if it's cupped upward.

# Concavity




Figure





**Figure:** A graph can have either increasing or decreasing behavior and be either concave up or down.



**Figure:** We can consider concavity at a point, but it's best thought of as a property over an interval. Many function's graphs have concavity that changes over the domain.

## Definition of Concavity

**Concave Up:** If the graph of a function  $f$  lies **above** all of its tangent lines over an interval  $I$ , then  $f$  is **concave up** on  $I$ .

**Concave Down:** If the graph of  $f$  lies **below** each of its tangent lines on an interval  $I$ ,  $f$  is **concave down** on  $I$ .

**Theorem:** (Second Derivative Test for Concavity)

Suppose  $f$  is twice differentiable on an interval  $I$ .

- ▶ If  $f''(x) > 0$  on  $I$ , then the graph of  $f$  is concave up on  $I$ .
- ▶ If  $f''(x) < 0$  on  $I$ , then the graph of  $f$  is concave down on  $I$ .

**Definition:** A point  $P$  on a curve  $y = f(x)$  is called an **inflection point** if  $f$  is continuous at  $P$  and the concavity of  $f$  changes at  $P$  (from down to up or from up to down). A point where  $f''(x) = 0$  would be a candidate for being an inflection point.

