

Section 4.4: Local Extrema and Concavity

Let's start with some review question inspired by this Thursday's quiz!

Question

The limit $\lim_{x \rightarrow 1} \frac{\sin(\pi x)}{\ln x}$ gives rise to

- (a) the indeterminate form $\frac{0}{0}$.
- (b) the indeterminate form $\frac{\infty}{\infty}$.
- (c) the indeterminate form $\infty - \infty$.
- (d) no indeterminate form, the limit is 1.

$$\sin(\pi) = 0$$

and

$$\ln 1 = 0$$

Question

Evaluate $\lim_{x \rightarrow 1} \frac{\sin(\pi x)}{\ln x}$ (use any applicable technique)

(a) $\lim_{x \rightarrow 1} \frac{\sin(\pi x)}{\ln x} = 1$

(b) $\lim_{x \rightarrow 1} \frac{\sin(\pi x)}{\ln x} = -\pi$

(c) $\lim_{x \rightarrow 1} \frac{\sin(\pi x)}{\ln x} = \infty$

(d) $\lim_{x \rightarrow 1} \frac{\sin(\pi x)}{\ln x} = 0$

Use l'H rule

$$\lim_{x \rightarrow 1} \frac{\cos(\pi x) \cdot \pi}{\frac{1}{x}} = \frac{\cos(\pi) \cdot \pi}{\frac{1}{1}}$$

$$= \frac{(-1) \pi}{1} = -\pi$$

Question

The critical numbers of the function $f(x) = x^3 - 12x$ are

(a) -2 and 2 .

(b) -2 , 0 , and -2 .

(c) nothing since f has no critical numbers.

(d) 0 and 4 .

$$\begin{aligned}f'(x) &= 3x^2 - 12 \\ &= 3(x^2 - 4)\end{aligned}$$

f' is defined everywhere

$$\begin{aligned}f'(x) = 0 &\Rightarrow 3(x^2 - 4) = 0 \\ &x = \pm 2.\end{aligned}$$

Question

Determine the absolute maximum and absolute minimum values of $f(x) = x^3 - 12x$ on the interval $[0, 3]$.

(a) The max is 15, and the min is -12.

(b) The max is 0, and the min is -9.

(c) The max is 0, and the min is -16.

(d) The max is 0, and there is no absolute minimum.

$$f(0) = 0 \leftarrow \text{max}$$

$$f(2) = -16 \leftarrow \text{min}$$

$$f(3) = -9$$

Our Latest Theorems & Definitions

Theorem: First Derivative Test: Let f be continuous and suppose that c is a critical number of f .

- ▶ If f' changes from $-$ to $+$ at c , then f has a local minimum at c .
- ▶ If f' changes from $+$ to $-$ at c , then f has a local maximum at c .
- ▶ If f' does not change signs at c , then f does not have a local extremum at c .

Definition: Concavity

Concave Up: If the graph of a function f lies **above** all of its tangent lines over an interval I , then f is **concave up** on I .

Concave Down: If the graph of f lies **below** each of its tangent lines on an interval I , f is **concave down** on I .

Our Latest Theorems & Definitions

Theorem: (Second Derivative Test for Concavity)

Suppose f is twice differentiable on an interval I .

- ▶ If $f''(x) > 0$ on I , then the graph of f is concave up on I .

- ▶ If $f''(x) < 0$ on I , then the graph of f is concave down on I .

Definition: A point P on a curve $y = f(x)$ is called an **inflection point** if f is continuous at P and the concavity of f changes at P (from down to up or from up to down).

A point where $f''(x) = 0$ would be a candidate for being an inflection point.

Example

Determine where the graph of $f(x) = x^4 - 4x^3$ is concave up, where it is concave down, and identify any x -values at which f has a point of inflection.

The domain of f is $(-\infty, \infty)$.

We'll do a sign analysis on $f''(x)$.

$$f'(x) = 4x^3 - 12x^2 = 4x^2(x-3).$$

$$f''(x) = 12x^2 - 24x = 12x(x-2)$$

f'' is defined everywhere.

$$f''(x) = 0 \quad \Rightarrow \quad 12x(x-2) = 0$$

$$x = 0 \quad \text{or} \quad x = 2$$

This divides our domain into the intervals
 $(-\infty, 0)$, $(0, 2)$, and $(2, \infty)$.

$$f''(x) = 12x(x-2)$$



Test pts :

-1	$f''(-1) = 12(-1)(-1-2)$
1	$f''(1) = 12(1)(1-2)$
3	$f''(3) = 12(3)(3-2)$

f is concave up on $(-\infty, 0) \cup (2, \infty)$.

f is concave down on $(0, 2)$.

f has inflection points @ $x=0$ and
at $x=2$.

Concavity and Extrema:

Theorem: (Second Derivative Test for Local Extrema)

Suppose $f'(c) = 0$ and that f'' is continuous near c . Then

- ▶ if $f''(c) > 0$, f takes a local minimum at c ,
- ▶ if $f''(c) < 0$, then f takes a local maximum at c .

If $f''(c) = 0$, then the test fails. f may or may not have a local extrema. You can go back to the first derivative test to find out.

Example: Consider $f(x) = x \ln x$

f has one local extreme value. Find the local extreme value and classify it as a local maximum or local minimum using the second derivative test.

The domain of f is $(0, \infty)$.

Find crit. #'s $f'(x) = 1 \cdot \ln x + x \cdot \frac{1}{x} = \ln x + 1$

f' is defined everywhere on $(0, \infty)$.

$$f'(x) = 0 \Rightarrow \ln x + 1 = 0$$

$$\Rightarrow \ln x = -1 \Rightarrow e^{\ln x} = e^{-1} \Rightarrow x = e^{-1}$$

our
one
critical
#

Will use the 2nd der. test.

$$f''(x) = \frac{1}{x}$$

Plug the crit. # into $f''(x)$

$$f''(e^{-1}) = \frac{1}{e^{-1}} = e = e$$

$$f''(e^{-1}) > 0 \quad \text{graph is concave up @ } e^{-1}$$

f takes a local minimum @ $x = e^{-1}$.

The minimum value is

$$f(e^{-1}) = e^{-1} \ln e^{-1} = e^{-1}(-1) = -\frac{1}{e}$$

Let's Analyze a Function

Consider the function $f(x) = xe^{3x}$. Let's determine

- (a) the intervals on which f is increasing and decreasing,
- (b) the intervals on which f is concave up and concave down,
- (c) identify critical points and classify any local extrema, and
- (d) identify any points of inflection.

Question

Find the first and second derivatives of $f(x) = xe^{3x}$.

(a) $f'(x) = 3e^{3x}$, $f''(x) = 9e^{3x}$

$$f'(x) = 1e^{3x} + x e^{3x} \cdot 3 \\ = e^{3x} (1 + 3x)$$

(b) $f'(x) = (1+3x)e^{3x}$, $f''(x) = (6+9x)e^{3x}$

$$f''(x) = (6+9x)e^{3x} \\ = 3(2+3x)e^{3x}$$

(c) $f'(x) = (1+3x)e^{3x}$, $f''(x) = (1+3x)e^{3x}$

(d) $f'(x) = e^{3x} + 3x^2 e^{2x}$, $f''(x) = 3e^{3x} + 6xe^{2x} + 6x^3 e^x$

Question

We found for $f(x) = xe^{3x}$ that $f'(x) = (1 + 3x)e^{3x}$. The critical number(s) of f are

(a) $\frac{1}{3}$ only

(b) $\frac{1}{3}$ and 0

(c) $-\frac{1}{3}$ only

(d) $-\frac{1}{3}$ and 0

f' is always defined

$$f'(x) = 0 \Rightarrow (1 + 3x)e^{3x} = 0$$

$$1 + 3x = 0 \quad \text{or} \quad e^{3x} = 0$$

$$x = -\frac{1}{3}$$

no
soln.

Question


We found for $f(x) = xe^{3x}$ that $f'(x) = (1 + 3x)e^{3x}$. And f has one critical number $-\frac{1}{3}$. f has the increasing/decreasing behavior

(a) f is decreasing on $(-\infty, -\frac{1}{3})$ and increasing on $(-\frac{1}{3}, \infty)$

(b) f is increasing on $(-\infty, -\frac{1}{3})$ and decreasing on $(-\frac{1}{3}, \infty)$

(c) f is increasing on $(-\infty, \infty)$

(d) f is decreasing on $(-\infty, \infty)$



Question


We found for $f(x) = xe^{3x}$ that $f''(x) = 3(2 + 3x)e^{3x}$. And $f''(x)$ has one root $-\frac{2}{3}$. f has the concavity

(a) f is concave up on $(-\infty, -\frac{2}{3})$ and concave down on $(-\frac{2}{3}, \infty)$

(b) f is concave down on $(-\infty, -\frac{2}{3})$ and concave up on $(-\frac{2}{3}, \infty)$

(c) f is concave up on $(-\infty, \infty)$

(d) f is concave down on $(-\infty, \infty)$







Figure: Plot of $y = xe^{3x}$.

$$f(x) = xe^{3x}$$

$f'(x) = (1 + 3x)e^{3x}$ and $f''(x) = 3(2 + 3x)e^{3x}$. f has one critical number $-1/3$. Observe how it can be classified as a local minimum.

Crit # $-\frac{1}{3}$

1st der. test



f takes a local minimum @ $-\frac{1}{3}$

2nd der. test

$$\begin{aligned} f''\left(-\frac{1}{3}\right) &= 3\left(2 + 3\left(-\frac{1}{3}\right)\right) e^{3\left(-\frac{1}{3}\right)} \\ &= 3(2-1) e^{-1} = 3e^{-1} > 0 \end{aligned}$$

conc. up

local minimum

Section 4.8: Antiderivatives; Differential Equations

We have a bunch of rules for taking derivatives, now we'll think about the other direction—if we know $f'(x)$, can we find f ?

Definition: A function F is called an antiderivative of f on an interval I if

$$F'(x) = f(x) \quad \text{for all } x \text{ in } I.$$

For example for $f(x) = 2x$, an antiderivative is $F(x) = x^2$
because $F'(x) = f(x)$.

General Antiderivative

Recall: The MVT told us that if $f'(x) = g'(x)$ on an interval, then they are equal except up to an added constant—i.e. $f(x) = g(x) + C$ for some constant C . We'll use this again.

Theorem: If F is any antiderivative of f on an interval I , then the *most general* antiderivative of f on I is

$$F(x) + C \quad \text{where } C \text{ is an arbitrary constant.}$$

For example, the most general antiderivative of
 $f(x) = 2x$ is $x^2 + C$. , well write this
as $F(x) = x^2 + C$

Find the most general antiderivative of f .

(i) $f(x) = 2e^{2x}$, $I = (-\infty, \infty)$ e^{2x} is an anti-der.

The most general is

$$F(x) = e^{2x} + C \quad \text{for constant } C.$$

(ii) $f(x) = \frac{1}{x}$, $I = (0, \infty)$

$$F(x) = \ln x + C$$

Question: Find the most general antiderivative of f .

(iii) $f(x) = \sin x$, $I = (-\infty, \infty)$

(a) $F(x) = \cos x$

(b) $F(x) = \cos x + C$

(c) $F(x) = -\cos x + C$

Question: Find the most general antiderivative of f .

(iv) $f(x) = \sec x \tan x, \quad I = \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

(a) $F(x) = \sec x$

(b) $F(x) = \sec x + C$

(c) $F(x) = \tan x + C$