



## Section 6.6: Graphing Trigonometric Functions with Transformations




$$f(x) = \sin(x)$$




$$f(x) = \cos(x)$$

**Figure:** Recall the basic graphs of the six trigonometric functions. Sine and Cosine

# Basic Trigonometric Graphs




$$f(x) = \tan(x)$$




$$f(x) = \cot(x)$$

Figure: Recall the basic graphs of the six trigonometric functions. Tangent and Cotangent

## Basic Trigonometric Graphs



$$f(x) = \csc(x)$$



$$f(x) = \sec(x)$$

**Figure:** Recall the basic graphs of the six trigonometric functions. Cosecant and Secant

## Question

The fundamental period of  $\sin x$  and  $\cos x$  is

(a)  $\pi$

(b)  $2\pi$

(c)  $2n\pi$  for integers  $n$

(d) there's no such thing as a *fundamental* period

## Question

$\sin x = 0$  whenever  $x$  is

- (a)  $\frac{\pi}{2}$
- (b) any negative number
- (c) any real number
- (d)  $n\pi$  whenever  $n$  is an integer

# Transformations on Sine and Cosine

Our goal is to graph functions of the form

$$f(x) = a \sin(bx - c) + d \quad \text{or} \quad f(x) = a \cos(bx - c) + d$$

Note: here we will be graphing points  $(x, y)$  on a curve  $y = f(x)$ .

# Amplitude

Consider:  $f(x) = a \sin(bx - c) + d$  or  $f(x) = a \cos(bx - c) + d$

**Definition:** Let  $a$  be any nonzero real number. The **amplitude** of the function  $f$  defined above is the value  $|a|$ .

Recall that this is half the distance between the maximum and minimum values.

If  $a < 0$  the graph is reflected in the  $x$ -axis. But the amplitude is still  $|a|$ .

# Period

Consider:  $f(x) = a \sin(bx - c) + d$  or  $f(x) = a \cos(bx - c) + d$

**Theorem:** Let  $b$  be any positive real number. The **fundamental period** of the function  $f$  above is given by

$$T = \frac{2\pi}{b}.$$

Recall that the fundamental period of  $\cos x$  and  $\sin x$  is  $2\pi$ .

Due to symmetry, we can always assume  $b > 0$ . Note for example


$$\sin(-3x + 2) = \sin(-(3x - 2)) = -\sin(3x - 2).$$

The period is **always positive**. The period in this example is  $\frac{2\pi}{3}$ .  
Allowing  $b$  to be signed, the period would be written as

$$T = \frac{2\pi}{|b|}.$$




# Period



**Figure:** Comparisons with  $b = 1/2, 1,$  and  $2$ . On the interval  $-2\pi < x < 2\pi$  we obtain one ( $b = 1/2$ ), two ( $b = 1$ ) or four ( $b = 2$ ) full cycles.

## Period



**Figure:** Comparisons with  $b = 1/2, 1$ , and  $2$ . On the interval  $-2\pi < x < 2\pi$  we obtain one ( $b = 1/2$ ), two ( $b = 1$ ) or four ( $b = 2$ ) full cycles.

## Example

Identify the period of each function.

(a)  $f(x) = 3 \sin(4x-2)+1$       period  $T = \frac{2\pi}{4} = \frac{\pi}{2}$   
 $b = 4$

(b)  $f(x) = -5 \sin\left(\frac{\pi x}{2}\right)+7$       period  $T = \frac{2\pi}{\pi/2} = 4$   
 $b = \frac{\pi}{2}$

(c)  $f(x) = 2-6 \cos(2x+3)$        $T = \frac{2\pi}{2} = \pi$   
 $b = 2$

# Frequency

Consider:  $f(x) = a \sin(bx - c) + d$  or  $f(x) = a \cos(bx - c) + d$

**Definition:** The reciprocal of the period is called the **frequency**. That is


$$\text{frequency} = \frac{1}{T} = \frac{b}{2\pi}.$$

If  $x$  represents time, then

- ▶ the period tells us how much time is required for one full cycle, and
- ▶ the frequency tells us how many cycles occur in one time unit.

If  $y = \cos(bx)$  (or  $y = \sin(bx)$ ), then  $b$  the number of cycles occurring in an interval of length  $2\pi$ .

## Question



The figure shows  $y = f(x)$  for  $0 < x < 2\pi$ .  
Which of the following is true?

(a)  $f(x) = 2 \sin\left(\frac{1}{3}x\right)$

(b)  $f(x) = 2 \sin(3x)$

(c)  $f(x) = \frac{1}{2} \sin\left(\frac{1}{3}x\right)$

(d)  $f(x) = \frac{1}{2} \sin(3x)$

Figure: Hint: Count the number of full cycles.



## Phase Shift (horizontal shift)

Consider:  $f(x) = a \sin(bx - c) + d$  or  $f(x) = a \cos(bx - c) + d$

**Definition:** A horizontal shift is called a **phase shift**. Again assuming that  $b > 0$ , the phase shift for  $f$  above is

$$\frac{|c|}{b} \text{ units}$$

to the **right** if  $c > 0$  and to the **left** if  $c < 0$ .



## Example

Identify the phase shift of each function. Determine if it is left or right.

(a)  $f(x) = 3 \sin(4x - 2) + 1$

$b = 4$     $c = 2$

phase shift  $\frac{|c|}{b} = \frac{2}{4} = \frac{1}{2}$   
to the right

(b)  $f(x) = -5 \sin\left(\frac{\pi x}{2}\right) + 7$

$b = \frac{\pi}{2}$     $c = 0$

No phase shift

(c)  $f(x) = 2 - 6 \cos(2x + 3)$

$b = 2$     $c = -3$

phase shift  $\frac{|c|}{b} = \frac{|-3|}{2} = \frac{3}{2}$   
to the left

## Question

Let  $f(x) = -3 \sin\left(\frac{\pi x}{4} - \frac{1}{3}\right) + 1$ . Then which of the following is true regarding the phase shift.

(a) The phase shift is  $\frac{4}{3\pi}$ , the graph is shifted to the left.

$$c = \frac{1}{3}$$

$$\frac{|c|}{b} = \frac{\frac{1}{3}}{\pi/4} = \frac{4}{3\pi}$$

(b) The phase shift is  $\frac{4}{3\pi}$ , the graph is shifted to the right.

(c) The phase shift is  $\frac{3\pi}{4}$ , the graph is shifted to the left.


(d) The phase shift is  $\frac{3\pi}{4}$ , the graph is shifted to the right.



## Vertical Shift

Consider:  $f(x) = a \sin(bx - c) + d$  or  $f(x) = a \cos(bx - c) + d$

**Definition:** If  $d$  is a nonzero number, then the function  $f$  has a **vertical shift** of  $|d|$  units **up** if  $d > 0$  and **down** if  $d < 0$ .



## Example

Identify the vertical shift of each function. Determine if it is up or down.

(a)  $f(x) = 3 \sin(4x-2)+1$

V. shift 1 unit up

$$d = 1$$

(b)  $f(x) = -5 \sin\left(\frac{\pi x}{2}\right) - 7$

V. Shift 7 units down


$$d = -7$$

(c)  $f(x) = 2 - 6 \cos(2x+3)$

V. shift 2 units up


$$d = 2$$

## Parent Plots



The period can be divided into four equal segments.  
For the sine function  $x\text{-int} \rightarrow \text{max} \rightarrow x\text{-int} \rightarrow \text{min} \rightarrow x\text{-int}$

## Parent Plots



The period can be divided into four equal segments.

For the cosine function  $\text{max} \rightarrow \text{x-int} \rightarrow \text{min} \rightarrow \text{x-int} \rightarrow \text{max}$  .

## Pulling it all Together!

Plot two full periods of the function  $f(x) = a \sin(bx - c) + d$  (or  $f(x) = a \cos(bx - c) + d$ ). Carry out each of the following steps:

- ▶ Identify the amplitude and determine if there is an  $x$ -axis reflection.
- ▶ Identify the period. Find the length of one fourth of the period.
- ▶ Identify any phase shift with its direction. Identify end points and points that divide the period into four equal parts.
- ▶ Identify any vertical shift with its direction.
- ▶ Use the basic plot of  $y = \sin x$  or  $y = \cos x$  to get the profile.

$$f(x) = 2 - 4 \cos\left(\pi x - \frac{\pi}{2}\right)$$

Identify the amplitude and vertical shift. Find the maximum and minimum values and determine if there is a reflection in the horizontal.

Amplitude  $A = |-4| = 4$

V. Shift 2 units up

max value  $d + |a| = 2 + 4 = 6$

min value  $d - |a| = 2 - 4 = -2$

There is a reflection in the horizontal

\* features will be min - int - max - int - min

$$f(x) = 2 - 4 \cos\left(\pi x - \frac{\pi}{2}\right)$$

Find the period and phase shift with direction.

$$\text{Period} \quad \frac{2\pi}{b} = \frac{2\pi}{\pi} = 2$$

$$\text{Phase shift} \quad \frac{|c|}{b} = \frac{\pi/2}{\pi} = \frac{1}{2} \quad \text{shift to the right}$$

$$f(x) = 2 - 4 \cos\left(\pi x - \frac{\pi}{2}\right)$$

Identify the beginning and end of one period, and divide it into fourths to determine where major points on the graph are (max/mins and translated x-intercepts)

We have a phase shift  $\frac{1}{2}$  unit to the right and a period of 2. A period should start @  $\frac{1}{2}$  and end @  $\frac{1}{2} + 2 = \frac{5}{2}$ .  $\cos(x)$  has a period starting @ 0 and ending @  $2\pi$ .

$$\pi x - \frac{\pi}{2} = 0 \Rightarrow \pi x = \frac{\pi}{2} \Rightarrow x = \frac{1}{2}$$

period start

$$\pi x - \frac{\pi}{2} = 2\pi \Rightarrow \pi x = \frac{5\pi}{2} \Rightarrow x = \frac{5}{2}$$

a period ends

$$\frac{1}{4} \text{ period} = \frac{1}{4} \cdot 2 = \frac{1}{2}$$

key points (x-values)

$\frac{1}{2}$   
min

1  
int

$\frac{3}{2}$   
max


2  
int

$\frac{5}{2}$   
min



$$f(x) = 2 - 4 \cos\left(\pi x - \frac{\pi}{2}\right)$$

Plot two periods of its graph.




## Section 7.4: Inverse Trigonometric Functions

**Question:** If someone asks "what is the sine of  $\frac{\pi}{6}$ ?" we can respond with the answer (from memory or perhaps using a calculator) " $\frac{1}{2}$ ". What if the question is reversed? What if someone asks

"What angle has a sine value of  $\frac{1}{2}$ ?"

An answer is  $\frac{\pi}{6}$ ; another is  $\frac{5\pi}{6}$ .

## Restricting the Domain of $\sin(x)$



**Figure:** To define an inverse sine function, we start by restricting the domain of  $\sin(x)$  to the interval  $[-\frac{\pi}{2}, \frac{\pi}{2}]$

# The Inverse Sine Function (a.k.a. arcsine function)

**Definition:** For  $x$  in the interval  $[-1, 1]$  the inverse sine of  $x$  is denoted by either

$$\sin^{-1}(x) \quad \text{or} \quad \arcsin(x)$$

and is defined by the relationship

$$y = \sin^{-1}(x) \iff x = \sin(y) \quad \text{where} \quad -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}.$$

**The Domain of the Inverse Sine is**  $-1 \leq x \leq 1$ .

**The Range of the Inverse Sine is**  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .

## Notation Warning!

**Caution:** We must remember not to confuse the superscript  $-1$  notation with reciprocal. That is

$$\sin^{-1}(x) \neq \frac{1}{\sin(x)}.$$

If we want to indicate a reciprocal, we should use parentheses or trigonometric identities

$$\frac{1}{\sin(x)} = (\sin(x))^{-1} \quad \text{or write} \quad \frac{1}{\sin(x)} = \csc(x).$$