

Nov. 28 Math 1190 sec. 52 Fall 2016

## Section 5.4: Properties of the Definite Integral

Suppose that  $f$  and  $g$  are integrable on  $[a, b]$  and let  $k$  be constant.

$$\text{I. } \int_a^b kf(x) dx = k \int_a^b f(x) dx$$

$$\text{II. } \int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$$

$$\text{II. } \int_a^b (f(x) - g(x)) dx = \int_a^b f(x) dx - \int_a^b g(x) dx$$

## The Sum/Difference in General

If  $f_1, f_2, \dots, f_n$  are integrable on  $[a, b]$  and  $k_1, k_2, \dots, k_n$  are constants, then

$$\int_a^b [k_1 f_1(x) + k_2 f_2(x) + \dots + k_n f_n(x)] dx =$$

$$k_1 \int_a^b f_1(x) dx + k_2 \int_a^b f_2(x) dx + \dots + k_n \int_a^b f_n(x) dx$$

## Properties of Definite Integrals Continued...

If  $f$  is integrable on any interval containing the numbers  $a$ ,  $b$ , and  $c$ , then

$$(IV) \quad \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

## Question

Suppose

$$\int_0^5 f(x) dx = 3, \quad \text{and} \quad \int_0^7 f(x) dx = 1.$$

Evaluate  $\int_5^7 f(x) dx$

$$\int_0^7 f(x) dx = \int_0^5 f(x) dx + \int_5^7 f(x) dx$$

$$1 = 3 + \int_5^7 f(x) dx$$

(a) 2

(b) -2

(c) 4

(d) Can't be determined without more information.

## Properties: Bounds on Integrals

(V) If  $f(x) \leq g(x)$  for  $a \leq x \leq b$ , then 
$$\int_a^b f(x) dx \leq \int_a^b g(x) dx$$


If  $f(x) > 0$  on  $[a, b]$  then 
$$\int_a^b f(x) dx > 0$$

(VI) And, as an immediate consequence of (V), if  $m \leq f(x) \leq M$  for  $a \leq x \leq b$ , then

$$m(b - a) \leq \int_a^b f(x) dx \leq M(b - a).$$

If  $f$  is continuous on  $[a, b]$ , we can take  $m$  to be the absolute minimum value and  $M$  the absolute maximum value of  $f$  as guaranteed by the Extreme Value Theorem.

## Bounding: For nonnegative function



## Average Value of a Function and the Mean Value Theorem

**Definition:** Let  $f$  be continuous on the closed interval  $[a, b]$ . Then the average value of  $f$  on  $[a, b]$  is

$$f_{avg} = \frac{1}{b-a} \int_a^b f(x) dx.$$

**Theorem:** (The Mean Value Theorem for Integrals) If  $f$  is continuous on the interval  $[a, b]$ , then there exists a number  $u$  in  $[a, b]$  such that

$$f(u) = f_{avg}, \quad \text{i.e.} \quad \int_a^b f(x) dx = f(u)(b-a).$$

## Question

Find the average value of  $f(x) = \sqrt{x}$  on the interval  $[0, 4]$ . That is, compute

$$f_{avg} = \frac{1}{4 - 0} \int_0^4 x^{1/2} dx$$

(a)  $\frac{16}{3}$

(b)  $\frac{4}{3}$

(c)  $\frac{1}{2}$

(d) 2

$$= \frac{1}{4} \left[ \frac{x^{3/2}}{3/2} \right]_0^4$$

$$= \frac{1}{4} \left( \frac{2}{3} 4^{3/2} - \frac{2}{3} 0^{3/2} \right)$$

$$= \frac{1}{4} \cdot \frac{2}{3} \cdot 8 = \frac{4}{3}$$



## Question

Find the value of  $f$  guaranteed by the MVT for integrals for  $f(x) = \sqrt{x}$  on the interval  $[0, 4]$ . That is, find  $u$  such that

$$f(u) = f_{avg} = \frac{1}{4} \int_0^4 x^{1/2} dx = \frac{4}{3}$$

(a)  $\sqrt{\frac{4}{3}}$

$$f(u) = \frac{4}{3}$$


(b)  $\frac{2}{\sqrt{3}}$

$$\sqrt{u} = \frac{4}{3} \Rightarrow u = \left(\frac{4}{3}\right)^2 = \frac{16}{9}$$

(c)  $\frac{16}{9}$

(d)  $\frac{16}{3}$

## MVT for Integrals Example



**Figure:** The area under the red curve is equal to the area in the rectangle with the blue top.

## Antiderivatives: $e^{ax}$ and $a^x$

If  $a$  is a nonzero constant

$$\frac{d}{dx} \left( \frac{1}{a} e^{ax} \right) = e^{ax}.$$

Hence  $\frac{1}{a} e^{ax}$  is an antiderivative of  $e^{ax}$ .

If  $a > 0$  and  $a \neq 1$ , then

$$\frac{d}{dx} \left( \frac{1}{\ln a} a^x \right) = a^x.$$

Hence  $\frac{1}{\ln a} a^x$  is an antiderivative of  $a^x$ .

## Evaluate Each Integral

$$(a) \int_1^2 (e^{2t} + 2^t) dt$$

$$= \frac{1}{2} e^{2t} + \frac{1}{\ln 2} 2^t \Big|_1^2$$

$$= \frac{1}{2} e^{2 \cdot 2} + \frac{1}{\ln 2} 2^2 - \left( \frac{1}{2} e^{2 \cdot 1} + \frac{1}{\ln 2} 2^1 \right)$$

$$= \frac{1}{2} e^4 + \frac{4}{\ln 2} - \frac{1}{2} e^2 - \frac{2}{\ln 2} = \frac{1}{2} e^4 + \frac{2}{\ln 2} - \frac{1}{2} e^2$$

$$(b) \int_1^0 (2x-1)^2 dx$$

$$(2x-1)^2 = 4x^2 - 4x + 1$$

$$= \int_1^0 (4x^2 - 4x + 1) dx$$

$$= 4 \frac{x^3}{3} - 4 \frac{x^2}{2} + x \Big|_1^0$$

$$= \frac{4}{3} \cdot 0^3 - 2 \cdot 0^2 + 0 - \left( \frac{4}{3} \cdot 1^3 - 2 \cdot 1^2 + 1 \right)$$

$$= 0 - \left( \frac{4}{3} - 2 + 1 \right) = -\frac{1}{3}$$

## Question

$$(c) \int_0^2 x^2(4x+6) dx = \int_0^2 (4x^3 + 6x^2) dx$$

$$(a) \quad 32 = 4 \cdot \frac{x^4}{4} + 6 \cdot \frac{x^3}{3} \Big|_0^2$$

$$(b) \quad \frac{160}{3} = x^4 + 2x^3 \Big|_0^2 = 2^4 + 2 \cdot 2^3 - (0^4 + 2 \cdot 0^3)$$

$$(c) \quad \frac{68}{3}$$

$$= 16 + 16 = 32$$

$$(d) \quad 68$$

$$(d) \int_0^{\pi/4} \tan^2 \theta \, d\theta$$

Recall that

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$\tan^2 \theta = \sec^2 \theta - 1$$

$$= \int_0^{\pi/4} (\sec^2 \theta - 1) \, d\theta$$

$$= \tan \theta - \theta \Big|_0^{\pi/4}$$

$$= \tan \pi/4 - \pi/4 - (\tan 0 - 0) = 1 - \pi/4$$

## Question

$$(e) \int_{\pi/4}^{\pi/2} \frac{dx}{\sin^2 x} = \int_{\pi/4}^{\pi/2} \csc^2 x \, dx = -\cot x \Big|_{\pi/4}^{\pi/2}$$

(a) -1

$$= -\cot \pi/2 - (-\cot \pi/4)$$

(b) 1

$$= 0 - (-1) = 1$$

(c) 0

(d) This can't be evaluated without more advanced techniques.



$$(f) \int_{-6}^{-2} \frac{x^{1/3} + 4x^3 + 2}{x} dx$$

$$= \int_{-6}^{-2} \left( \frac{x^{1/3}}{x} + \frac{4x^3}{x} + \frac{2}{x} \right) dx$$

$$= \int_{-6}^{-2} \left( x^{-2/3} + 4x^2 + \frac{2}{x} \right) dx$$

$$\frac{x^{1/3}}{1/3} + 4 \frac{x^3}{3} + 2 \ln|x| \Big|_{-6}^{-2}$$

$$3(-2)^{\frac{1}{3}} + \frac{4}{3}(-2)^3 + 2\ln|-2| -$$

$$\left( 3(-6)^{\frac{1}{3}} + \frac{4}{3}(-6)^3 + 2\ln|-6| \right)$$

$$= 3\sqrt[3]{-2} + \frac{4}{3}(-8) + 2\ln 2 - \left( 3\sqrt[3]{-6} - \frac{4}{3}(216) + 2\ln 6 \right)$$

$$= -3\sqrt[3]{2} - \frac{32}{3} + 2\ln 2 - (-3\sqrt[3]{6}) + \frac{864}{3} - 2\ln 6$$

$$= 3\sqrt[3]{6} - 3\sqrt[3]{2} + \frac{832}{3} + 2\ln 2 - 2\ln 6$$