

November 9 MATH 1113 sec. 51 Fall 2018

Section 6.5: Trigonometric Functions of a Real Variable and Their Graphs




Figure: It's worth noting that major features (maximum, minimum, x-intercepts) divide the period into four equal parts.

Plot of $y = f(x) = \sin x$




Figure: $y = \sin x$ over four full periods. Note the periodicity and the odd symmetry.

Plot of $y = f(x) = \cos x$




Figure: $y = \cos x$ over one full period.

Plot of $y = f(x) = \cos x$




Figure: $y = \cos x$ over four full periods. Note the periodicity and the even symmetry

Plot of $f(x) = \cos x$ and $g(x) = \sin x$ Together




Figure: Note that the cofunction property $\cos(x) = \sin(\pi/2 - x)$ can be seen in the graphs.

A word of caution on using the unit circle

Our goal is to know and understand the properties of the trigonometric functions as **real valued functions of real variables**.

- ▶ They appear in many models of periodic processes (sound/water waves, seasonal processes, predator-prey interactions) that have no connection to triangles or circles.
- ▶ Right triangles and the unit circle provide powerful mnemonic devices.
- ▶ You'll often see x used as the independent variable—in this class and in Calculus to be sure! The point $(x, y) = (\cos s, \sin s)$ on the unit circle are not the same (x, y) as in $y = \cos x$. Get used to it!

Summary of Sine and Cosine Properties

Let $f(x) = \sin x$ and $g(x) = \cos x$. Then

- ▶ Both f and g are periodic with fundamental period 2π .
- ▶ The domain of both f and g is $(-\infty, \infty)$.
- ▶ The range of both f and g is $[-1, 1]$; they have amplitude 1.
- ▶ f is an odd function, i.e. $f(-x) = \sin(-x) = -\sin x = -f(x)$.
- ▶ g is an even function, i.e. $g(-x) = \cos(-x) = \cos x = g(x)$.
- ▶ Both f and g are continuous on their entire domain $(-\infty, \infty)$.
- ▶ The zeros of f are integer multiples of π , i.e. $f(n\pi) = 0$ for $n = 0, \pm 1, \pm 2, \dots$
- ▶ The zeros of g are odd integer multiples of $\pi/2$, $g\left(\frac{m\pi}{2}\right) = 0$ for $m = \pm 1, \pm 3, \dots$

Question

Let $f(x) = \sin x$, and suppose that $f(a) = -0.62$ for some real number a . Which of the following is true?

(a) $f(a + 2\pi) = -0.62$ ✓

$\sin(x + 2\pi) = \sin x$ for all real x

(b) $f(2a) = -1.24$ ← outside of the range

(c) $f(-a) = 0.62$ ✓

$\sin(-x) = -\sin(x)$ for all real x

(d) Only (a) and (b) are true.

(e) Only (a) and (c) are true.

Question

True or False: The equation $\cos x = 2$ has no real solutions.

- (a) True, and I'm confident.
- (b) True, but I'm not confident.
- (c) False, and I'm confident.
- (d) False, but I'm not confident.

The Tangent

The function $\tan s = \frac{\sin s}{\cos s}$. Recall that

$$\cos s = 0 \quad \text{whenever} \quad s = \frac{m\pi}{2} \quad \text{for} \quad m = \pm 1, \pm 3, \pm 5, \dots$$

When $\cos s = 0$, $\sin s$ is either 1 or -1 . Hence

Domain: The domain of the tangent function is all real number **except** odd multiples of $\pi/2$. We can write this as

$$\left\{ s \mid s \neq \frac{\pi}{2} + k\pi, k = 0, \pm 1, \pm 2, \dots \right\}$$

Moreover, the graph of the tangent function has vertical asymptotes at each odd multiple of $\pi/2$.

The Tangent

Range: The range of the tangent function is **all real numbers**.

Symmetry: The function $f(s) = \tan s$ is odd. That is

$$f(-s) = \tan(-s) = -\tan s = -f(s).$$

Periodicity: The tangent function is periodic with fundamental period π . That is

$$\tan(s + \pi) = \tan s \quad \text{for all } s \text{ in the domain.}$$

Note: The period of the tangent function is π . This is different from the period of the sine and cosine.

The Tangent

A few key tangent values:

s	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
$\tan s$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	undef.

And due to symmetry

s	$-\frac{\pi}{2}$	$-\frac{\pi}{3}$	$-\frac{\pi}{4}$	$-\frac{\pi}{6}$	0
$\tan s$	undef.	$-\sqrt{3}$	-1	$-\frac{1}{\sqrt{3}}$	0

Here is an applet to plot two periods of the function $f(s) = \tan s$.

GeoGebra Graph Applet: Sine, Cosine, and Tangent

Basic Plot $f(x) = \tan x$




Figure: Plot of several periods of $f(x) = \tan x$. Note that the interval between adjacent asymptotes is the period π .

Cotangent: Using the Cofunction ID and Symmetry

$$\cot s = \tan\left(\frac{\pi}{2} - s\right) = \tan\left(-\left(s - \frac{\pi}{2}\right)\right) = -\tan\left(s - \frac{\pi}{2}\right).$$

So the graph of $f(s) = \cot s$ is the graph of $g(s) = \tan s$ under a horizontal shift $\pi/2$ units to the right followed by a reflection in the s -axis.




Figure: Plot of $f(x) = \cot x$. Note that the lines $x = n\pi$ for $n = 0, \pm 1, \pm 2, \dots$ are vertical asymptotes to the graph. The dashed curve is $y = \tan x$.

Cosecant and Secant

Domains: Since $\sin(n\pi) = 0$ for integers n ,

$$\text{Domain}(\csc s) = \{s \mid s \neq n\pi, \text{ for integers } n\}.$$

Since $\cos\left(\frac{\pi}{2} + n\pi\right) = 0$ for integers n , the domain of $\sec s$ is

$$\text{Domain}(\sec s) = \left\{s \mid s \neq \frac{\pi}{2} + n\pi, \text{ for integers } n\right\}.$$

Ranges: Note that

$$|\csc s| = \frac{1}{|\sin s|} \geq 1 \quad \text{and} \quad |\sec s| = \frac{1}{|\cos s|} \geq 1$$

so the range of both $\csc s$ and $\sec s$ is

$$(-\infty, -1] \cup [1, \infty).$$

Cosecant: Using $\csc s = \frac{1}{\sin s}$




Figure: Two periods of $f(s) = \csc s$. The dashed curve is $y = \sin s$. Note the asymptotes $s = n\pi$ for integers n where $\sin s$ takes its zeros. The curves meet at the relative extrema and have the same period 2π .

Secant: Using $\sec s = \frac{1}{\cos s}$




Figure: Two periods of $f(s) = \sec s$. The dashed curve is $y = \cos s$. Note the asymptotes $s = \pi/2 + n\pi$ for integers n where $\cos s$ takes its zeros. The curves meet at the relative extrema and have the same period 2π .

Question

Recall that the **cosine is an even function** and the **sine is an odd function**. Moreover,

$$\sec x = \frac{1}{\cos x} \quad \text{and} \quad \csc x = \frac{1}{\sin x}$$

Which of the following statements is true about the symmetry of the secant and cosecant.

- (a) Both of the secant and cosecant are even functions.
- (b) Both the secant and cosecant are odd functions.
- (c) Secant is even, and cosecant is odd.
- (d) Secant is odd, and cosecant is even.
- (e) Symmetry for secant and cosecant can't be deduced from symmetry of cosine and sine.

Section 6.6: Graphing Trigonometric Functions with Transformations

Our goal is to graph functions of the form

$$f(x) = a \sin(bx - c) + d \quad \text{or} \quad f(x) = a \cos(bx - c) + d$$

Note: here we will be graphing points (x, y) on a curve $y = f(x)$.

Amplitude


Consider: $f(x) = a \sin(bx - c) + d$ or $f(x) = a \cos(bx - c) + d$

Definition: Let a be any nonzero real number. The **amplitude** of the function f defined above is the value $|a|$.


Recall that this is half the distance between the maximum and minimum values.

If $a < 0$ the graph is reflected in the x -axis. But the amplitude is still $|a|$.


Amplitude




$$f(x) = a \sin x$$



..... $y = \sin(x)$ — $y = 2\sin(x)$ — $y = \frac{1}{2}\sin(x)$



$$f(x) = a \cos x$$



..... $y = \cos(x)$ — $y = 2\cos(x)$ — $y = \frac{1}{2}\cos(x)$

Example


Identify the amplitude A of each function. Determine if the graph is reflected in the x -axis.

(a) $f(x) = 3 \sin(4x-2)+1$ $A = 3$ not reflected
3 is positive

(b) $f(x) = -5 \sin\left(\frac{\pi x}{2}\right)+7$ $A = 5$ yes reflected as
-5 is negative

(c) $f(x) = 2-6 \cos(2x+3)$ $A = 6$ yes reflected as
-6 is negative
 $= -6 \cos(2x+3) + 2$

Question



The figure shows two periods of the plot of $y = f(x)$ where

(a) $f(x) = 4 \sin x$

(b) $f(x) = -4 \sin x$

(c) $f(x) = 4 \cos x$

(d) $f(x) = -4 \cos x$