

### Review for Exam 3

#### MATH 2306 (Ritter)

Sections Covered: 8, 9, 10, 11, 12, 13, 14

*This review is provided as a courtesy to give some idea of what material is covered. Nothing else is intended or implied.*

(1) Solve each IVP

(a)  $y'' - 3y' + 2y = 0$   $y(0) = 0$ ,  $y'(0) = 2$

(b)  $y'' + 2y' = 0$   $y(1) = 0$ ,  $y'(1) = 1$

(c)  $y'' - 2y' + 5y = 0$   $y(0) = 0$ ,  $y'(0) = 2$

(2) Find the general solution of each nonhomogeneous equation

(a)  $y'' + 6y' + 9y = e^x + 3e^{-3x}$

(b)  $y'' + y' - 12y = 2x$

(c)  $y'' + y = 4 \cos x$

(3) Determine the **form** of the particular solution. (Do not bother trying to find any of the coefficients  $A$ ,  $B$ , etc.)

(a)  $y'' - 4y' + 5y = x \cos 2x$

(b)  $y'' + y = x^3 + e^x$

(c)  $y'' - 4y' + 5y = xe^{2x} \sin x$

(d)  $y'' - 2y' + y = 1 + e^x$

(4) For each homogeneous equation, write out the characteristic equation. If the equation doesn't have a characteristic equation, briefly state why.

(a)  $3\frac{d^4y}{dx^4} - 2\frac{d^3y}{dx^3} + \frac{dy}{dx} - 4y = 0$

(b)  $4y'' + 2xy' + e^x y = 0$

(c)  $x^3 y''' + 2x^2 y'' - 4xy' + y = 0$

(d)  $y^{(6)} + 16y^{(4)} - 12y'' + y = 0$

(5) Consider the nonhomogeneous equation  $x^2 y'' + xy' - y = x$ . In standard form, this equation is  $y'' + \frac{1}{x}y' - \frac{1}{x^2}y = x^{-1}$ .

(a) Try using the method of undetermined coefficients to find a particular solution of the form

$$y_p = Ax + B.$$

(b) Maybe the form was wrong. Try it again with  $y_p = Ax^{-1}$ .

(c) Why is this approach failing?

(d) It can be shown that  $y_c = c_1x + c_2x^{-1}$ . Use an appropriate method to find  $y_p$ .

(6) A certain spring is 1 ft long with no mass attached. An object weighing 10 lbs is attached to the spring. The length of the spring with the mass attached is then 18 inches.

(a) Compute the mass  $m$  in slugs and the spring constant  $k$  in lbs/ft.

(b) If the object is initially at equilibrium and given a downward velocity of 1 ft/sec, find the displacement for  $t > 0$ .

(c) Next assume that a driving force of  $f(t) = \cos(\gamma t)$  is applied to the object. What value of  $\gamma$  will result in pure resonance?

(d) Let  $f(t) = \cos(3t)$ . Determine the displacement for  $t > 0$  assuming the object started from rest at equilibrium .

(7) A 64 lb object is attached to a spring whose spring constant is 26 lb/ft. A dashpot provides damping that is numerically equal to 8 times the instantaneous velocity.

(a) Determine the mass  $m$  of the object in slugs.

(b) Assuming there is no external applied force, set up the differential equation for the displacement and determine if the motion is overdamped, underdamped or critically damped.

(c) If the object is initially displaced 6 inches above equilibrium and given an initial upward velocity of 2 ft/sec, determine the displacement for  $t > 0$ .

(8) A 200 volt battery is applied to a series circuit with inductance 2 henries, resistance 26 ohms and capacitance  $\frac{1}{80}$  farads. Find the charge on the capacitor  $q(t)$  for  $t > 0$  assuming the initial charge and current are zero,  $q(0) = 0$ ,  $i(0) = 0$ .

(9) Consider the nonhomogeneous equation  $x^2y'' + xy' - 4y = 20x^3$ .

(a) One solution of the associated homogeneous equation is  $y_1 = x^2$ . Find a second linearly independent one  $y_2$ .

(b) Find a particular solution  $y_p$  of the nonhomogeneous equation.

(c) Solve the IVP:  $x^2y'' + xy' - 4y = 20x^3$ ,  $y(1) = 3$ ,  $y'(1) = 6$ .

(10) Use the method of variation of parameters to find a particular solution for each nonhomogeneous equation.

(a)  $y'' + y = \sec \theta \tan \theta$ ,  $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$

(b)  $y'' + 3y' + 2y = \sin(e^x)$

(11) Use the definition (i.e. compute an integral) to show that  $\mathcal{L}\{e^{at}\} = \frac{1}{s-a}$  for  $s > a$ .

(12) Compute the transform or inverse transform as indicated. (Use the table of Laplace transforms along with any necessary algebra or identities.)

(a)  $\mathcal{L}\{(2t-3)^2\}$

(b)  $\mathcal{L}\{\cos^2 t - \sin^2 t\}$  (hint: double angle formula)

(c)  $\mathcal{L}\{e^{3t} + 7 \sin(2t) - t^5\}$

(d)  $\mathcal{L}^{-1}\left\{\frac{1}{s^4} - \frac{s}{s^2 + 5}\right\}$

(e)  $\mathcal{L}^{-1}\left\{\frac{5s + 3}{s^2 + s}\right\}$