

Exam I Math 3260 sec. 57

Fall 2017

Name: _____ *Solutions*

Your signature (required) confirms that you agree to practice academic honesty.

Signature: _____

Problem	Points
1	
2	
3	
4	
5	
6	
7	
8	

INSTRUCTIONS: There are 8 problems; the point values are listed with the problems. You may use a calculator with matrix capabilities. **No wifi enabled device can be used as a calculator.** There are no notes, or books allowed. **Illicit use of a smart phone, tablet, device that runs apps, or hand written notes will result in a grade of zero on this exam as well as a formal allegation of academic misconduct.** To receive full credit, you must clearly justify your answer.

(1) (15 points) Solve the linear system by using row reduction on the associated augmented matrix. **Show each step in the row reduction process—i.e. do this one by hand and show all of your work.** Give your answer in parametric or vector parametric form, your choice.

$$\begin{array}{rrcr} x_1 & & -x_3 & = & 2 \\ 2x_1 & + & x_2 & + & 2x_3 & = & -6 \\ 3x_1 & + & 2x_2 & + & 2x_3 & = & -5 \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 2 \\ 2 & 1 & 2 & -6 \\ 3 & 2 & 2 & -5 \end{array} \right] \quad \begin{array}{l} -2R_1 + R_2 \rightarrow R_2 \\ -3R_1 + R_3 \rightarrow R_3 \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 2 \\ 0 & 1 & 4 & -10 \\ 0 & 2 & 5 & -11 \end{array} \right]$$

$$-2R_2 + R_3 \rightarrow R_3$$

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 2 \\ 0 & 1 & 4 & -10 \\ 0 & 0 & -3 & 9 \end{array} \right]$$

$$-\frac{1}{3}R_3 \rightarrow R_3$$

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 2 \\ 0 & 1 & 4 & -10 \\ 0 & 0 & 1 & -3 \end{array} \right]$$

$$-4R_3 + R_2 \rightarrow R_2$$

$$R_3 + R_1 \rightarrow R_1$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -3 \end{array} \right]$$

\Rightarrow

$$\begin{array}{l} x_1 = -1 \\ x_2 = 2 \\ x_3 = -3 \end{array}$$

In vector parametric

$$\vec{x} = \begin{bmatrix} -1 \\ 2 \\ -3 \end{bmatrix}$$

(2) (15 points) The matrices A and B shown below are row equivalent. Use this to find the solution set of the given system of equations.

$$A = \begin{bmatrix} 3 & -6 & 1 & 6 & 14 \\ -1 & 2 & 0 & -1 & -4 \\ 2 & -4 & 1 & 5 & 10 \end{bmatrix} \quad B = \begin{bmatrix} 1 & -2 & 0 & 1 & 4 \\ 0 & 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{aligned} 3x_1 - 6x_2 + x_3 + 6x_4 &= 14 \\ -x_1 + 2x_2 - x_4 &= -4 \\ 2x_1 - 4x_2 + x_3 + 5x_4 &= 10 \end{aligned}$$

From B

$$x_1 = 4 + 2x_2 - x_4$$

x_2 - free

$$x_3 = 2 - 3x_4$$

x_4 - free

In vector parametric

$$\vec{x} = \begin{bmatrix} 4 \\ 0 \\ 2 \\ 0 \end{bmatrix} + t \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -1 \\ 0 \\ -3 \\ 1 \end{bmatrix} \quad \text{s.t. in } \mathbb{R}$$

(3) (15 points) Find the standard matrix for the linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ defined by

$$T(x_1, x_2, x_3) = (-x_3, 0, x_2, x_1)$$

$$T(1, 0, 0) = (0, 0, 0, 1)$$

$$T(0, 1, 0) = (0, 0, 1, 0)$$

$$T(0, 0, 1) = (-1, 0, 0, 0)$$

The standard matrix is

$$\begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

(4) (5 points) Suppose the homogeneous equation $Ax = \mathbf{0}$ has the two nontrivial solutions \mathbf{u} and \mathbf{v} . Show that every vector in $\text{Span}\{\mathbf{u}, \mathbf{v}\}$ also solves the homogeneous equation.

$$A\vec{u} = \vec{0} \quad \text{and} \quad A\vec{v} = \vec{0}$$

Let $\vec{x} \in \text{span}\{\vec{u}, \vec{v}\}$. So $\vec{x} = c_1\vec{u} + c_2\vec{v}$ for some numbers c_1 and c_2 .

$$\begin{aligned} A\vec{x} &= A(c_1\vec{u} + c_2\vec{v}) = c_1 A\vec{u} + c_2 A\vec{v} \\ &= c_1\vec{0} + c_2\vec{0} = \vec{0} \end{aligned}$$

Hence \vec{x} in $\text{span}\{\vec{u}, \vec{v}\}$ solves the homogeneous equation.

(5) (5 points) Suppose that the vector \mathbf{u}_1 solves the nonhomogeneous equation $Ax = \mathbf{b}_1$ and that the vector \mathbf{u}_2 solves the nonhomogeneous equation $Ax = \mathbf{b}_2$. Show that the vector $\mathbf{u}_1 + \mathbf{u}_2$ solves the nonhomogeneous equation $Ax = \mathbf{b}_1 + \mathbf{b}_2$.

Directly from the properties of matrix multiplication

$$\begin{aligned} A(\vec{u}_1 + \vec{u}_2) &= A\vec{u}_1 + A\vec{u}_2 \\ &= \vec{b}_1 + \vec{b}_2 \quad \text{as expected,} \end{aligned}$$

(6) (15 points) Consider the set of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

$$\mathbf{v}_1 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix}$$

(a) Is the set $\{\mathbf{v}_1, \mathbf{v}_2\}$ linearly dependent or linearly independent? (Justify)

They are dependent, in particular

$$\vec{v}_2 = 2\vec{v}_1$$

$$\text{i.e. } 2\vec{v}_1 - \vec{v}_2 = \vec{0}$$

(b) Is the vector $\mathbf{b} = (1, 0, 0)$ in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$? (Justify)

No. If $\vec{b} = c_1\vec{v}_1 + c_2\vec{v}_2$ then

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} c_1 \cdot 0 + c_2 \cdot 0 \\ : \\ : \end{bmatrix} \quad \text{which requires} \\ 1 = 0.$$

(c) Is the set $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ linearly dependent or linearly independent? (Justify)

It's dependent by part (a).

$$\text{Note } 2\vec{v}_1 - \vec{v}_2 + 0\vec{v}_3 = \vec{0}$$

Is a lin. dependence relation.

(7) (20 points) Use the given matrices to evaluate each expression. If an expression does not exist, state why it does not exist.

$$A = \begin{bmatrix} 1 & -1 & 4 \\ 0 & 2 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 0 \\ 2 & -3 \end{bmatrix}, \quad C = \begin{bmatrix} 2 & 2 \\ 1 & -3 \\ 0 & 4 \end{bmatrix}$$

(a) $-2B = \begin{bmatrix} -2 & 0 \\ -4 & 6 \end{bmatrix}$

(b) AB
 $2 \times 3 \quad 2 \times 2$
 Undefined. \cdot # rows in B doesn't match # columns in A

(c) $BA = \begin{bmatrix} 1 & 0 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} 1 & -1 & 4 \\ 0 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -1 & 4 \\ 2 & -8 & 5 \end{bmatrix}$

(d) Ax where $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$
 Undefined Ax requires \vec{x} in \mathbb{R}^3

(e) $C^T + A = \begin{bmatrix} 2 & 1 & 0 \\ 2 & -3 & 4 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 4 \\ 0 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 4 \\ 2 & -1 & 5 \end{bmatrix}$

(8) (10 points) The homogeneous equation $Ax = 0$ is always consistent because it always has the trivial solution. Let

$$A = \begin{bmatrix} 1 & b \\ c & d \end{bmatrix}, \quad \text{for some real numbers } b, c, \text{ and } d.$$

Determine whether the homogeneous equation $Ax = 0$ has one solution or infinitely many solutions

(a) If $d = bc$.

$$\begin{bmatrix} 1 & b & 0 \\ c & d & 0 \end{bmatrix} \quad -cR_1 + R_2 \rightarrow R_2$$

$$\begin{bmatrix} 1 & b & 0 \\ 0 & d - cb & 0 \end{bmatrix}$$

If $d = bc$
the matrix is $\begin{bmatrix} 1 & b & 0 \\ 0 & 0 & 0 \end{bmatrix}$

The system has infinitely many solutions.

(b) If $d = bc + 1$.

If $d = bc + 1$ the above is

$$\begin{bmatrix} 1 & b & 0 \\ 0 & 1 & 0 \end{bmatrix} \Rightarrow x_1 = x_2 = 0$$

Only the trivial solution would exist.