

Exam III Math 3260 sec. 56

Spring 2018

Name: _____ *Solution* _____

Your signature (required) confirms that you agree to practice academic honesty.

Signature: _____

Problem	Points
1	
2	
3	
4	
5	
6	
7	

INSTRUCTIONS: There are 7 problems; the point values are listed with the problems. You may use a calculator with matrix capabilities. **No wifi enabled device can be used as a calculator.** There are no notes, or books allowed. **Illicit use of a smart phone, tablet, device that runs apps, or hand written notes will result in a grade of zero on this exam as well as a formal allegation of academic misconduct.** To receive full credit, you must clearly justify your answer.

(1) (15 points) Find an **orthonormal** basis for the column space of the matrix $A = \begin{bmatrix} 1 & 4 \\ 2 & -7 \\ 2 & -4 \end{bmatrix}$.

Orthogonal: \vec{v}_1, \vec{v}_2

$$\vec{v}_1 = \vec{u}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$$

$$\vec{v}_2 = \vec{u}_2 - \frac{\vec{u}_2 \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \vec{v}_1 = \begin{bmatrix} 4 \\ -7 \\ -4 \end{bmatrix} - \frac{-18}{9} \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 4 \\ -7 \\ -4 \end{bmatrix} + \begin{bmatrix} 2 \\ 4 \\ 4 \end{bmatrix} = \begin{bmatrix} 6 \\ -3 \\ 0 \end{bmatrix}$$

Orthogonal basis $\left\{ \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 6 \\ -3 \\ 0 \end{bmatrix} \right\}$

$$\|\vec{v}_1\| = \sqrt{9} = 3, \quad \|\vec{v}_2\| = \sqrt{36+9} = \sqrt{45}$$

Orthonormal basis $\left\{ \frac{1}{3} \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \frac{1}{\sqrt{45}} \begin{bmatrix} 6 \\ -3 \\ 0 \end{bmatrix} \right\}$

(2) (10 points) The given set \mathcal{B} is a basis for \mathbb{R}^2 . Determine the change of coordinates matrix $P_{\mathcal{B}}$ and its inverse (use the order presented here). Then use this to find the indicated coordinate vectors.

$$\mathcal{B} = \left\{ \begin{bmatrix} 3 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$$

$$P_{\mathcal{B}} = \begin{bmatrix} 3 & -1 \\ 1 & 1 \end{bmatrix}, \quad \det(P_{\mathcal{B}}) = 4, \quad P_{\mathcal{B}}^{-1} = \frac{1}{4} \begin{bmatrix} 1 & 1 \\ -1 & 3 \end{bmatrix}$$

Determine $[\mathbf{x}]_{\mathcal{B}}$ for (a) $\mathbf{x} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$,

(b) $\mathbf{x} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$

$$\begin{aligned} [\mathbf{x}]_{\mathcal{B}} &= P_{\mathcal{B}}^{-1} \mathbf{x} = \frac{1}{4} \begin{bmatrix} 1 & 1 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \\ &= \frac{1}{4} \begin{bmatrix} 0 \\ -4 \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} [\mathbf{x}]_{\mathcal{B}} &= P_{\mathcal{B}}^{-1} \mathbf{x} = \frac{1}{4} \begin{bmatrix} 1 & 1 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \end{bmatrix} \\ &= \frac{1}{4} \begin{bmatrix} 4 \\ 4 \end{bmatrix} \\ &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} \end{aligned}$$

(3) (15 points) Determine the dimension of each vector space described.

(a) The space \mathbb{P}_5 of polynomials of degree at most 5.

$$\dim(\mathbb{P}_5) = 6$$

(b) The subspace of \mathbb{R}^5 of vectors of the form $(x_1, x_2, 0, x_4, 0)$.

The dimension is 3

(c) The null space of a 8×11 matrix A if $\text{rank}(A) = 5$.

$$\text{nullity} + 5 = 11 \Rightarrow \text{nullity} = 6$$

(4) (10 points) Consider the vector $\mathbf{v} = \begin{bmatrix} 3 \\ 0 \\ 2 \\ 1 \end{bmatrix}$.

(a) Determine the norm $\|\mathbf{v}\|$.

$$\|\mathbf{v}\| = \sqrt{9 + 4 + 1} = \sqrt{14}$$

(b) Find a unit vector parallel to \mathbf{v} . (Call the new vector \mathbf{u} .)

$$\vec{u} = \frac{1}{\|\vec{v}\|} \vec{v} = \begin{bmatrix} \frac{3}{\sqrt{14}} \\ 0 \\ \frac{2}{\sqrt{14}} \\ \frac{1}{\sqrt{14}} \end{bmatrix}$$

(5) (25 points) Find bases for each of $\text{Col}A$, $\text{Row}A$, and $\text{Nul}A$. State the rank and the nullity of A where

$$A = \begin{bmatrix} 1 & 3 & 2 & 7 \\ 2 & 6 & 4 & 14 \\ 2 & 6 & 3 & 4 \end{bmatrix} \xrightarrow{\text{rref}} \begin{bmatrix} 1 & 3 & 0 & -13 \\ 0 & 0 & 1 & 10 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

In $\text{Nul}A$ \vec{x} satisfies
 $\vec{x}_1 = -3x_2 + 13x_4$
 $x_3 = 10x_4$
 x_2, x_4 - free

(Be sure to write any calculator output on this page to show that upon which your conclusions are based.)

The following are bases

$$\text{Nul}A = \text{Span} \left\{ \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 13 \\ 0 \\ 10 \\ 1 \end{bmatrix} \right\}$$

$$\text{Col}A = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 3 \end{bmatrix} \right\}$$

$$\text{Row}A = \text{Span} \left\{ \begin{bmatrix} 1 \\ 3 \\ 0 \\ -13 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 10 \end{bmatrix} \right\}$$

$$\text{rank } A = 2, \quad \text{nullity } A = 2$$

(6) (15 points) Find the orthogonal projection of $\mathbf{y} = \begin{bmatrix} -1 \\ 5 \\ 3 \end{bmatrix}$ onto the line $L = \text{Span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$, and determine the distance from \mathbf{y} to L . (Hint: Write $\mathbf{y} = \hat{\mathbf{y}} + \mathbf{z}$ where $\hat{\mathbf{y}}$ is the orthogonal projection.)

$$\hat{\mathbf{y}} = \frac{\mathbf{y} \cdot \mathbf{u}}{\mathbf{u} \cdot \mathbf{u}} \mathbf{u} = \frac{4}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}$$

The orthogonal projection is $\begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}$.

$$\mathbf{y} - \hat{\mathbf{y}} = \begin{bmatrix} -1 \\ 5 \\ 3 \end{bmatrix} - \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} -3 \\ 3 \\ 3 \end{bmatrix}$$

The distance from \mathbf{y} to L is

$$\left\| \begin{bmatrix} -3 \\ 3 \\ 3 \end{bmatrix} \right\| = \sqrt{9+9+9} = 3\sqrt{3}$$

(7) (10 points) Let $A = \begin{bmatrix} 2 & 1 \\ -2 & 5 \end{bmatrix}$. One of the eigenvalues of A is $\lambda = 4$. Find a basis for the associated eigenspace.

$$A\mathbf{x} = 4\mathbf{x} \Rightarrow \begin{bmatrix} 2-4 & 1 \\ -2 & 5-4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -2 & 1 \\ -2 & 1 \end{bmatrix} \xrightarrow{\text{row}} \begin{bmatrix} 1 & -1/2 \\ 0 & 0 \end{bmatrix} \quad \begin{array}{l} x_1 = \frac{1}{2}x_2 \\ x_2 \text{-free} \end{array}$$

A basis for the eigen space is $\left\{ \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} \right\}$.

The Gram-Schmidt Process: Given a set of basis vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ in some inner product space \mathbf{V} , we can obtain an orthonormal set $\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n\}$ that spans the same subspace. First, we obtain the orthogonal set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ where

$$\begin{aligned}\mathbf{v}_1 &= \mathbf{u}_1 \\ \mathbf{v}_2 &= \mathbf{u}_2 - \frac{\mathbf{u}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 \\ \mathbf{v}_3 &= \mathbf{u}_3 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2 \\ &\vdots \\ \mathbf{v}_n &= \mathbf{u}_n - \sum_{k=1}^{n-1} \frac{\mathbf{u}_n \cdot \mathbf{v}_k}{\mathbf{v}_k \cdot \mathbf{v}_k} \mathbf{v}_k\end{aligned}$$

Finally, we obtain the orthonormal set by dividing each \mathbf{v}_i by its norm. That is, for $i = 1, \dots, n$ set

$$\mathbf{w}_i = \frac{1}{\sqrt{\mathbf{v}_i \cdot \mathbf{v}_i}} \mathbf{v}_i.$$