

Exam 4 Math 1113 sec. 51 Fall 2018

Name: Solutions

Your signature (required) confirms that you agree to practice academic honesty.

Signature: _____

Problem	Points
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
Total	

INSTRUCTIONS: There are 10 problems worth 10 points each. No calculator use is allowed on any part of this exam. There are no notes, or books allowed. **Illicit use of a calculator, smart phone, tablet, device that runs apps, or notes will result in a grade of zero on this exam as well as a formal allegation of academic misconduct.** To receive full credit, answers must be clear, complete, and written using proper notation.

You may assume the following six IDs. Choose the signs with same placements.

$$\begin{aligned} \cos(u \pm v) &= \cos u \cos v \mp \sin u \sin v \\ \sin(u \pm v) &= \sin u \cos v \pm \sin v \cos u \\ \tan(u \pm v) &= \frac{\tan u \pm \tan v}{1 \mp \tan u \tan v} \end{aligned}$$

1. Complete the table of trigonometric values.

$\theta =$	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan \theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	undef.

2. Evaluate each expression exactly.

$$(a) \cos\left(\frac{4\pi}{3}\right) = -\frac{1}{2}$$

$$(b) \tan\left(\frac{5\pi}{4}\right) = 1$$

$$(c) \csc\left(\frac{5\pi}{6}\right) = 2$$

$$(d) \cot\left(-\frac{\pi}{3}\right) = \frac{1}{\sqrt{3}}$$

$$(e) \sec\left(\frac{\pi}{4}\right) = \sqrt{2}$$

3. Complete the table. Use interval notation.

Function	Domain	Range
$y = \sin^{-1} x$	$[-1, 1]$	$[-\pi/2, \pi/2]$
$y = \cos^{-1} x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1} x$	$(-\infty, \infty)$	$(-\pi/2, \pi/2)$

4. Evaluate each expression exactly. Recall that certain formulas have been provided on the first page of this exam.

$$(a) \sin(40^\circ) \cos(50^\circ) + \sin(50^\circ) \cos(40^\circ) = \sin(90^\circ) = 1$$

$$(b) \cos(70^\circ) \cos(25^\circ) + \sin(70^\circ) \sin(25^\circ) = \cos(45^\circ) = \frac{1}{\sqrt{2}}$$

$$(c) \tan^{-1}(\sqrt{3}) = \frac{\pi}{3}$$

$$(d) \cos^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{6}$$

$$(e) \tan\left(\frac{\pi}{12}\right) \quad (\text{hint: } \frac{1}{12} = \frac{1}{3} - \frac{1}{4})$$




$$= \frac{\tan\left(\frac{\pi}{3}\right) - \tan\left(\frac{\pi}{4}\right)}{1 + \tan\frac{\pi}{3} \tan\frac{\pi}{4}} = \frac{\sqrt{3} - 1}{1 + \sqrt{3}}$$

5. Plot at least two full periods of each function on the grids provided.



(a) $f(x) = \sin x$

(b) $g(x) = \cos x$

(c) $h(x) = \tan x$



6. Suppose $\frac{3\pi}{2} < \theta < 2\pi$, and $\cos \theta = \frac{2}{3}$. Also suppose that $0 < \phi < \frac{\pi}{2}$ and $\csc \phi = 4$. Draw a representative diagram for each angle from which trigonometric values can be deduced.



Evaluate each expression exactly. Your answers should be simplified, but it is not necessary to rationalize denominators.

$$(a) \cos(\phi - \theta) = \cos \phi \cos \theta + \sin \phi \sin \theta = \frac{\sqrt{15}}{4} \left(\frac{2}{3}\right) + \frac{1}{4} \left(\frac{-\sqrt{5}}{3}\right) = \frac{2\sqrt{15} - \sqrt{5}}{12}$$

$$(b) \csc(\theta + \phi) = \frac{1}{\sin(\theta + \phi)} = \frac{12}{2 - \sqrt{75}}$$

$$\begin{aligned} \sin(\theta + \phi) &= \sin \theta \cos \phi + \sin \phi \cos \theta = \frac{-\sqrt{5}}{3} \left(\frac{\sqrt{15}}{4}\right) + \left(\frac{1}{4}\right) \left(\frac{2}{3}\right) \\ &= \frac{2 - \sqrt{75}}{12} \end{aligned}$$

7. For each function, identify the amplitude A , period T , phase shift ϕ (with direction), and vertical shift V (with direction). Write *none* if a function does not have a specific characteristic.

$$(a) f(x) = 4 \cos\left(\frac{\pi x}{4}\right) - 2 \quad 2\pi / (\pi/4) = 8$$

$$A = \underline{4}, \quad T = \underline{8}, \quad \phi = \underline{\text{none}}, \quad V = \underline{2 \text{ down}}$$

$$(b) y = 2 - \frac{1}{3} \sin\left(x - \frac{\pi}{4}\right)$$

$$A = \underline{\frac{1}{3}}, \quad T = \underline{2\pi}, \quad \phi = \underline{\pi/4 \text{ right}}, \quad V = \underline{2 \text{ up}}$$

8. The hour hand on a certain clock is 3 inches long. (Provide exact answers with the factor π if necessary.)

(a) Determine the distance traversed by the tip of the hour hand over the course of 7 hours.

Distance $s = r\theta$ where $r = 3\text{in}$ and $\theta = \frac{7}{12} \cdot (2\pi) = \frac{7\pi}{6}$

So $s = 3\text{in} \left(\frac{7\pi}{6}\right) = \frac{7\pi}{2} \text{in}$

(b) Find the area of the sector swept out by the hour hand during this 7 hours.

The area $A = \frac{1}{2}r^2\theta$ for the same r, θ

$A = \frac{1}{2}(3\text{in})^2 \left(\frac{7\pi}{6}\right) = \frac{21\pi}{4} \text{in}^2$

9. Evaluate each expression exactly. Your answers should be justified by a diagram or other demonstration of your process.

(a) $\cos(\tan^{-1} 6)$

$= \frac{1}{\sqrt{37}}$


if $\theta = \tan^{-1} 6, \tan \theta = 6$
 $0 < \theta < \frac{\pi}{2}$



(b) $\cot\left(\sin^{-1} -\frac{1}{5}\right)$

$= -\sqrt{24}$

if $\theta = \sin^{-1}\left(-\frac{1}{5}\right) \sin \theta = -\frac{1}{5}$
 $-\frac{\pi}{2} < \theta < 0$



10. Prove each identity.

(a) $\cos^4 x - \sin^4 x = \cos^2 x - \sin^2 x$

From the left

$$\cos^4 x - \sin^4 x = (\cos^2 x - \sin^2 x)(\cos^2 x + \sin^2 x) \quad \text{Diff of squares}$$

$$= (\cos^2 x - \sin^2 x) \cdot 1$$

$$= \cos^2 x - \sin^2 x \quad \text{as expected}$$

(b) $\frac{1 + \tan \theta}{1 + \cot \theta} = \frac{\sin \theta}{\cos \theta}$

From the left

$$\frac{1 + \tan \theta}{1 + \cot \theta} = \frac{\frac{\cos \theta}{\cos \theta} + \frac{\sin \theta}{\cos \theta}}{\frac{\sin \theta}{\sin \theta} + \frac{\cos \theta}{\sin \theta}}$$

$$= \frac{\frac{\cos \theta + \sin \theta}{\cos \theta}}{\frac{\sin \theta + \cos \theta}{\sin \theta}}$$

$$= \frac{\cos \theta + \sin \theta}{\cos \theta} \cdot \frac{\sin \theta}{\sin \theta + \cos \theta}$$

$$= \frac{\sin \theta}{\cos \theta} \quad \text{as expected.}$$