

## Section 15: Shift Theorems

We have two major shift (a.k.a. translation) theorems.

**Theorem:** Suppose  $\mathcal{L}\{f(t)\} = F(s)$ . Then for any real number  $a$


$$\mathcal{L}\{e^{at}f(t)\} = F(s - a).$$

**Theorem:** If  $F(s) = \mathcal{L}\{f(t)\}$  and  $a > 0$ , then

$$\mathcal{L}\{f(t - a)\mathcal{U}(t - a)\} = e^{-as}F(s).$$

## Recall the Unit Step Function

$$\text{For } a > 0, \quad \mathcal{U}(t - a) = \begin{cases} 0, & 0 \leq t < a \\ 1, & t \geq a \end{cases}$$



**Figure:** The function  $\mathcal{U}(t - 0) = \mathcal{U}(t) = 1$  for  $t > 0$ . Here, we are taking  $\mathcal{U}(0) = 1$ , but depending on the context, it might be taken as 0 or  $\frac{1}{2}$ .

## Alternative Form for Translation in $t$

It is often the case that we wish to take the transform of a product of the form

$$g(t)\mathcal{U}(t - a)$$

in which the function  $g$  is not translated.

The main theorem statement

$$\mathcal{L}\{f(t - a)\mathcal{U}(t - a)\} = e^{-as}F(s).$$

can be restated as

$$\mathcal{L}\{g(t)\mathcal{U}(t - a)\} = e^{-as}\mathcal{L}\{g(t + a)\}.$$

This is based on the observation that

$$g(t) = g((t + a) - a).$$

## Example

$$\mathcal{L}\{g(t)\mathcal{U}(t-a)\} = e^{-as}\mathcal{L}\{g(t+a)\}$$

Example: Find  $\mathcal{L}\{\cos t \mathcal{U}\left(t - \frac{\pi}{2}\right)\} = e^{-\frac{\pi}{2}s} \mathcal{L}\left\{\cos\left(t + \frac{\pi}{2}\right)\right\}$

$$a = \frac{\pi}{2}$$

$$g(t) = \cos t$$

$$= e^{-\frac{\pi}{2}s} \mathcal{L}\{-\sin t\}$$

$$= -e^{-\frac{\pi}{2}s} \frac{1}{s^2 + 1} = \frac{-e^{-\frac{\pi}{2}s}}{s^2 + 1}$$

$$\cos\left(t + \frac{\pi}{2}\right) = \underbrace{\cos t}_{0''} \cos \frac{\pi}{2} - \underbrace{\sin t}_{1''} \sin \frac{\pi}{2} = -\sin t$$

## Example

$$\mathcal{L}^{-1}\{e^{-as}F(s)\} = f(t-a)\mathcal{U}(t-a)$$

Example: Find  $\mathcal{L}^{-1}\left\{\frac{e^{-2s}}{s(s+1)}\right\}$

We need  $f(t) = \mathcal{L}^{-1}\{F(s)\}$  where  $F(s) = \frac{1}{s(s+1)}$

Partial fractions

$$\frac{1}{s(s+1)} = \frac{A}{s} + \frac{B}{s+1}$$

$$1 = A(s+1) + Bs$$

$$\begin{array}{l} \text{Set } s=0 \\ s=-1 \end{array} \quad \begin{array}{l} 1 = A \\ 1 = -B \end{array} \quad \frac{1}{s(s+1)} = \frac{1}{s} - \frac{1}{s+1}$$

$$\begin{aligned} f(t) &= \mathcal{L}^{-1} \left\{ \frac{1}{s} - \frac{1}{s+1} \right\} \\ &= \mathcal{L}^{-1} \left\{ \frac{1}{s} \right\} - \mathcal{L}^{-1} \left\{ \frac{1}{s+1} \right\} \\ &= 1 - e^{-t} \end{aligned}$$

$$f(t) = 1 - e^{-t}$$

$$\mathcal{L}^{-1}\{e^{-as}F(s)\} = f(t-a)\mathcal{U}(t-a)$$

$$\mathcal{L}^{-1}\left\{\frac{e^{-2s}}{s(s+1)}\right\} = \left(1 - e^{-(t-2)}\right)\mathcal{U}(t-2)$$

$$a = 2$$