

## Section 5.2: Eigenvectors and Eigenvalues

### Definition:

Let  $A$  be an  $n \times n$  matrix. A nonzero vector  $\mathbf{x}$  such that

$$A\mathbf{x} = \lambda\mathbf{x}$$

for some scalar  $\lambda$  is called an **eigenvector** of the matrix  $A$ .

A scalar  $\lambda$  such that there exists a nonzero vector  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$  is called an **eigenvalue** of the matrix  $A$ . Such a nonzero vector  $\mathbf{x}$  is an *eigenvector corresponding to*  $\lambda$ .

# Eigenspace

## Definition:

Let  $A$  be an  $n \times n$  matrix and  $\lambda$  an eigenvalue of  $A$ . The set of all eigenvectors corresponding to  $\lambda$  together with the zero vector—i.e. the set

$$\{\mathbf{x} \in \mathbb{R}^n \mid \text{and } A\mathbf{x} = \lambda\mathbf{x}\}, = \text{Nul}(A - \lambda I)$$

is called the **eigenspace of  $A$  corresponding to  $\lambda$** .

**Remark:** The video mentioned something called an **Eigenbasis**. When possible, an **eigenbasis** will be constructed by taking bases for all eigenspaces for a matrix and combining them.

We'll get back to this in section 5.3 when we talk about *diagonalizability*.

## Matrices with Nice Structure

### Theorem:

If  $A$  is an  $n \times n$  triangular matrix, then the eigenvalues of  $A$  are its diagonal elements.

**Example:** Find the eigenvalues of the matrix  $A = \begin{bmatrix} 4 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 \\ 1 & 2 & -8 & 0 \\ 1 & 2 & 3 & 489 \end{bmatrix}$

They are

$$\lambda_1 = 4, \lambda_2 = 2, \lambda_3 = -8$$

$$\text{and } \lambda_4 = 489.$$

## Example

Suppose  $\lambda = 0$  is an eigenvalue<sup>1</sup> of a matrix  $A$ . Argue that  $A$  is not invertible.

If  $\lambda = 0$  is an eigenvalue, then there's a non zero vector  $\vec{x}$  such that

$$A\vec{x} = 0\vec{x} \Rightarrow A\vec{x} = \vec{0}.$$

Since the homogeneous equation has a nontrivial solution,  $A$  must not be invertible.

---

<sup>1</sup>Eigenvectors must be nonzero vectors, but it is perfectly legitimate to have a zero eigenvalue!

# Theorems

## Theorem:

A square matrix  $A$  is invertible if and only if zero is **not** an eigenvalue.

## Theorem:

If  $\mathbf{v}_1, \dots, \mathbf{v}_r$  are eigenvectors of a matrix  $A$  corresponding to distinct eigenvalues,  $\lambda_1, \dots, \lambda_r$ , then the set  $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$  is linearly independent.

# Linear Independence

Suppose  $\mathbf{v}_1$  and  $\mathbf{v}_2$  are eigenvectors of a matrix  $A$  corresponding to distinct eigenvalues  $\lambda_1$  and  $\lambda_2$  (i.e.  $\lambda_1 \neq \lambda_2$ ).

Show that  $\{\mathbf{v}_1, \mathbf{v}_2\}$  is linearly independent.

We know  $\vec{v}_1 \neq \vec{0}$ ,  $\vec{v}_2 \neq \vec{0}$ ,  $\lambda_1 \neq \lambda_2$ ,  $A\vec{v}_1 = \lambda_1\vec{v}_1$

and  $A\vec{v}_2 = \lambda_2\vec{v}_2$ . Consider the homogeneous vector equation

$$c_1\vec{v}_1 + c_2\vec{v}_2 = \vec{0} \quad (*)$$

We need to show that  $c_1 = 0$  and  $c_2 = 0$  is necessarily true. We'll create two equations.

① Multiply (\*) by  $A$ ,

② Multiply (\*) by  $\lambda_1$ .

$$\textcircled{1} \quad A(c_1 \vec{v}_1 + c_2 \vec{v}_2) = A \vec{0}$$

$$c_1 A \vec{v}_1 + c_2 A \vec{v}_2 = \vec{0}$$

$$c_1 \lambda_1 \vec{v}_1 + c_2 \lambda_2 \vec{v}_2 = \vec{0} \quad \leftarrow \text{save this.}$$

$$\textcircled{2} \quad \lambda_1 (c_1 \vec{v}_1 + c_2 \vec{v}_2) = \lambda_1 \vec{0}$$

$$c_1 \lambda_1 \vec{v}_1 + c_2 \lambda_1 \vec{v}_2 = \vec{0} \quad \leftarrow \text{save this}$$

Subtract one equation from the other

$$c_1 \lambda_1 \vec{v}_1 + c_2 \lambda_2 \vec{v}_2 = \vec{0}$$

$$- \quad c_1 \lambda_1 \vec{v}_1 + c_2 \lambda_1 \vec{v}_2 = \vec{0}$$

---

$$\vec{0} + c_2 \lambda_2 \vec{v}_2 - c_2 \lambda_1 \vec{v}_2 = \vec{0}$$

$$c_2 (\lambda_2 - \lambda_1) \vec{v}_2 = \vec{0}$$

So  $c_2 = 0$ , or  $\lambda_2 - \lambda_1 = 0$ , or  $\vec{v}_2 = \vec{0}$ .

$\vec{v}_2 \neq \vec{0}$  as an eigenvector,  $\lambda_2 - \lambda_1 \neq 0$  since  $\lambda_1 \neq \lambda_2$ .

So  $c_2 = 0$ . Then equation (\*) becomes

$$c_1 \vec{v}_1 + 0 \vec{v}_2 = \vec{0} \Rightarrow c_1 \vec{v}_1 = \vec{0}$$

Since  $\vec{v}_1 \neq \vec{0}$ , it must be that  $c_1 = 0$ .

Because  $c_1 = c_2 = 0$  is the only solution to (\*) )  $\{ \vec{v}_1, \vec{v}_2 \}$  is linearly

independent.



## Another Addendum to the Invertible Matrix Thm.

### Theorem:

The  $n \times n$  matrix  $A$  is invertible if and only if<sup>a</sup>

- (s) The number 0 is not an eigenvalue of  $A$ .
- (t) The determinant of  $A$  is nonzero.

---

<sup>a</sup>This is nothing new, we're just adding to the list.

## Section 5.2: The Characteristic Equation

Find the eigenvalues of  $A = \begin{bmatrix} 2 & 3 \\ 3 & -6 \end{bmatrix}$  by appealing to the fact that the equation  $A\mathbf{x} = \lambda I\mathbf{x}$  can be restated as:

Find a nontrivial solution of the homogeneous equation

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

we need  $\det(A - \lambda I) = 0$

$$A - \lambda I = \begin{bmatrix} 2 & 3 \\ 3 & -6 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} = \begin{bmatrix} 2-\lambda & 3 \\ 3 & -6-\lambda \end{bmatrix}$$

$$\det(A - \lambda I) = (2-\lambda)(-6-\lambda) - 3 \cdot 3$$

$$= -\lambda^2 + 4\lambda - 12 - 9$$

$$= \lambda^2 + 4\lambda - 21$$

set to zero,

$$\lambda^2 + 4\lambda - 21 = 0$$

$$(\lambda + 7)(\lambda - 3) = 0$$

we find two eigenvalues

$$\lambda_1 = -7, \lambda_2 = 3$$

# Characteristic Equation

## Definition:

For  $n \times n$  matrix  $A$ , the expression  $\det(A - \lambda I)$  is an  $n^{\text{th}}$  degree polynomial in  $\lambda$ . It is called the **characteristic polynomial** of  $A$ .

## Definition:

The equation  $\det(A - \lambda I) = 0$  is called the **characteristic equation** of  $A$ .

## Theorem:

The scalar  $\lambda$  is an eigenvalue of the matrix  $A$  if and only if it is a root of the characteristic equation.