

# August 14 Math 2306 sec. 51 Fall 2024

## Section 1: Concepts and Terminology

Suppose  $y = \phi(x)$  is a differentiable function. We know that  $\frac{dy}{dx} = \phi'(x)$  is another (related) function.

For example, if  $y = \cos(2x)$ , then  $y$  is differentiable on  $(-\infty, \infty)$ . In fact,

$$\frac{dy}{dx} = -2 \sin(2x).$$

Even  $dy/dx$  is differentiable with  $d^2y/dx^2 = -4 \cos(2x)$ . Note that this function happens to satisfy the equation

$$\frac{d^2y}{dx^2} + 4y = 0.$$

$$y'' = -4 \cos(2x) \quad , \quad 4y = 4 \cos(2x)$$

Equations like

$$\frac{dr}{dt} = \frac{\beta}{3\rho} \left( \frac{3\alpha}{\beta} - r \right) \quad \text{and} \quad \frac{d^2y}{dx^2} + 4y = 0$$

are examples of differential equations.

Some questions we can ask about this second example are

### Questions

1. If we only started with the equation, how could we determine that  $\cos(2x)$  satisfies it?
2. Also, is  $\cos(2x)$  the only possible function that  $y$  could be?

We'll be able to answer these and many more questions as we proceed.

# Definition

## Definition

A **Differential Equation** is an equation containing the derivative(s) of one or more dependent variables, with respect to one or more independent variables.

We can characterize the *variables* in calculus terms:

## Independent and Dependent Variables

An **Independent Variable**: will appear as one that derivatives are taken **with respect to**. (The  $x$  in  $y = f(x)$ .)

A **Dependent<sup>a</sup> Variable**: will appear as one that derivatives are taken **of**. (The  $y$  in  $y = f(x)$ .)

---

<sup>a</sup>Think of dependent variables as the functions.

## Independent and Dependent Variables

Identify the independent and dependent variables in each differential equation shown.

$$\frac{d^2y}{dx^2} + 4y = 0$$

Independent x      Dependent y

$$t^2y'' - 2ty' + y = \sin(t)$$

Independent t      Dependent y

$$2\frac{d^2x}{du^2} = x^2 + u^2$$

Independent u      Dependent x

Sometimes the independent variable isn't explicitly stated, e.g., consider  $y'' + 4y = 0$ . I'll usually just call it  $x$  or maybe  $t$ .

# Classifications: Type ODE or PDE

## ODEs

An **ordinary differential equation (ODE)** has exactly one independent variable<sup>a</sup>. For example

$$\frac{dy}{dx} - y^2 = 3x, \quad \text{or} \quad \frac{dy}{dt} + 2\frac{dx}{dt} = t, \quad \text{or} \quad y'' + 4y = 0$$

---

<sup>a</sup>These are the subject of this course.

## PDEs

A **partial differential equation (PDE)** has two or more independent variables. For example

$$\frac{\partial y}{\partial t} = \frac{\partial^2 y}{\partial x^2}, \quad \text{or} \quad \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$$

## Classifications: Order

### Definition

**Order:** The order of a differential equation is the same as the highest order derivative appearing anywhere in the equation.

Examples: Identify the order of each differential equation.

$$\frac{dy}{dx} - y^2 = 3x$$

Order = 1<sup>st</sup> ODE

$$y''' + (y')^4 = x^3$$

Order = 3<sup>rd</sup> ODE

$$\frac{\partial y}{\partial t} = \frac{\partial^2 y}{\partial x^2}$$

Order = 2<sup>nd</sup>

PDE

## Notations and Symbols

We'll mostly use standard derivative notations:

$$\text{Leibniz: } \frac{dy}{dx}, \quad \frac{d^2y}{dx^2}, \quad \dots \quad \frac{d^ny}{dx^n}, \quad \text{or}$$

$$\text{Prime \& superscripts: } y', \quad y'', \quad \dots \quad y^{(n)}.$$

Newton's **dot notation** is a special notation that is reserved for derivatives **with respect to time**. For example, if  $s(t)$  is the position of a moving particle at the time  $t$ , then

$$\text{velocity is } \frac{ds}{dt} = \dot{s}, \quad \text{and acceleration is } \frac{d^2s}{dt^2} = \ddot{s}$$

Note that the dot is like a prime, but it's placed on top of the variable.

## Notations and Symbols

On occasion, we'll want to reference a generic ODE. We have a couple of formats for that.

An  $n^{\text{th}}$  order ODE, with independent variable  $x$  and dependent variable  $y$  can always be expressed as an equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

where  $F$  is some real valued function of  $n + 2$  variables.

**Example:** Express the equation  $y'' + 4y = 0$  in the form

$$F(x, y, y', \dots, y^{(n)}) = 0$$

this form with  $y'' + 4y = 0$  is in  
 $F(x, y, y', y'') = y'' + 4y$



## Notations and Symbols

### Normal Form

**Normal Form:** If it is possible to isolate the highest derivative term, then we can write a **normal form** of the equation

$$\frac{d^n y}{dx^n} = f(x, y, y', \dots, y^{(n-1)}).$$

E.g. if  $n = 1$   $\frac{dy}{dx} = f(x, y)$ ,      if  $n = 2$   $\frac{d^2 y}{dx^2} = f(x, y, y')$ .

**Example:** Express the equation  $y'' + 4y = 0$  in the normal form.

Subtract  $4 \cdot y$        $y'' = -4y$

$$f(x, y, y') = -4y$$

## Notations and Symbols

### Differential Form

If  $M(x, y)$  and  $N(x, y)$  are functions of the variables  $x$  and  $y$ , then an expression of the form

$$M(x, y) dx + N(x, y) dy$$

is called a **Differential Form**. A first order equation may be written in terms of a differential form as follows:

$$M(x, y) dx + N(x, y) dy = 0$$

Note that this can be rearranged into a couple<sup>1</sup> of different normal forms

$$\frac{dy}{dx} = -\frac{M(x, y)}{N(x, y)} \quad \text{or} \quad \frac{dx}{dy} = -\frac{N(x, y)}{M(x, y)}$$

---

<sup>1</sup>We have to assume that  $N$  or  $M$  is nonzero as needed.

# Classifications: Linearity

## Linear Differential Equation

**Linearity:** An  $n^{\text{th}}$  order differential equation is said to be **linear** if it can be written in the form

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x).$$

**Example First Order:**

$$a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

**Example Second Order:**

$$a_2(x) \frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

# Properties of a Linear ODE

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x).$$

## Properties of Linear ODEs

- ▶ Each of the coefficients  $a_0, \dots, a_n$  and the right hand side  $g$  may depend on the independent variable but not the dependent one.
- ▶  $y$ , and its derivatives can only appear as themselves (not squared, square rooted, inside some other function).
- ▶ The characteristic structure of the left side is that

$$y, \quad \frac{dy}{dx}, \quad \frac{d^2 y}{dx^2}, \quad \cdots, \quad \frac{d^n y}{dx^n}$$

are multiplied by functions of the independent variable and added together.

## Examples (Linear -vs- Nonlinear)

Verify that the two equations here are **linear**.

$$y'' + 4y = 0$$

$$t^2 \frac{d^2 x}{dt^2} + 2t \frac{dx}{dt} - x = e^t$$

$$a_2(x) \frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

$$g(x) = 0$$

$$a_0(x) = 4$$

$$a_1(x) = 0$$

$$a_2(x) = 1$$

$$g(t) = e^t$$

$$a_0(t) = -1$$

$$a_1(t) = 2t$$

$$a_2(t) = t^2$$

## Examples (Linear -vs- Nonlinear)

Determine why the following two equations are **nonlinear**.

$$\frac{d^3y}{dx^3} + \left(\frac{dy}{dx}\right)^4 = x^3$$

$$y''' + (y')^3 y' = x^3$$

nonlinear  
term

$$u'' + u' = \cos u$$

$u$  is dependent

so  $\cos u$

is a nonlinear

term.

$$* u'' + u' = \cos t$$

would be  
linear.

## Example: Classification

Identify the independent and dependent variables. Determine the order of the equation. State whether it is linear or nonlinear.

(a)  $y'' + 2ty' = \cos t + y - y'''$

rearrange

$$y''' + y'' + 2ty' - y = \cos t$$

- ▶ independent var. t
- ▶ dependent var. y
- ▶ order 3<sup>rd</sup>
- ▶ linear/nonlinear linear

Identify the independent and dependent variables. Determine the order of the equation. State whether it is linear or nonlinear.

(b)  $\ddot{\theta} + \frac{g}{l} \sin \theta = 0$   $g$  and  $l$  are constant

$\theta$  is dependent  
 $\sin \theta$  is a  
nonlinear  
term

- ▶ independent var. time  $t$
- ▶ dependent var.  $\theta$
- ▶ order 2<sup>nd</sup>
- ▶ linear/nonlinear non