

Section 1: Concepts and Terminology

Suppose $y = \phi(x)$ is a differentiable function. We know that $dy/dx = \phi'(x)$ is another (related) function.

For example, if $y = \cos(2x)$, then y is differentiable on $(-\infty, \infty)$. In fact,

$$\frac{dy}{dx} = -2 \sin(2x).$$

Even dy/dx is differentiable with $d^2y/dx^2 = -4 \cos(2x)$.

Suppose $y = \cos(2x)$ \leftarrow this function makes this equation true.

Note that $\frac{d^2y}{dx^2} + 4y = 0$.

$$y = \cos(2x) \quad \text{so} \quad \frac{d^2y}{dx^2} = -4 \cos(2x)$$

$$\text{so} \quad \frac{d^2y}{dx^2} + 4y =$$

$$-4 \cos(2x) + 4 \cos(2x) = 0$$

A differential equation

The equation

$$\frac{d^2y}{dx^2} + 4y = 0.$$

is an example of a **differential equation**.

Questions: If we only started with the equation, how could we determine that $\cos(2x)$ satisfies it? Also, is $\cos(2x)$ the only possible function that y could be?

no
there are
others

yes

Definition

A **Differential Equation** is an equation containing the derivative(s) of one or more dependent variables, with respect to one or more independent variables.

Solving a differential equation refers to determining the dependent variable(s)—as function(s).

Independent Variable: will appear as one that derivatives are taken **with respect to**.

Dependent Variable: will appear as one that derivatives are taken **of**.

*↑
functions*

Independent and Dependent Variables

Often, the derivatives indicate which variable is which:

$$\frac{dy}{dx}$$

implies \rightarrow
 y is a
function of x
 y is dependent
 x is independent

$$\frac{du}{dt}$$

\leftarrow dep.
 \uparrow
indep.

$$\frac{dx}{dr}$$

\uparrow
 x is a
function of
 r

Classifications

Type: An **ordinary differential equation (ODE)** has exactly one independent variable¹. For example

$$\frac{dy}{dx} - y^2 = 3x, \quad \text{or} \quad \frac{dy}{dt} + 2\frac{dx}{dt} = t, \quad \text{or} \quad y'' + 4y = 0$$

A **partial differential equation (PDE)** has two or more independent variables. For example

$$\frac{\partial y}{\partial t} = \frac{\partial^2 y}{\partial x^2}, \quad \text{or} \quad \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$$

⓪ partial symbol

¹These are the subject of this course.

Classifications

Order: The order of a differential equation is the same as the highest order derivative appearing anywhere in the equation.

$$\frac{dy}{dx} - y^2 = 3x$$

← 1st derivative

1st order ODE

$$y''' + (y')^4 = x^3$$

3rd order ODE

$$\frac{\partial y}{\partial t} = \frac{\partial^2 y}{\partial x^2}$$

2nd order PDE

Notations and Symbols

We'll use standard derivative notations:

Leibniz: $\frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^ny}{dx^n},$ or

Prime & superscripts: $y', y'', \dots y^{(n)}.$

Newton's **dot notation** may be used if the independent variable is time. For example if s is a position function, then

velocity is $\frac{ds}{dt} = \dot{s},$ and acceleration is $\frac{d^2s}{dt^2} = \ddot{s}$

Notations and Symbols

An n^{th} order ODE, with independent variable x and dependent variable y can always be expressed as an equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

where F is some real valued function of $n + 2$ variables.

Normal Form: If it is possible to isolate the highest derivative term, then we can write a **normal form** of the equation

$$\frac{d^n y}{dx^n} = f(x, y, y', \dots, y^{(n-1)}).$$

$$\frac{d^2y}{dx^2} + 4y = 0$$

This is already written as

$$F(x, y, y', y'') = 0$$

$$\text{Here } F(x, y, y', y'') = y'' + 4y$$

In normal form this is

$$\frac{d^2y}{dx^2} = -4y, \quad f(x, y, y') = -4y$$


Notations and Symbols

If $n = 1$ or $n = 2$, an equation in normal form would look like

$$\frac{dy}{dx} = f(x, y) \quad \text{or} \quad \frac{d^2y}{dx^2} = f(x, y, y').$$

Differential Form: A first order equation may appear in the form

$$M(x, y) dx + N(x, y) dy = 0$$

differential
form

$$M(x, y) dx + N(x, y) dy = 0$$

Differential forms may be written in normal form in a couple of ways.

$$N(x, y) dy = -M(x, y) dx$$

$$dy = \frac{dy}{dx} dx$$

Divide by $N(x, y) dx$

$$\frac{dy}{dx} = \frac{-M(x, y)}{N(x, y)} \quad \text{as long as } N(x, y) \neq 0$$

$$\text{or} \quad \frac{dx}{dy} = \frac{-N(x, y)}{M(x, y)} \quad \text{for } M(x, y) \neq 0$$

Classifications

Linearity: An n^{th} order differential equation is said to be **linear** if it can be written in the form

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x).$$

- ▶ Each of the coefficients a_0, \dots, a_n and the right hand side g may depend on the independent variable but not the dependent one.
- ▶ y , and its derivatives can only appear as themselves (not squared, square rooted, inside some other function).
- ▶ The characteristic structure of the left side is that

$$y, \quad \frac{dy}{dx}, \quad \frac{d^2 y}{dx^2}, \dots, \frac{d^n y}{dx^n}$$

are multiplied by functions of the independent variable and added together.

Examples (Linear -vs- Nonlinear) Both linear

$$y'' + 4y = 0$$

$$a_2(x)y'' + a_1(x)y' + a_0(x)y = g(x)$$

$$a_2(x) = 1$$

$$a_1(x) = 0$$

$$a_0(x) = 4$$

$$g(x) = 0$$

$$t^2 \frac{d^2 x}{dt^2} + 2t \frac{dx}{dt} - x = e^t$$

$$a_2(t)x'' + a_1(t)x' + a_0(t)x = g(t)$$

$$a_2(t) = t^2$$

$$a_1(t) = 2t$$

$$a_0(t) = -1$$

$$g(t) = e^t$$

Examples (Linear -vs- Nonlinear)

$$\frac{d^3 y}{dx^3} + \left(\frac{dy}{dx}\right)^4 = x^3$$

nonlinear
term

nonlinear

$$u'' + u' = \cos u$$

nonlinear
term

u is
dependent