

# August 23 Math 2306 sec. 52 Fall 2021

We'll close out section 1 (Concepts & Terminology) with a few terms.

- ▶ A **parameter** is an unspecified constant such as  $c_1$  and  $c_2$  in the last example.
- ▶ A **family of solutions** is a collection of solution functions that only differ by a parameter.
- ▶ An  **$n$ -parameter family of solutions** is one containing  $n$  parameters (e.g.  $c_1 x + \frac{c_2}{x}$  is a 2 parameter family).

## Some final terms

- ▶ A **particular solution** is one with no arbitrary constants in it.
- ▶ The **trivial solution** is the simple constant function  $y = 0$ .
- ▶ An **integral curve** is the graph of one solution (perhaps from a family).

## Section 2: Initial Value Problems

An initial value problem consists of an ODE with additional conditions.

Solve the equation <sup>1</sup>

$$\frac{d^n y}{dx^n} = f(x, y, y', \dots, y^{(n-1)}) \quad (1)$$

subject to the *initial conditions*

$$y(x_0) = y_0, \quad y'(x_0) = y_1, \quad \dots, y^{(n-1)}(x_0) = y_{n-1}. \quad (2)$$

The problem (1)–(2) is called an *initial value problem* (IVP).

---

<sup>1</sup>on some interval  $I$  containing  $x_0$ .

# IVPs

First order case:

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0$$

1st  
order  
ODE  $\nearrow$

$\uparrow$   
one initial  
condition

The initial condition tells us that the point  $(x_0, y_0)$  is on the solution curve,  $y$ , if  $y$  solves the IVP.

# IVPs

Second order case:

$$\frac{d^2 y}{dx^2} = f(x, y, y'), \quad y(x_0) = y_0, \quad y'(x_0) = y_1$$

2nd  
order  
ODE  $\rightarrow$

2 initial  
conditions

For example, if  $y$  is the position of a particle moving along a line at time  $x$ , the ODE tells us about the acceleration,  $y_0$  is the initial position, and  $y_1$  is the initial velocity.

## Example

Given that  $y = c_1 x + \frac{c_2}{x}$  is a 2-parameter family of solutions of  $x^2 y'' + xy' - y = 0$ , solve the IVP

$$x^2 y'' + xy' - y = 0, \quad y(1) = 1, \quad y'(1) = 3$$

We already have all the solutions to the ODE. They are in the family

$y = c_1 x + \frac{c_2}{x}$ . Our task is to find numbers  $c_1$  and  $c_2$  such that  $y(1) = 1$  and  $y'(1) = 3$ .

$$y = c_1 x + \frac{c_2}{x}, \quad y' = c_1 - \frac{c_2}{x^2}$$

$$y(1) = 1 \Rightarrow 1 = c_1(1) + \frac{c_2}{1} \Rightarrow c_1 + c_2 = 1$$

$$y'(1) = 3 \Rightarrow 3 = c_1 - \frac{c_2}{1^2} \Rightarrow c_1 - c_2 = 3$$

we'll solve the system

$$c_1 + c_2 = 1 \quad \text{add} \quad 2c_1 = 4$$


$$c_1 - c_2 = 3 \quad c_1 = 2$$

$$c_2 = -1$$

The solution to the IVP is

$$y = 2x - \frac{1}{x}$$

# Graphical Interpretation



**Figure:** Each curve solves  $y' + 2xy = 0$ ,  $y(0) = y_0$ . Each colored curve corresponds to a different value of  $y_0$



# A Numerical Solution


Consider a first order initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0.$$

**Euler's Method** is a scheme for finding an approximate solution to this IVP. The basic idea is that we

- ▶ Start with the known point  $(x_0, y_0)$  on the solution curve,
- ▶ use the slope (given by  $\frac{dy}{dx}$ ) to get a tangent line there, and
- ▶ approximate a nearby point on the curve by the tangent line.
- ▶ march forward a little bit, and repeat.


Example  $\frac{dy}{dx} = xy, \quad y(0) = 1$



**Figure:** We know that the point  $(x_0, y_0) = (0, 1)$  is on the curve. And the slope of the curve at  $(0, 1)$  is  $m_0 = f(0, 1) = 0 \cdot 1 = 0$ .


Note: The gray curve is the true solution to this IVP. It's shown for reference.

Example  $\frac{dy}{dx} = xy, \quad y(0) = 1$




**Figure:** So we draw a little tangent line (we know the point and slope). Then we increase  $x$ , say  $x_1 = x_0 + h$ , and approximate the solution value  $y(x_1)$  with the value on the tangent line  $y_1$ . So  $y_1 \approx y(x_1)$ .

Example  $\frac{dy}{dx} = xy, \quad y(0) = 1$




**Figure:** We take the approximation to the true function  $y$  at the point  $x_1 = x_0 + h$  to be the point on the tangent line.

Example  $\frac{dy}{dx} = xy, \quad y(0) = 1$




**Figure:** When  $h$  is very small, the true solution and the tangent line point will be close. Here, we've zoomed in to see that there is some error between the exact  $y$  value and the approximation from the tangent line.

Example  $\frac{dy}{dx} = xy, \quad y(0) = 1$




**Figure:** Now we start with the point  $(x_1, y_1)$  and repeat the process. We get the slope  $m_1 = f(x_1, y_1)$  and draw a tangent line through  $(x_1, y_1)$  with slope  $m_1$ .

Example  $\frac{dy}{dx} = xy, \quad y(0) = 1$



**Figure:** We go out  $h$  more units to  $x_2 = x_1 + h$ . Pick the point on the tangent line  $(x_2, y_2)$ , and use this to approximate  $y(x_2)$ . So  $y_2 \approx y(x_2)$


Example  $\frac{dy}{dx} = xy, \quad y(0) = 1$



**Figure:** If we zoom in, we can see that there is some error. But as long as  $h$  is small, the point on the tangent line approximates the point on the actual solution curve.



Example  $\frac{dy}{dx} = xy, \quad y(0) = 1$



**Figure:** We can repeat this process at the new point to obtain the next point. We build an approximate solution by advancing the independent variable and connect the points  $(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$ .

# Euler's Method: An Algorithm & Error

We start with the IVP

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0.$$

We build a sequence of points that approximates the true solution  $y$

$$(x_0, y_0), (x_1, y_1), (x_2, y_2), \dots, (x_N, y_N).$$

We'll take the  $x$  values to be equally spaced with a common difference of  $h$ . That is

$$x_1 = x_0 + h$$

$$x_2 = x_1 + h = x_0 + 2h$$

$$x_3 = x_2 + h = x_0 + 3h$$

$$\vdots$$

$$x_n = x_0 + nh$$

# Euler's Method: An Algorithm

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0.$$

## Notation:

- ▶  $y_n$  will denote our approximation, and
- ▶  $y(x_n)$  will denote the exact solution (that we don't know)

To build a formula for the approximation  $y_1$ , let's approximate the derivative at  $(x_0, y_0)$ .

$$f(x_0, y_0) = \left. \frac{dy}{dx} \right|_{(x_0, y_0)} \approx \frac{y_1 - y_0}{x_1 - x_0}$$

(Notice that's the standard formula for slope. )

# Euler's Method: An Algorithm

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0.$$

Let's get a formula for  $y_1$ .  $x_1 - x_0 = h$

$$\frac{y_1 - y_0}{x_1 - x_0} = \frac{y_1 - y_0}{h} = f(x_0, y_0)$$

$$y_1 - y_0 = h f(x_0, y_0)$$

$$\text{so } y_1 = y_0 + h f(x_0, y_0)$$

# Euler's Method: An Algorithm

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0.$$

We can continue this process. So we use

$$\frac{y_2 - y_1}{h} = f(x_1, y_1) \implies y_2 = y_1 + hf(x_1, y_1)$$

and so forth. We have

**Euler's Method Formula:** The  $n^{\text{th}}$  approximation  $y_n$  to the exact solution  $y(x_n)$  is given by

$$y_n = y_{n-1} + hf(x_{n-1}, y_{n-1})$$

with  $(x_0, y_0)$  given in the original IVP and  $h$  the choice of step size.

Euler's Method Example:  $\frac{dy}{dx} = xy$ ,  $y(0) = 1$

Take  $h = 0.25$  to find an approximation to  $y(1)$ .

$$x_0 = 0 \text{ and } h = 0.25, \text{ so } x_4 = 1$$

$$x_0 = 0, \quad y_0 = 1 \quad h = 0.25$$

$$y_1 = y_0 + h f(x_0, y_0)$$

$$= 1 + 0.25(0.1) = 1$$

$$f(x, y) = xy$$

$$x_1 = 0.25, \quad y_1 = 1$$

$$y_2 = y_1 + h f(x_1, y_1) = 1 + 0.25(0.25 \cdot 1)$$

$$= 1.0625$$

$$x_2 = 0.5, \quad y_2 = 1.0625$$

$$\begin{aligned} y_3 &= y_2 + h f(x_2, y_2) \\ &= 1.0625 + 0.25(0.5 \cdot 1.0625) = 1.19531 \end{aligned}$$

$$x_3 = 0.75, \quad y_3 = 1.19531$$

$$\begin{aligned} y_4 &= y_3 + h f(x_3, y_3) \\ &= 1.19531 + 0.25(0.75 \cdot 1.19531) \\ &= 1.41943 \end{aligned}$$

$$y(1) \approx y_4 = 1.41943$$

The true,  $y(1) = \sqrt{e} \approx 1.64872$

We'll talk about the error on Wednesday.