

Section 3: Separation of Variables

We said that a first order equation of the form

$$\frac{dy}{dx} = g(x)h(y)$$

is called a **separable** differential equation.

A solution (usually implicit) is found by **separating the variables**.

$$\int \frac{dy}{h(y)} = \int g(x) dx$$

(assuming $h(y)$ is nonzero on the interval of interest)

An IVP¹

$$\frac{dQ}{dt} = -2(Q-70), \quad Q(0) = 180$$

$$\frac{1}{Q-70} \frac{dQ}{dt} = -2$$

$$\frac{1}{Q-70} \frac{dQ}{dt} dt = -2 dt$$

$$\int \frac{1}{Q-70} dQ = \int -2 dt$$

$$\ln |Q-70| = -2t + C$$

Let's isolate Q :

$$\frac{dQ}{dt} = g(t) h(Q)$$

$$g(t) = -2$$

$$h(Q) = Q-70$$

a 1-parameter
family
implicit

¹Recall IVP stands for *initial value problem*.

$$e^{\ln|Q-70|} = e^{-zt+C} = e^{-zt} \cdot e^C$$

$$|Q-70| = e^C e^{-zt} \quad \text{let } k = \pm e^C$$

$$Q-70 = k e^{-zt}$$

$$Q(t) = 70 + k e^{-zt}$$

1-parameter
family explicit

$$\text{Apply } Q(0) = 180 \Rightarrow 180 = 70 + k e^0 \Rightarrow k = 110$$

The solution to the IVP is

$$Q = 70 + 110 e^{-zt}$$

Find an explicit solution to the IVP.

$$\frac{dy}{dx} = 4x\sqrt{y}, \quad y(0) = 0$$

Here $g(x) = 4x$

$$h(y) = \sqrt{y}$$

$$\frac{1}{\sqrt{y}} \frac{dy}{dx} = 4x$$

$$\frac{1}{\sqrt{y}} \frac{dy}{dx} dx = 4x dx$$

$$\int \frac{1}{\sqrt{y}} dy = \int 4x dx \Rightarrow \int y^{-1/2} dy = \int 4x dx$$

$$\frac{y^{1/2}}{1/2} = 2x^2 + C$$

$$y^{1/2} = x^2 + \frac{1}{2}C$$

$$\Rightarrow \sqrt{y} = x^2 + k$$

$$k = \frac{1}{2}C$$

Apply $y(0)=0$ $\sqrt{0} = 0^2 + k \Rightarrow k = 0$

An implicit solution is

$$\sqrt{y} = x^2$$

Solving for y , $(\sqrt{y})^2 = (x^2)^2 \Rightarrow y = x^4$

An explicit solution is

$$y = x^4$$

Missed Solution

We made an assumption about being able to divide by $h(y)$ when solving $\frac{dy}{dx} = g(x)h(y)$. This may cause us to not find valid solutions.

The IVP $\frac{dy}{dx} = 4x\sqrt{y}$, $y(0) = 0$ has two distinct solutions

$$y = x^4, \quad \text{and} \quad y(x) = 0.$$

The second solution **CANNOT** be found by separation of variables.

Why?

$\frac{1}{\sqrt{y}}$ is not valid if $y = 0$

Missed Solutions $\frac{dy}{dx} = g(x)h(y)$.

Theorem: If the number c is a zero of the function h , i.e. $h(c) = 0$, then the constant function $y(x) = c$ is a solution to the differential equation $\frac{dy}{dx} = g(x)h(y)$.

Remark: Such a constant solution may or may not be recovered by separation of variables. We can always look for such solutions in addition to separation of variables.

Solutions Defined by Integrals

The Fundamental Theorem of Calculus tells us that: If g and $\frac{dy}{dx}$ are continuous on an interval $[x_0, b)$ and x is in this interval, then

$$\frac{d}{dx} \int_{x_0}^x g(t) dt = g(x) \quad \text{and} \quad \int_{x_0}^x \frac{dy}{dt} dt = y(x) - y(x_0).$$

Theorem: If g is continuous on some interval containing x_0 , then the function

$$y = y_0 + \int_{x_0}^x g(t) dt$$

is a solution of the initial value problem

$$\frac{dy}{dx} = g(x), \quad y(x_0) = y_0$$

Example

Express the solution of the IVP in terms of an integral.

$$\frac{dy}{dx} = \sin(x^2), \quad y(\sqrt{\pi}) = 1$$

$$g(x) = \sin(x^2)$$

$$x_0 = \sqrt{\pi} \text{ and } y_0 = 1$$

$$y = y_0 + \int_{x_0}^x g(t) dt$$

$$\text{The solution is } y = 1 + \int_{\sqrt{\pi}}^x \sin(t^2) dt$$

Let's verify that this solves the IVP.

Does $y(\sqrt{\pi})=1$?

$$y(\sqrt{\pi}) = 1 + \int_{\sqrt{\pi}}^{\sqrt{\pi}} \sin(t^2) dt$$

$$= 1 + 0 = 1$$

this does
satisfy the
IC

Is $\frac{dy}{dx} = \sin(x^2)$?

$$\frac{dy}{dx} = \frac{d}{dx} \left(1 + \int_{\sqrt{\pi}}^x \sin(t^2) dt \right)$$

$$= \frac{d}{dx} (1) + \frac{d}{dx} \int_{\sqrt{\pi}}^x \sin(t^2) dt$$

$$= 0 + \sin(x^2)$$

$$= \sin(x^2)$$

yes it
satisfies the
ODE