

Section 3: Separation of Variables

Separable Differential Equations

Recall that a first order equation of the form

$$\frac{dy}{dx} = g(x)h(y) \quad (1)$$

is called **separable**.

- ▶ If  $h(c) = 0$ , the  $y = c$  is a constant solution to (1).
- ▶ The equation (1) may be solved by separation of variables,

$$\int \frac{dy}{h(y)} = \int g(x) dx$$

## Solutions Expressed as Integrals

**Theorem:** If  $g$  is continuous on some interval containing  $x_0$ , then the function

$$y = y_0 + \int_{x_0}^x g(t) dt$$

is a solution of the initial value problem  $\frac{dy}{dx} = g(x)$ ,  $y(x_0) = y_0$ .

### Generalizing

If  $p$  and  $g$  are sufficiently continuous then

$$\int_{y_0}^y p(z) dz = \int_{x_0}^x g(t) dt \quad \text{solves} \quad \frac{dy}{dx} = \frac{g(x)}{p(y)}, \quad y(x_0) = y_0$$

Example: Express the solution of the IVP in terms of an integral.

$$\frac{dy}{dx} = \sin(x^2), \quad y(\sqrt{\pi}) = 1$$

Identify the pieces

$$g(x) = \sin(x^2), \quad x_0 = \sqrt{\pi}, \quad y_0 = 1$$

$$y = y_0 + \int_{x_0}^x g(t) dt$$

The solution to the IVP is

$$y = 1 + \int_{\sqrt{\pi}}^x \sin(t^2) dt$$



## Section 4: First Order Equations: Linear

Recall that a first order linear equation has the form

$$a_1(x) \frac{dy}{dx} + a_0(x)y = g(x).$$

If  $g(x) = 0$  the equation is called **homogeneous**. Otherwise it is called **nonhomogeneous**.

### Standard Form

Provided  $a_1(x) \neq 0$  on the interval  $I$  of definition of a solution, we can write the **standard form** of the equation

$$\frac{dy}{dx} + P(x)y = f(x).$$

$$P(x) = \frac{a_0(x)}{a_1(x)}$$

$$f(x) = \frac{g(x)}{a_1(x)}$$

We'll be interested in equations (and intervals  $I$ ) for which  $P$  and  $f$  are continuous on  $I$ .

# The Solutions of $\frac{dy}{dx} + P(x)y = f(x)$

The solution to a first order linear ODE always has the same basic structure

$$y(x) = y_c(x) + y_p(x) \quad \text{where}$$

- ▶  $y_c$  is called the **complementary** solution. The complementary solution solves **associated homogeneous** equation,  $\frac{dy}{dx} + P(x)y = 0$ , and
- ▶  $y_p$  is called the **particular** solution. The particular solution depends heavily on  $f$  and is zero if  $f(x) = 0$ .

With higher order equations, we'll have to find  $y_c$  and  $y_p$  separately, but for first order equations we have a process for finding the whole solution.

## Motivating Example

Find the solutions of  $x^2 \frac{dy}{dx} + 2xy = e^x$ .

This is not in standard form, but will work w/ it as is. The left side is the derivative of the product  $x^2 y$ .

$$\frac{d}{dx} (x^2 y) = x^2 \frac{dy}{dx} + 2xy$$

So the ODE is

$$\frac{d}{dx} (x^2 y) = e^x$$

The goal is to find  $y$ . Integrate

$$\int \frac{d}{dx} (x^2 y) dx = \int e^x dx$$

$$x^2 y = e^x + C$$

Isolate  $y$

$$y = \frac{e^x + C}{x^2}$$

this is a  
1-parameter  
family of  
solutions

$$y = \frac{C}{x^2} + \frac{e^x}{x^2}$$

$$y_c = \frac{C}{x^2}, \quad y_p = \frac{e^x}{x^2}$$





# Derivation of Solution via Integrating Factor

Solve the equation in standard form

$$\frac{dy}{dx} + P(x)y = f(x)$$

we will manipulate this equation so that the left side becomes a derivative of a product. Multiply the equation by a special function  $\mu(x)$ , called an integrating factor, so that the left side is  $\frac{d}{dx}(\mu(x)y)$ . Multiply by  $\mu$  assuming it exists.

$$\mu \left( \frac{dy}{dx} + P(x)y \right) = \mu f(x)$$

$$\mu \frac{dy}{dx} + P(x)\mu y = \mu f(x)$$

this should equal  $\frac{d}{dx}(\mu y)$

$$\frac{d}{dx}(\mu y) = \mu \frac{dy}{dx} + \frac{d\mu}{dx} y$$

Compare

$$\underline{\mu \frac{dy}{dx}} + \underline{\frac{d\mu}{dx} y} = \underline{\mu \frac{dy}{dx}} + \underline{P(x)\mu y}$$

Pink terms must match!

$$\frac{d\mu}{dx} y = P(x) \mu y$$

For  $y \neq 0$

$$\frac{d\mu}{dx} = P(x) \mu$$

1st  
order  
sep and ble  
ODE for

$\mu$

$$\int \frac{1}{\mu} d\mu = \int P(x) dx$$

$$\ln|\mu| = \int P(x) dx$$

suppose

$$\mu > 0$$

$$\mu = e^{\int p(x) dx}$$

This is  
the integrating  
factor

# Integrating Factor

## Integrating Factor

For the first order, linear ODE in standard form

$$\frac{dy}{dx} + P(x)y = f(x),$$

the integrating factor

$$\mu(x) = \exp\left(\int P(x) dx\right).$$

Let's list the steps involved in solving a first order linear ODE.