

## 1.3.2 Span

We'll recall that for vectors  $\vec{x} = \langle x_1, x_2, \dots, x_n \rangle$  and  $\vec{y} = \langle y_1, y_2, \dots, y_n \rangle$  in  $R^n$  and scalar  $c$  in  $R$ , vector addition and scalar multiplication are defined by

$$\vec{x} + \vec{y} = \langle x_1 + y_1, x_2 + y_2, \dots, x_n + y_n \rangle, \quad \text{and} \quad c\vec{x} = \langle cx_1, cx_2, \dots, cx_n \rangle.$$

### Linear Combination in $R^n$

Let  $S = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k\}$  be a set of one or more ( $k \geq 1$ ) vectors in  $R^n$ . A **linear combination** of these vectors is any vector of the form

$$c_1 \vec{v}_1 + c_2 \vec{v}_2 + \dots + c_k \vec{v}_k,$$

where  $c_1, \dots, c_k$  are scalars. The coefficients,  $c_1, \dots, c_k$ , are often called the **weights**. They can also be called **coefficients**.

## 1.3.2 Span: Let's Recall Some Examples

Last week, we determined that the **set of all linear combinations of the vector**  $\vec{e}_1 = \langle 1, 0 \rangle$  in  $R^2$  is the horizontal (i.e.,  $x_1$ ) axis.

In a previous example, we showed that the set of all vectors in  $R^3$  that are orthogonal to  $\vec{x} = \langle 1, -1, 2 \rangle$  in  $R^3$  have the form

$$\vec{y} = y_2 \langle 1, 1, 0 \rangle + y_3 \langle -2, 0, 1 \rangle,$$

where the scalars  $y_2$  and  $y_3$  can be any real numbers. We can call this **the set of all linear combinations of**  $\langle 1, 1, 0 \rangle$  and  $\langle -2, 0, 1 \rangle$ .

We'll call a vector like  $\langle 1, 1, 0 \rangle$  (with numbers but not variables in it) a **fixed vector**.

## 1.3.2 Span

### Definition: Span

Let  $S = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k\}$  be a set of one or more ( $k \geq 1$ ) vectors in  $R^n$ . The set of all possible linear combinations of the vectors in  $S$  is called the **subspace of  $R^n$  spanned by  $S$** . We often refer to this as **span** of  $S$ . It is denoted

$$\text{Span}(S) \quad \text{or by} \quad \text{Span}\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k\}.$$

**Example:** If  $S$  is a set of one vector,  $S = \{\vec{v}\}$ , then

$$\text{Span}(S) = \{c\vec{v} \mid c \in R\}.$$


That is,  $\text{Span}(S) = \text{Span}\{\vec{v}\}$  is the set of all scalar multiples of the vector  $\vec{v}$ . If  $S = \{\vec{u}, \vec{v}\}$ , so  $S$  is a set of two vectors, then

$$\text{Span}(S) = \{c_1\vec{u} + c_2\vec{v} \mid c_1, c_2 \in R\},$$

and so forth.

## Example

If  $S = \{ \langle \frac{1}{4}, 1 \rangle \}$ , then  $\text{Span}(S)$  is the set of all vectors that are orthogonal to  $\vec{x} = \langle 4, -1 \rangle$ .




**Figure:** For nonzero  $\vec{v}$ ,  $\text{Span}\{\vec{v}\}$  is a line in  $\mathbb{R}^2$  through the origin and parallel to  $\vec{v}$ . Left: Standard representation of  $\langle 4, -1 \rangle$ . Right:  $\langle 4, -1 \rangle$ ,  $\langle 1/4, 1 \rangle$ , and  $\text{Span}\{\langle 1/4, 1 \rangle\}$ .

## Example

If  $S = \{\langle 1, 1, 0 \rangle, \langle -2, 0, 1 \rangle\}$ , then

$\text{Span}(S)$  is the set of all vectors in  $R^3$  that are orthogonal to  $\vec{x} = \langle 1, -1, 2 \rangle$ .



**Figure:** The set of all vectors orthogonal to  $\langle 1, -1, 2 \rangle$  is a plane containing the origin. This plane contains the points  $(1, 1, 0)$  and  $(-2, 0, 1)$ .

## What does it mean for a vector to be in a subspace spanned by a set?

If  $S = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k\}$ , then to say that the vector  $\vec{x}$  is in  $\text{Span}(S)$ —written  $\vec{x} \in \text{Span}(S)$ , means that there exist some scalars  $c_1, \dots, c_k$  such that

$$\vec{x} = c_1 \vec{v}_1 + c_2 \vec{v}_2 + \dots + c_k \vec{v}_k.$$

The symbol “ $\in$ ” means “*is an element of.*” It can be read as “*is in,*” e.g.,

$$x \in \dots \quad \text{“}x \text{ is in...”}$$

Example: Let  $S = \{\langle 1, 0, 0 \rangle, \langle 0, 1, 0 \rangle\}$ .

1. Which of the following vectors are in  $S$ ?

$\langle 1, 0, 0 \rangle$ ,  $\langle 2, 0, 0 \rangle$ ,  $\langle 0, 1, 0 \rangle$ ,  $\langle 1, 1, 0 \rangle$

2. Which of the following vectors<sup>1</sup> are in  $\text{Span}(S)$ ?

$\langle 1, 0, 0 \rangle$ ,  $\langle 2, 0, 0 \rangle$ ,  $\langle 0, 1, 0 \rangle$ ,  $\langle 1, 1, 0 \rangle$

3. Are there vectors in  $\mathbb{R}^3$  that are not in  $\text{Span}(S)$ ?

$$\vec{x} \in \text{Span}(S) \Rightarrow \vec{x} = c_1 \langle 1, 0, 0 \rangle + c_2 \langle 0, 1, 0 \rangle = \langle c_1, c_2, 0 \rangle$$

$\langle 0, 0, 1 \rangle$  is not in  $\text{Span}(S)$ , so yes

there are vectors in  $\mathbb{R}^3$  not in  $\text{Span}(S)$ .

---

<sup>1</sup>Note that  $\text{Span}(S) = \{c_1 \langle 1, 0, 0 \rangle + c_2 \langle 0, 1, 0 \rangle \mid c_1, c_2 \in \mathbb{R}\}$ .

## Example

Let  $S = \{\vec{u}, \vec{v}\}$  where  $\vec{u} = \langle 1, 1 \rangle$  and  $\vec{v} = \langle 1, -1 \rangle$ .

1. Show that the vector  $\vec{x} = \langle 2, 3 \rangle$  is in  $\text{Span}(S)$ .
2. Show that  $\mathbb{R}^2 = \text{Span}(S)$ .

1. we need to show that there are scalars  $c_1, c_2$  such that  $\vec{x} = c_1\vec{u} + c_2\vec{v}$ .

$$\langle 2, 3 \rangle = c_1 \langle 1, 1 \rangle + c_2 \langle 1, -1 \rangle$$

$$\langle 2, 3 \rangle = \langle c_1 + c_2, c_1 - c_2 \rangle$$

This is true if

$$c_1 + c_2 = 2 \text{ and}$$

$$c_1 - c_2 = 3$$

add  
subtract

$$\begin{array}{rcl} c_1 + c_2 = 2 & \text{and} & \\ c_1 - c_2 = 3 & & \\ \hline 2c_1 = 5 & \Rightarrow & c_1 = \frac{5}{2} \\ 2c_2 = -1 & \Rightarrow & c_2 = -\frac{1}{2} \end{array}$$



$$\vec{x} = \frac{5}{2} \langle 1, 1 \rangle + \frac{-1}{2} \langle 1, -1 \rangle \quad \text{so } \vec{x} \in \text{Span}(S).$$

2. Show that  $\mathbb{R}^2 = \text{Span}(S)$ .

we need to show that every vector  $\vec{x}$  in  $\mathbb{R}^2$  is in  $\text{Span}(S)$ : let  $\vec{x} = \langle x_1, x_2 \rangle$  for any  $x_1$  and  $x_2$ . We need to find  $c_1, c_2$  such that  $\vec{x} = c_1 \vec{u} + c_2 \vec{v}$ .

$$\begin{aligned} \langle x_1, x_2 \rangle &= c_1 \langle 1, 1 \rangle + c_2 \langle 1, -1 \rangle \\ &= \langle c_1 + c_2, c_1 - c_2 \rangle \end{aligned}$$

This requires

$$C_1 + C_2 = X_1 \quad \text{ad}$$

$$C_1 - C_2 = X_2$$

$$\text{add} \quad \frac{\quad}{2C_1 = X_1 + X_2} \Rightarrow C_1 = \frac{X_1 + X_2}{2}$$

$$\text{subtract} \quad 2C_2 = X_1 - X_2 \Rightarrow C_2 = \frac{X_1 - X_2}{2}$$

$$\text{So } \vec{x} = \left( \frac{X_1 + X_2}{2} \right) \langle 1, 1 \rangle + \left( \frac{X_1 - X_2}{2} \right) \langle 1, -1 \rangle$$

showing that  $\vec{x} \in \text{Span}(S)$ .

$$\text{Check: } \left( \frac{X_1 + X_2}{2} \right) \langle 1, 1 \rangle + \left( \frac{X_1 - X_2}{2} \right) \langle 1, -1 \rangle$$

$$= \left\langle \frac{x_1 + x_2}{2}, \frac{x_1 + x_2}{2} \right\rangle + \left\langle \frac{x_1 - x_2}{2}, -\left(\frac{x_1 - x_2}{2}\right) \right\rangle$$

$$= \left\langle \frac{x_1 + x_2}{2} + \frac{x_1 - x_2}{2}, \frac{x_1 + x_2}{2} - \frac{x_1 - x_2}{2} \right\rangle$$

$$= \left\langle \frac{x_1 + x_2 + x_1 - x_2}{2}, \frac{x_1 + x_2 - x_1 + x_2}{2} \right\rangle$$

$$= \left\langle \frac{2x_1}{2}, \frac{2x_2}{2} \right\rangle = \langle x_1, x_2 \rangle$$

Since we can write  $\vec{x} = c_1 \vec{u} + c_2 \vec{v}$  for any  $\vec{x} \in \mathbb{R}^2$ ,  
all of  $\mathbb{R}^2$  is in  $\text{Span}(S)$ .