

January 19 Math 3260 sec. 51 Spring 2022

Section 1.2: Row Reduction and Echelon Forms

Recall the examples of augmented matrices for which it was easy to determine whether the system was consistent or inconsistent and what solutions it might have.

$$\begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -2 \end{bmatrix}, \quad \begin{bmatrix} 1 & 2 & 0 & 3 \\ 0 & 1 & -1 & 4 \\ 0 & 0 & 0 & 3 \end{bmatrix}, \quad \begin{bmatrix} 1 & 0 & -2 & -3 \\ 0 & 1 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Some key features we can look for are

- ▶ Columns having only one nonzero entry that is a 1, and
- ▶ A row with all zeroes or all zeros except for the last entry which is nonzero.

Row Reduction and Echelon Forms

Definition:

A matrix is in **echelon form** (a.k.a. **row echelon form**) if the following properties hold

- i Any row of all zeros are at the bottom.
- ii The first nonzero number (called the *leading entry*) in a row is to the right of the first nonzero number in all rows above it.
- iii All entries below a leading entry are zeros.

ref

Is

$$\begin{bmatrix} 2 & 1 & 3 \\ 0 & -1 & 1 \\ 0 & 0 & 7 \end{bmatrix}$$

Is Not

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & -3 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

Reduced Echelon Form

Definition: A matrix is in **reduced echelon form** (a.k.a. **reduced row echelon form**) if it is in echelon form and the following additional properties hold

- iv The leading entry of each row is 1 (called a *leading 1*), and
- v each leading 1 is the only nonzero entry in its column.

Is

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Is Not

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Notation for Elementary Row Operations

The **Elementary Row Operations** are

- i Interchange any two rows (**row swap**).

$$\text{Swap rows } i \text{ and } j: \quad R_i \leftrightarrow R_j$$

- ii Multiply a row by any nonzero constant (**scaling**).

$$\text{Scale row } i \text{ by } k: \quad kR_i \rightarrow R_i$$

- iii Replace a row with the sum of itself and a multiple of another row (**replacement**).

Replace row j with the sum of itself and k times row i :

$$kR_i + R_j \rightarrow R_j$$

Example (finding ref's and rref's)

Find an echelon form for the following matrix using elementary row operations.

$$\begin{bmatrix} 2 & 1 & 3 \\ 4 & 3 & 6 \\ 0 & 3 & 2 \end{bmatrix}$$

- Want nonzero entry in row 1 column 1.

We already have that.

- Want zeros below that 1st leading entry. (using that leading entry,

$$-2R_1 + R_2 \rightarrow R_2$$

$$\begin{bmatrix} 2 & 1 & 3 \\ 0 & 1 & 0 \\ 0 & 3 & 2 \end{bmatrix}$$

Scratch

$$\begin{array}{ccc} -4 & -2 & -6 \\ 4 & 3 & 6 \end{array}$$

- Now we want a nonzero entry in the 2nd row 2nd column.
- Get zeros below it.

$$-3R_2 + R_3 \rightarrow R_3$$

$$\begin{bmatrix} 2 & 1 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

This is a row echelon form. (ref.)

Example (rref)

Find the reduced echelon form for the following matrix.

$$\begin{bmatrix} 2 & 1 & 3 \\ 4 & 3 & 6 \\ 0 & 3 & 2 \end{bmatrix}$$

we can start w/ the rref
we just found.

$$\begin{bmatrix} 2 & 1 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

• need all leading entries
to be 1 and have all
zeros above them.

$$\frac{1}{2} R_3 \rightarrow R_3$$

$$\begin{bmatrix} 2 & 1 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Clear third column $-3R_3 + R_1 \rightarrow R_1$

$$\begin{bmatrix} 2 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Clear column 2. $-R_2 + R_1 \rightarrow R_1$

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Finally, get leading 1 in to p row

$$\frac{1}{2} R_1 \rightarrow R_1 \quad \cdot \quad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

This is an rref.