

January 21 Math 3260 sec. 52 Spring 2022

We defined the **row echelon form** and **reduced row echelon forms** of matrices.

Definition: A matrix is in **row echelon form** (ref) if the following properties hold

- i Any row of all zeros are at the bottom.
- ii The first nonzero number (called the *leading entry*) in a row is to the right of the first nonzero number in all rows above it.
- iii All entries below a leading entry are zeros.

And a matrix is in **reduced row echelon form** (rref) if it is in echelon form and the following additional properties hold

- iv The leading entry of each row is 1 (called a *leading 1*), and
- v each leading 1 is the only nonzero entry in its column.

Example (finding ref's and rref's)

$$\begin{bmatrix} 2 & 1 & 3 \\ 4 & 3 & 6 \\ 0 & 3 & 2 \end{bmatrix}$$

Using elementary row operations, an ref for this matrix is

$$\begin{bmatrix} 2 & 1 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \quad \frac{1}{2}R_3 \rightarrow R_3 \quad \begin{bmatrix} 2 & 1 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

The rref for this matrix is

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Theorem

Theorem: Each matrix is row equivalent to exactly one reduced row echelon matrix.

This says that the rref of a matrix is unique, and it allows the following unambiguous definition:

Definition: A **pivot position** in a matrix A is a location that corresponds to a leading 1 in the reduced echelon form of A . A **pivot column** of a matrix A is a column of A that contains a pivot position.

Identify the pivot position and columns given...

A

rref of A

$$\begin{bmatrix} 0 & -3 & -6 & 4 & 9 \\ -1 & -2 & -1 & 3 & 1 \\ -2 & -3 & 0 & 3 & -1 \\ 1 & 4 & 5 & -9 & -7 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & -3 & 0 & 5 \\ 0 & 1 & 2 & 0 & -3 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

○ these are the pivot positions

The pivot columns are columns 1, 2, and 4.

Complete Row Reduction isn't needed to find Pivots

Find the pivot positions and pivot columns of the matrix

$$\begin{bmatrix} \textcircled{1} & 1 & 4 \\ -2 & \textcircled{1} & -2 \\ 1 & 0 & 2 \end{bmatrix}$$

$\textcircled{0}$ - pivot positions

The pivot columns are 1 and 2.

This matrix has an ref and rref

$$\begin{bmatrix} 1 & 1 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \quad \text{respectively.}$$

Row Reduction Algorithm

To obtain an echelon form, we work from left to right beginning with the top row working downward.

$$\begin{bmatrix} 0 & 3 & -6 & 4 & 6 \\ 3 & -7 & 8 & 8 & -5 \\ 3 & -9 & 12 & 6 & -9 \end{bmatrix}$$

Step 1: Unless the left most column is all zeros, the left most column is a pivot column. The top position is a pivot position.

Step 2: Get a nonzero entry in the top left position by row swapping if needed.

Let's do $R_1 \leftrightarrow R_3$

$$\begin{bmatrix} 3 & -9 & 12 & 6 & -9 \\ 3 & -7 & 8 & 8 & -5 \\ 0 & 3 & -6 & 4 & 6 \end{bmatrix}$$

Row Reduction Algorithm

$$\begin{bmatrix} 3 & -9 & 12 & 6 & -9 \\ 3 & -7 & 8 & 8 & -5 \\ 0 & 3 & -6 & 4 & 6 \end{bmatrix}$$

Step 3: Use row operations to get zeros in all entries below the pivot.

$$- R_1 + R_2 \rightarrow R_2$$

$$\begin{bmatrix} 3 & -9 & 12 & 6 & -9 \\ 0 & 2 & -4 & 2 & 4 \\ 0 & 3 & -6 & 4 & 6 \end{bmatrix}$$

Row Reduction Algorithm

$$\begin{bmatrix} 3 & -9 & 12 & 6 & -9 \\ 0 & 2 & -4 & 2 & 4 \\ 0 & 3 & -6 & 4 & 6 \end{bmatrix}$$

Step 4: Ignore the row with a pivot, all rows above it, the pivot column, and all columns to its left, and repeat steps 1-3 for each pivot position.

Let's do $\frac{1}{2} R_2 \rightarrow R_2$

$$\begin{bmatrix} 3 & -9 & 12 & 6 & -9 \\ 0 & 1 & -2 & 1 & 2 \\ 0 & 3 & -6 & 4 & 6 \end{bmatrix}$$

$-3R_2 + R_3 \rightarrow R_3$

$$\begin{bmatrix} 0 & -3 & 6 & -3 & -6 \\ 0 & 3 & -6 & 4 & 0 \end{bmatrix}$$

Row Reduction Algorithm

$$\begin{bmatrix} 3 & -9 & 12 & 6 & -9 \\ 0 & 1 & -2 & 1 & 2 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

This is
an ref

Row Reduction Algorithm

To obtain a reduced row echelon form: Starting with the right most pivot and working up and to the left, use row operations to get a zero in each position above a pivot. Scale to make each pivot a 1.

$$\begin{bmatrix} 3 & -9 & 12 & 6 & -9 \\ 0 & 1 & -2 & 1 & 2 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

Clear out the 6 and 1
in column 4 above the last
pivot

$$-R_3 + R_2 \rightarrow R_2 \quad \text{and} \quad -6R_3 + R_1 \rightarrow R_1$$

$$\begin{bmatrix} 3 & -9 & 12 & 0 & -9 \\ 0 & 1 & -2 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

Row Reduction Algorithm

Clear out the -9 about the pivot position row 2 column 2.

$$9R_2 + R_1 \rightarrow R_1$$

$$\begin{bmatrix} 0 & 9 & -18 & 0 & 18 \\ 3 & -9 & 12 & 0 & -9 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 0 & -6 & 0 & 9 \\ 0 & 1 & -2 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

Scale row 1 $\frac{1}{3}R_1 \rightarrow R_1$

$$\begin{bmatrix} 1 & 0 & -2 & 0 & 3 \\ 0 & 1 & -2 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

This is the rref.

Echelon Form & Solving a System

Remark: The row operations used to get an rref correspond to an **equivalent** system!

Consider the reduced echelon matrix, and describe the solution set for the associated system of equations (the one who'd have this as its augmented matrix).

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 3 \\ 0 & 0 & 1 & 0 & -2 & 4 \\ 0 & 0 & 0 & 1 & 0 & -9 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$x_1 + x_2 = 3$$

$$x_3 - 2x_5 = 4$$

$$x_4 = -9$$

$$0 = 0$$

The solutions are given by

$$x_1 = 3 - x_2$$

$$x_3 = 4 + 2x_5$$

$$x_4 = -9$$

x_2 & x_5 can be any
real numbers

Basic & Free Variables

Suppose a system has m equations and n variables, x_1, x_2, \dots, x_n . The first n columns of the augmented matrix correspond to the n variables.

- ▶ If the i^{th} column is a pivot column, then x_i is called a **basic variable**.
- ▶ If the i^{th} column is NOT a pivot column, then x_i is called a **free variable**.

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 3 \\ 0 & 0 & 1 & 0 & -2 & 4 \\ 0 & 0 & 0 & 1 & 0 & -9 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The system would have 4 equations in 5 variables. The basic variables are x_1 , x_3 and x_4 . The free variables are x_2 and x_5 .