

Section 3: Separation of Variables

We defined the first order equation as being **separable** if it has the form

$$\frac{dy}{dx} = g(x)h(y).$$

To solve a separable equation, we separate the variables. That is, we convert the ODE to

$$\int \frac{dy}{h(y)} = \int g(x) dx$$

and integrate to obtain a one-parameter family of solutions (usually defined implicitly).

Solutions Defined by Integrals

The Fundamental Theorem of Calculus tells us that: If g and $\frac{dy}{dx}$ are continuous on an interval $[x_0, b)$ and x is in this interval, then

$$\frac{d}{dx} \int_{x_0}^x g(t) dt = g(x) \quad \text{and} \quad \int_{x_0}^x \frac{dy}{dt} dt = y(x) - y(x_0).$$

Theorem: If g is continuous on some interval containing x_0 , then the function

$$y = y_0 + \int_{x_0}^x g(t) dt$$

is a solution of the initial value problem

$$\frac{dy}{dx} = g(x), \quad y(x_0) = y_0$$

Example

Express the solution of the IVP in terms of an integral.

$$\frac{dy}{dx} = \sin(x^2), \quad y(\sqrt{\pi}) = 1$$

$$y = y_0 + \int_{x_0}^x g(t) dt$$

Here, $g(t) = \sin(t^2)$, $x_0 = \sqrt{\pi}$ and $y_0 = 1$

The solution to the IVP is

$$y = 1 + \int_{\sqrt{\pi}}^x \sin(t^2) dt$$

Let's verify:

$$\text{Is } y(\sqrt{\pi}) = 1?$$

yes!

$$y(\sqrt{\pi}) = 1 + \int_{\sqrt{\pi}}^{\sqrt{\pi}} \sin(t^2) dt = 1 + 0 = 1$$

Is $\frac{dy}{dx} = \sin(x^2)$? *yes it does!*

$$\frac{dy}{dx} = \frac{d}{dx} \left(1 + \int_{\sqrt{\pi}}^x \sin(t^2) dt \right)$$

$$= \frac{d}{dx} (1) + \frac{d}{dx} \int_{\sqrt{\pi}}^x \sin(t^2) dt.$$

$$= 0 + \sin(x^2)$$

$$\Rightarrow \frac{dy}{dx} = \sin(x^2)$$

Section 4: First Order Equations: Linear

A first order linear equation has the form

$$a_1(x) \frac{dy}{dx} + a_0(x)y = g(x).$$

If $g(x) = 0$ the equation is called **homogeneous**. Otherwise it is called **nonhomogeneous**.

Provided $a_1(x) \neq 0$ on the interval I of definition of a solution, we can write the **standard form** of the equation

$$\frac{dy}{dx} + P(x)y = f(x).$$

$$P(x) = \frac{a_0(x)}{a_1(x)}$$

$$f(x) = \frac{g(x)}{a_1(x)}$$

We'll be interested in equations (and intervals I) for which P and f are continuous on I .

Solutions (the General Solution)

$$\frac{dy}{dx} + P(x)y = f(x).$$

It turns out the solution will always have a basic form of $y = y_c + y_p$ where

- ▶ y_c is called the **complementary** solution and would solve the equation

$$\frac{dy}{dx} + P(x)y = 0$$

(called the associated homogeneous equation), and

- ▶ y_p is called the **particular** solution, and is heavily influenced by the function $f(x)$.

The cool thing is that our solution method will get both parts in one process—we won't get this benefit with higher order equations!

Motivating Example

$$x^2 \frac{dy}{dx} + 2xy = e^x$$

This is not in standard form.

The left side is the derivative of $x^2 y$. Note that

$$\frac{d}{dx}(x^2 y) = x^2 \frac{dy}{dx} + 2xy$$

So the ODE can be written as

$$\frac{d}{dx}(x^2 y) = e^x$$

Integrate with respect to x

$$\int \frac{d}{dx}(x^2 y) dx = \int e^x dx$$

$$x^2 y = e^x + C$$

So the solutions are

$$y = \frac{e^x + C}{x^2}$$

1 parameter
family
of
solutions

$$y = \frac{e^x}{x^2} + \frac{C}{x^2}$$

$$y_c = \frac{C}{x^2}, \quad y_p = \frac{e^x}{x^2}$$