

## Section 3.1: Introduction to Determinants

For an  $n \times n$  matrix  $A$ , we defined a number  $\det(A)$  called the **determinant** of the matrix. This number is a function of the entries in the matrix and it was defined so that

- ▶ if  $\det(A) \neq 0$ , then  $A$  is nonsingular (a.k.a invertible), and
- ▶ if  $\det(A) = 0$ , then  $A$  is singular.

## $2 \times 2$ and $3 \times 3$ Cases

$$2 \times 2 \quad \det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = a_{11}a_{22} - a_{21}a_{12}.$$

$$3 \times 3 \quad \det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} =$$
$$= a_{11} \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$

## Minors & Cofactors

**Definition:** Let  $n \geq 2$  and let  $A = [a_{ij}]$  be an  $n \times n$  matrix. The  $i, j^{\text{th}}$  **minor** of  $A$  is

$$M_{ij} = \det(A_{ij}),$$

where  $A_{ij}$  is the  $(n - 1) \times (n - 1)$  matrix obtained from  $A$  by removing the  $i^{\text{th}}$  row and the  $j^{\text{th}}$  column.

The  $i, j^{\text{th}}$  **cofactor** of  $A$  is  $C_{ij} = (-1)^{i+j} M_{ij}$ .

Using the notation of cofactor and minors, the determinant of a  $3 \times 3$  matrix  $A$  has the simple formula

$$\begin{aligned} \det(A) &= a_{11}M_{11} - a_{12}M_{12} + a_{13}M_{13} \\ &= a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13} \end{aligned}$$

# The Determinant Defined

**Definition:** For  $n \geq 2$ , the **determinant** of the  $n \times n$  matrix  $A = [a_{ij}]$  is the number

$$\begin{aligned}\det(A) &= a_{11}C_{11} + a_{12}C_{12} + \cdots + a_{1n}C_{1n} & (1) \\ &= \sum_{j=1}^n (-1)^{1+j} a_{1j} M_{1j}\end{aligned}$$

The expression in equation (1) is called a **cofactor expansion**.

## Example

Find all values of  $x$  such that  $A$  is singular where

$$A = \begin{bmatrix} 3-x & 2 & 1 \\ 0 & 2-x & 4 \\ 0 & 3 & 1-x \end{bmatrix}$$

We can use  $\det(A)$   
since  $\det(A) = 0$   
if  $A$  is singular.

$$\det(A) = (3-x) \det \begin{bmatrix} 2-x & 1 \\ 3 & 1-x \end{bmatrix} - 2 \det \begin{bmatrix} 0 & 4 \\ 0 & 1-x \end{bmatrix} + 1 \det \begin{bmatrix} 0 & 2-x \\ 0 & 3 \end{bmatrix}$$

$$= (3-x) [(2-x)(1-x) - 12]$$

$$= (3-x) [2 - 3x + x^2 - 12]$$

$$= (3-x)(x^2-3x-10)$$

We need to know when  $\det(A) = 0$

$$\begin{aligned}\det(A) &= (3-x)(x^2-3x-10) \\ &= (3-x)(x-5)(x+2)\end{aligned}$$

$$\det(A) = 0 \Rightarrow x = 3, x = 5, \text{ or } x = -2$$

So  $A$  is singular if  $x = 3$ ,  $x = 5$ ,  
or  $x = -2$ .

## Theorem:

The determinant of an  $n \times n$  matrix can be computed by cofactor expansion across any row or down any column.

We can fix any row  $i$  of a matrix  $A$  and then

$$\det(A) = \sum_{j=1}^n (-1)^{i+j} a_{ij} M_{ij}$$

Or, we can fix any column  $j$  of a matrix  $A$  and then

$$\det(A) = \sum_{i=1}^n (-1)^{i+j} a_{ij} M_{ij}$$

# Triangular Matrices

**Definition:** The  $n \times n$  matrix  $A = [a_{ij}]$  is said to be **upper triangular** if  $a_{ij} = 0$  for all  $i > j$ .

It is said to be **lower triangular** if  $a_{ij} = 0$  for all  $j > i$ . A matrix that is both upper and lower triangular is a diagonal matrix.

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a_{22} & a_{23} & \cdots & a_{2n} \\ 0 & 0 & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_{nn} \end{bmatrix}$$

**Upper Triangular**

$$\begin{bmatrix} a_{11} & 0 & 0 & \cdots & 0 \\ a_{21} & a_{22} & 0 & \cdots & 0 \\ a_{31} & a_{32} & a_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{nn} \end{bmatrix}$$

**Lower Triangular**



# Determinants of Triangular<sup>1</sup> Matrices

**Theorem:** For  $n \geq 2$ , the determinant of an  $n \times n$  triangular matrix is the product of its diagonal entries.

That is, if  $A = [a_{ij}]$  is a triangular matrix, then

$$\det(A) = a_{11}a_{22}a_{33} \cdots a_{nn}$$

---

<sup>1</sup>We'll use the catch-all name **triangular matrix** to refer to a matrix that is either upper triangular or lower triangular.

## Example: Compute $\det(A)$

$$(i) \quad A = \begin{bmatrix} -1 & 3 & 4 & 0 & 2 \\ 0 & 2 & -3 & 0 & -4 \\ 0 & 0 & 3 & 7 & 5 \\ 0 & 0 & 0 & -4 & 6 \\ 0 & 0 & 0 & 0 & 6 \end{bmatrix}$$

upper triangular

$$\begin{aligned} \det(A) &= (-1)(2)(3)(-4)(6) \\ &= 144 \end{aligned}$$

$$(ii) \quad A = \begin{bmatrix} 7 & 0 & 0 & 0 \\ 3 & 6 & 0 & 0 \\ 0 & -1 & 2 & 0 \\ 4 & 2 & 2 & 2 \end{bmatrix}$$

Lower triangular

$$\begin{aligned} \det(A) &= (7)(6)(2)(2) \\ &= 168 \end{aligned}$$

## Section 3.2: Properties of Determinants

**Theorem:** Let  $A$  be an  $n \times n$  matrix, and suppose the matrix  $B$  is obtained from  $A$  by performing a single elementary row operation<sup>2</sup>. Then

- (i) If  $B$  is obtained by adding a multiple of a row of  $A$  to another row of  $A$  (row replacement), then

$$\det(B) = \det(A).$$

- (ii) If  $B$  is obtained from  $A$  by swapping any pair of rows (row swap), then

$$\det(B) = -\det(A).$$

- (iii) If  $B$  is obtained from  $A$  by scaling any row by the constant  $k$  (scaling), then

$$\det(B) = k\det(A).$$

---

<sup>2</sup>If "row" is replaced by "column" in any of the operations, the conclusions still follow. ↻

## Example: Using Row Operations

Use row operations to obtain a triangular matrix, and then find the determinant.

$$\begin{bmatrix} 2 & 5 & -7 & 3 \\ 0 & 2 & 4 & -2 \\ 0 & 3 & 6 & 2 \\ -2 & -5 & 4 & -2 \end{bmatrix}$$

Call this  $A$ .

Let's get an ref.

$$R_1 + R_4 \rightarrow R_4$$

$$\begin{bmatrix} 2 & 5 & -7 & 3 \\ 0 & 2 & 4 & -2 \\ 0 & 3 & 6 & 2 \\ 0 & 0 & -3 & 1 \end{bmatrix}$$

$$\frac{1}{2} R_2 \rightarrow R_2$$

Effect

① no effect

② factor of  $\frac{1}{2}$

$$\begin{bmatrix} 2 & 5 & -7 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 3 & 6 & 2 \\ 0 & 0 & -3 & 1 \end{bmatrix}$$

$$-3R_2 + R_3 \rightarrow R_3$$

$$\begin{bmatrix} 2 & 5 & -7 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & -3 & 1 \end{bmatrix}$$

$$R_3 \leftrightarrow R_4$$

$$\begin{bmatrix} 2 & 5 & -7 & 3 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & -3 & 1 \\ 0 & 0 & 0 & 5 \end{bmatrix}$$

③ no effect.

④ factor -1

Call this matrix B.

$$\det(B) = 2(1)(-3)(5) = -30$$

From our operations

$$\det(B) = \frac{1}{2}(-1) \det(A)$$

$$\begin{aligned} \Rightarrow \det(A) &= -2 \det(B) \\ &= -2(-30) = 60 \end{aligned}$$

## Some Theorems:

**Theorem:** The  $n \times n$  matrix  $A$  is invertible if and only if  $\det(A) \neq 0$ .

**Theorem:** For  $n \times n$  matrix  $A$ ,  $\det(A^T) = \det(A)$ .

**Theorem:** For  $n \times n$  matrices  $A$  and  $B$ ,  $\det(AB) = \det(A) \det(B)$ .

## Example

Show that if  $A$  is an  $n \times n$  invertible matrix, then

$$\det(A^{-1}) = \frac{1}{\det(A)}.$$

Suppose  $A^{-1}$  exists, then  $\det(A) \neq 0$  and

$$A^{-1}A = I. \quad \text{Hence} \quad \det(A^{-1}A) = \det(I) = 1$$

From the last theorem on the previous slide

$$\det(A^{-1}) \det(A) = 1$$

$$\Rightarrow \det(A^{-1}) = \frac{1}{\det(A)}$$



## Example

Let  $A$  be an  $n \times n$  matrix, and suppose there exists invertible matrix  $P$  such that

$$B \doteq P^{-1}AP.$$

Show that

$$\det(B) = \det(A).$$

$$\begin{aligned}\det(B) &= \det(P^{-1}AP) \\ &= \det(P^{-1}) \det(A) \det(P) \\ &= \det(P^{-1}) \det(P) \det(A) \\ &= \underbrace{\frac{1}{\det(P)} \det(P)}_1 \det(A) \\ &= \det(A)\end{aligned}$$

*these  
are  
scalars*