

Section 4.4: Coordinate Systems

Definition: Coordinate Vectors

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be an **ordered** basis of the vector space V . For each \mathbf{x} in V we define the **coordinate vector of \mathbf{x} relative to the basis \mathcal{B}** to be the unique vector (c_1, \dots, c_n) in \mathbb{R}^n whose entries are the weights $\mathbf{x} = c_1\mathbf{b}_1 + \dots + c_n\mathbf{b}_n$.

We'll use the notation $[\mathbf{x}]_{\mathcal{B}}$; that is $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$.

Different Coordinates on \mathbb{R}^n

We can use alternative coordinate systems to describe vectors in \mathbb{R}^n .

Definition

Given an ordered basis \mathcal{B} in \mathbb{R}^n , the matrix $P_{\mathcal{B}} = [\mathbf{b}_1 \ \mathbf{b}_2 \ \cdots \ \mathbf{b}_n]$ is called the **change of coordinates matrix** for the basis \mathcal{B} (or from the basis \mathcal{B} to the standard basis).

Theorem

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be an ordered basis of \mathbb{R}^n . Then the change of coordinate mapping $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ is the one to one linear transformation from \mathbb{R}^n onto \mathbb{R}^n defined by

$$[\mathbf{x}]_{\mathcal{B}} = P_{\mathcal{B}}^{-1} \mathbf{x}$$

$$[\mathbf{x}]_{\mathcal{B}} = P_{\mathcal{B}}^{-1} [\mathbf{x}]_{\mathcal{B}}$$

Theorem: Coordinate Mapping

Theorem

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be an ordered basis for a vector space V . Then the coordinate mapping $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ is a **one to one** mapping of V **onto** \mathbb{R}^n .

Remark:

- ▶ For a vector space V , if there exists a coordinate mapping from $V \rightarrow \mathbb{R}^n$, we say that V is **isomorphic** to \mathbb{R}^n .
- ▶ Properties of subsets of V , such as linear dependence, can be discerned from the coordinate vectors in \mathbb{R}^n .

\mathbb{P}_3 is Isomorphic to \mathbb{R}^4

We saw that using the ordered basis $\mathcal{B} = \{1, t, t^2, t^3\}$ that any vector

$$\mathbf{p}(t) = p_0 + p_1 t + p_2 t^2 + p_3 t^3$$

in \mathbb{P}_3 has coordinate vector

$$[\mathbf{p}]_{\mathcal{B}} = \begin{bmatrix} p_0 \\ p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

in \mathbb{R}^4 .

Example

Use coordinate vectors to determine if the set $\{\mathbf{p}, \mathbf{q}, \mathbf{r}\}$ is linearly dependent or independent in \mathbb{P}_2 .

$$\mathbf{p}(t) = 1 - 2t^2, \quad \mathbf{q}(t) = 3t + t^2, \quad \mathbf{r}(t) = 1 + t$$

We need a basis of \mathbb{P}_2 to define coordinate vectors. Let's use $\mathcal{B} = \{1, t, t^2\}$ in this order.

$$[\vec{p}]_{\mathcal{B}} = \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}, \quad [\vec{q}]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 3 \\ 1 \end{bmatrix}, \quad [\vec{r}]_{\mathcal{B}} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

We can use a matrix to determine if

these vectors in \mathbb{R}^3 are lin. independent.

$$\text{Let } A = \left[\begin{array}{ccc} [\vec{p}]_{\mathcal{B}} & [\vec{q}]_{\mathcal{B}} & [\vec{r}]_{\mathcal{B}} \end{array} \right]$$
$$= \begin{bmatrix} 1 & 0 & 1 \\ 0 & 3 & 1 \\ -2 & 1 & 0 \end{bmatrix}$$

We can use a determinant. (across row 1)

$$\det(A) = 1 \begin{vmatrix} 3 & 1 \\ 1 & 0 \end{vmatrix} - 0 \begin{vmatrix} \dots & \dots \end{vmatrix} + 1 \begin{vmatrix} 0 & 3 \\ -2 & 1 \end{vmatrix}$$

$$= 1(0-1) + 1(0+6) = -1 + 6 = 5.$$

Since $\det(A) \neq 0$, the columns of A

are lin. independent.

$\{ [\vec{p}]_{\mathcal{B}}, [\vec{q}]_{\mathcal{B}}, [\vec{r}]_{\mathcal{B}} \}$ is

linearly independent in \mathbb{R}^3 , so

$\{ \vec{p}, \vec{q}, \vec{r} \}$ is lin. independent

in \mathbb{P}_2 .