

Section 13: The Laplace Transform

A quick word about functions of 2-variables:

Suppose $G(s, t)$ is a function of two independent variables (s and t) defined over some rectangle in the plane $a \leq t \leq b$, $c \leq s \leq d$. If we compute an integral with respect to one of these variables, say t ,

$$\int_{\alpha}^{\beta} G(s, t) dt$$

- ▶ the result is a function of the remaining variable s , and
- ▶ the variable s is treated as a constant while integrating with respect to t .

For Example...

Assume that $s \neq 0$ and $b > 0$. Compute the integral

$$\begin{aligned}\int_0^b e^{-st} dt &= \frac{1}{-s} e^{-st} \Big|_0^b = \frac{1}{-s} e^{-sb} - \frac{1}{-s} e^{-s(0)} \\ &= \frac{1}{s} - \frac{1}{s} e^{-sb}\end{aligned}$$

Treat s like a constant

Integral Transform

An **integral transform** is a mapping that assigns to a function $f(t)$ another function $F(s)$ via an integral of the form

$$\int_a^b K(s, t)f(t) dt.$$

- ▶ The function K is called the **kernel** of the transformation.
- ▶ The limits a and b may be finite or infinite.
- ▶ The integral may be improper so that convergence/divergence must be considered.
- ▶ This transform is **linear** in the sense that

$$\int_a^b K(s, t)(\alpha f(t) + \beta g(t)) dt = \alpha \int_a^b K(s, t)f(t) dt + \beta \int_a^b K(s, t)g(t) dt.$$

The Laplace Transform

Definition: Let $f(t)$ be defined on $[0, \infty)$. The Laplace transform of f is denoted and defined by

$$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt = F(s).$$

The domain of the transformation $F(s)$ is the set of all s such that the integral is convergent.

Note: The **kernel** for the Laplace transform is $K(s, t) = e^{-st}$.

Limits at Infinity e^{-st}

If $s > 0$, evaluate

$$\lim_{t \rightarrow \infty} e^{-st} = 0$$

$$-st < 0 \quad -st \rightarrow -\infty$$

If $s < 0$, evaluate

$$\lim_{t \rightarrow \infty} e^{-st} = \infty$$

$$-st > 0 \quad -st \rightarrow +\infty$$

Find¹ the Laplace transform of $f(t) = 1$.

$$\mathcal{L}\{1\} = \int_0^{\infty} e^{-st} 1 dt$$

Suppose $s=0$ the integral is

$$\int_0^{\infty} 1 dt = \lim_{b \rightarrow \infty} \int_0^b dt = \lim_{b \rightarrow \infty} t \Big|_0^b = \lim_{b \rightarrow \infty} b = \infty$$

The integral diverges \Rightarrow zero is not in the domain of $\mathcal{L}\{1\}$.

Suppose $s \neq 0$

$$\mathcal{L}\{1\} = \int_0^{\infty} e^{-st} dt$$

¹Unless stated otherwise, the domain for each example is $[0, \infty)$.

$$= \lim_{b \rightarrow \infty} \int_0^b e^{-st} dt$$

$$= \lim_{b \rightarrow \infty} \frac{1}{s} - \frac{1}{s} e^{-sb}$$

This only converges if $s > 0$

for $s > 0$ $\mathcal{L}\{1\} = \frac{1}{s} - 0 = \frac{1}{s}$

$$\mathcal{L}\{1\} = \frac{1}{s} \text{ for } s > 0$$

Find the Laplace transform of $f(t) = t$.

$$\mathcal{L}\{t\} = \int_0^{\infty} e^{-st} t dt$$

If $s=0$, the integral is

$$\int_0^{\infty} t dt = \infty \text{ divergent.}$$

For $s \neq 0$

$$\mathcal{L}\{t\} = \int_0^{\infty} e^{-st} t dt$$

$$u = t, \quad du = dt$$

$$dv = e^{-st} dt$$

$$v = -\frac{1}{s} e^{-st}$$

$$= \frac{-1}{s} e^{-st} t \Big|_0^{\infty} - \int_0^{\infty} \frac{-1}{s} e^{-st} dt$$

for $s > 0$, $\lim_{t \rightarrow \infty} e^{-st} t = 0$

for $s > 0$

$$\mathcal{L}\{t\} = \frac{1}{s} \underbrace{\int_0^{\infty} e^{-st} dt}_{\mathcal{L}\{1\}}$$


$$= \frac{1}{s} \left(\frac{1}{s} \right) = \frac{1}{s^2}$$

$$\mathcal{L}\{t\} = \frac{1}{s^2}, \quad s > 0$$

A piecewise defined function

Find the Laplace transform of f defined by

$$f(t) = \begin{cases} 2t, & 0 \leq t < 10 \\ 0, & t \geq 10 \end{cases}$$



$$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

$$= \int_0^{10} e^{-st} f(t) dt + \int_{10}^{\infty} e^{-st} f(t) dt$$

$$= \int_0^{10} e^{-st} (2t) dt + \int_{10}^{\infty} e^{-st} (0) dt$$

When $s=0$, we get

$$\int_0^{10} 2t \, dt = t^2 \Big|_0^{10} = 100$$

When $s \neq 0$, we set

$$2 \int_0^{10} e^{-st} t \, dt$$
$$= 2 \left(\frac{-1}{s} t e^{-st} \Big|_0^{10} - \int_0^{10} \frac{-1}{s} e^{-st} \, dt \right)$$

$$u = t, \quad du = dt$$

$$dv = e^{-st} \, dt$$

$$v = \frac{-1}{s} e^{-st}$$

$$= 2 \left(\frac{-1}{s} (10) e^{-10s} - 0 + \frac{1}{s} \left(\frac{-1}{s} e^{-st} \right) \Big|_0^{10} \right)$$

$$= 2 \left(\frac{-10}{s} e^{-10s} - \frac{1}{s^2} (e^{-10s} - e^0) \right)$$

$$= \frac{2}{s^2} - \frac{2}{s^2} e^{-10s} - \frac{20}{s} e^{-10s}$$

$$f(t) = \begin{cases} 2t, & 0 \leq t < 10 \\ 0, & t \geq 10 \end{cases}$$

$$\mathcal{L}\{f(t)\} = \begin{cases} \frac{2}{s^2} - \frac{2}{s^2} e^{-10s} - \frac{20}{s} e^{-10s}, & s \neq 0 \\ 100, & s = 0 \end{cases}$$

The Laplace Transform is a Linear Transformation

Some basic results include:

$$\blacktriangleright \mathcal{L}\{\alpha f(t) + \beta g(t)\} = \alpha F(s) + \beta G(s)$$

$$\blacktriangleright \mathcal{L}\{1\} = \frac{1}{s}, \quad s > 0$$

$$\blacktriangleright \mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}, \quad s > 0 \text{ for } n = 1, 2, \dots$$

$$\blacktriangleright \mathcal{L}\{e^{at}\} = \frac{1}{s-a}, \quad s > a$$

$$\blacktriangleright \mathcal{L}\{\cos kt\} = \frac{s}{s^2+k^2}, \quad s > 0$$

$$\blacktriangleright \mathcal{L}\{\sin kt\} = \frac{k}{s^2+k^2}, \quad s > 0$$

Evaluate the Laplace transform $\mathcal{L}\{f(t)\}$ if

$$\mathcal{L}\{\cos kt\} = \frac{s}{s^2+k^2}, \quad s > 0$$

(a) $f(t) = \cos(\pi t)$

Here $k = \pi$

$$\mathcal{L}\{\cos(\pi t)\} = \frac{s}{s^2 + \pi^2}$$

Evaluate the Laplace transform $\mathcal{L}\{f(t)\}$ if

$$\mathcal{L}\{1\} = \frac{1}{s}, \quad s > 0$$

(b) $f(t) = 2t^4 - e^{-5t} + 3$

$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}, \quad s > 0 \text{ for } n \geq 0$$

$$\mathcal{L}\{e^{at}\} = \frac{1}{s-a}, \quad s > a$$

$$\mathcal{L}\{2t^4 - e^{-5t} + 3\}$$

$$= 2\mathcal{L}\{t^4\} - \mathcal{L}\{e^{-5t}\} + 3\mathcal{L}\{1\}$$

$$= 2 \frac{4!}{s^{4+1}} - \frac{1}{s - (-5)} + 3 \frac{1}{s}$$

$$= \frac{48}{s^5} - \frac{1}{s+5} + \frac{3}{s} \quad \text{for } s > 0$$