

# March 8 Math 3260 sec. 52 Spring 2024

## Section 4.1: Vector Spaces and Subspaces

### Definition: Vector Space

A **vector space** is a nonempty set  $V$  of objects called *vectors* together with two operations called *vector addition* and *scalar multiplication* that satisfy the following ten axioms:

For all  $\mathbf{u}, \mathbf{v}$ , and  $\mathbf{w}$  in  $V$ , and for any scalars  $c$  and  $d$

1. The sum  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
2.  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ .
4. There exists a **zero** vector  $\mathbf{0}$  in  $V$  such that  $\mathbf{u} + \mathbf{0} = \mathbf{u}$ .
5. For each vector  $\mathbf{u}$  there exists a vector  $-\mathbf{u}$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
6. For each scalar  $c$ ,  $c\mathbf{u}$  is in  $V$ .
7.  $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8.  $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ .
9.  $c(d\mathbf{u}) = d(c\mathbf{u}) = (cd)\mathbf{u}$ .
10.  $1\mathbf{u} = \mathbf{u}$

## An Example of a Vector Space: “P two”

### “P two”

$$\mathbb{P}_2 = \left\{ \mathbf{p}(t) = p_0 + p_1 t + p_2 t^2 \mid p_0, p_1, p_2 \in \mathbb{R} \right\}.$$

Consider  $t$  to be some real variable, and consider the scalars to be  $\mathbb{R}$ . Let  $\mathbb{P}_2$  be the set of all polynomials with real coefficients of degree at most two.

Examples of elements of  $\mathbb{P}_2$  include things like

$$\mathbf{p}(t) = 1 + t - 3t^2, \quad \mathbf{q}(t) = -2 + 5t + 12t^2, \quad \text{and} \quad \mathbf{r}(t) = \pi + \frac{1}{\pi}t.$$

**Remark:** It doesn't make sense to state that  $\mathbb{P}_2$  is a vector space until we define **scalar multiplication** and **vector addition**.

## An Example of a Vector Space: “P two”

Let  $\mathbf{p}(t) = p_0 + p_1t + p_2t^2$  and  $\mathbf{q}(t) = q_0 + q_1t + q_2t^2$  be polynomials in  $\mathbb{P}_2$  and  $c$  be a scalar. We define the two operations as follows:

$$\text{Scalar Multiplication: } (c\mathbf{p})(t) = c\mathbf{p}(t) = cp_0 + cp_1t + cp_2t^2.$$

$$\begin{aligned} \text{Vector Addition: } (\mathbf{p} + \mathbf{q})(t) &= \mathbf{p}(t) + \mathbf{q}(t) \\ &= (p_0 + q_0) + (p_1 + q_1)t + (p_2 + q_2)t^2. \end{aligned}$$

**Remark:** It can be shown that  $\mathbb{P}_2$  with these operations satisfies the ten vector space axiom. Note, this means that

the polynomials ARE vectors.

## Example

$$\mathbf{p}(t) = 1 + t - 3t^2, \quad \mathbf{q}(t) = -2 + 5t + 12t^2, \quad \text{and} \quad \mathbf{r}(t) = \pi + \frac{1}{\pi}t.$$

Evaluate

$$1. (\mathbf{p} + \mathbf{q})(t) = \vec{p}(t) + \vec{q}(t) = (1-2) + (1+5)t + (-3+12)t^2 \\ = -1 + 6t + 9t^2$$

$$2. (-1\mathbf{r})(t) = -1\vec{r}(t) = -1\pi + (-1)\frac{1}{\pi}t = -\pi - \frac{1}{\pi}t$$

$$3. (-1\mathbf{r} + \mathbf{r})(t) = -1\vec{r}(t) + \vec{r}(t) = (-\pi + \pi) + \left(-\frac{1}{\pi} + \frac{1}{\pi}\right)t \\ = 0 + 0t$$

## The Zero Vector in $\mathbb{P}_2$

Let  $\mathbf{0}(t) = a_0 + a_1 t + a_2 t^2$  be the zero vector in  $\mathbb{P}_2$ .

Use the property<sup>1</sup> in Axiom 4 to identify the values of the coefficients  $a_0$ ,  $a_1$ , and  $a_2$ .

Let  $\vec{p}(t) = p_0 + p_1 t + p_2 t^2$  be any vector in  $\mathbb{P}_2$ .

$$\begin{aligned}(\vec{0} + \vec{p})(t) &= \vec{0}(t) + \vec{p}(t) = (a_0 + p_0) + (a_1 + p_1)t + (a_2 + p_2)t^2 \\ &= p_0 + p_1 t + p_2 t^2\end{aligned}$$

This requires

$$a_0 + p_0 = p_0 \Rightarrow a_0 = 0$$

---

<sup>1</sup>Axiom 4 says that  $\mathbf{p} + \mathbf{0} = \mathbf{p}$  for every vector  $\mathbf{p}$  in  $\mathbb{P}_2$ .

$$a_1 + p_1 = p_1 \Rightarrow a_1 = 0$$

$$a_2 + p_2 = p_2 \Rightarrow a_2 = 0.$$

That is,  $a_0 = a_1 = a_2 = 0$  so

$$\vec{0}(t) = 0 + 0t + 0t^2$$

$\mathbb{P}_n$ 

For an integer  $n \geq 0$ , let  $\mathbb{P}_n$  denote the set of all polynomials with real coefficients of degree at most  $n$ .

$$\mathbb{P}_n = \left\{ \mathbf{p}(t) = p_0 + p_1 t + \cdots + p_n t^n \mid p_0, p_1, \dots, p_n \in \mathbb{R} \right\},$$

For  $\mathbf{p}$  and  $\mathbf{q}$  in  $\mathbb{P}_n$  and scalar  $c$ , define scalar multiplication and vector addition by

$$(c\mathbf{p})(t) = c\mathbf{p}(t) = cp_0 + cp_1 t + \cdots + cp_n t^n,$$

$$(\mathbf{p} + \mathbf{q})(t) = \mathbf{p}(t) + \mathbf{q}(t) = (p_0 + q_0) + (p_1 + q_1)t + \cdots + (p_n + q_n)t^n.$$


**Remark:** It can readily be shown that the zero vector  $\mathbf{0}(t) = 0 + 0t + \cdots + 0t^n$ , and the additive inverse of  $\mathbf{p}(t) = p_0 + p_1 t + \cdots + p_n t^n$  is

$$-\mathbf{p}(t) = -p_0 - p_1 t - \cdots - p_n t^n.$$

## A set that is not a Vector Space

Let  $S = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid x \leq 0, y \leq 0 \right\}$  with regular vector addition and scalar multiplication in  $\mathbb{R}^2$ .

Geometrically, what is the set  $S$ ?



$S$  is the third quadrant including the negative  $x$  and  $y$  axes.



## A set that is not a Vector Space

$S = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid x \leq 0, y \leq 0 \right\}$  with regular vector addition and scalar multiplication in  $\mathbb{R}^2$ .

Does Axiom<sup>2</sup> 1. hold for  $S$ ?

Let  $\vec{u} = \begin{bmatrix} x \\ y \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} u \\ v \end{bmatrix}$  be in  $S$ .

Then  $x \leq 0$ ,  $y \leq 0$ ,  $u \leq 0$ , and  $v \leq 0$ .


$\vec{u} + \vec{v} = \begin{bmatrix} x+u \\ y+v \end{bmatrix}$ . Is  $\vec{u} + \vec{v}$  in  $S$ ?

Yes,  $x+u \leq 0$  and  $y+v \leq 0$  so  $\vec{u} + \vec{v} \in S$ .

---

<sup>2</sup>Axiom 1 is closure under vector addition.

$S$  is closed under vector addition.



## A set that is not a Vector Space

$S = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \mid x \leq 0, y \leq 0 \right\}$  with regular vector addition and scalar multiplication in  $\mathbb{R}^2$ .

Does Axiom<sup>3</sup> 6. hold for  $S$ ?


Consider  $\vec{u} = \begin{bmatrix} -1 \\ -3 \end{bmatrix}$  which is in  $S$ .

$-2\vec{u} = \begin{bmatrix} 2 \\ 6 \end{bmatrix}$  is not in  $S$ .

$S$  is not closed under scalar mult.

---

<sup>3</sup>Axiom 6 is closure under scalar multiplication.



# Some Algebraic Properties of Vector Spaces

## Theorem

Let  $V$  be a vector space. For each  $\mathbf{u}$  in  $V$  and scalar  $c$

(i)  $0\mathbf{u} = \mathbf{0}$

(ii)  $c\mathbf{0} = \mathbf{0}$

(iii)  $-1\mathbf{u} = -\mathbf{u}$

Proof of (i):

Let  $\vec{u} \in V$  be arbitrary. Note that the scalar  $0 = 0 + 0$ .

$$0\vec{u} = (0+0)\vec{u}$$

$$= \vec{0} + \vec{0}\vec{u}$$

By axiom 5, there exists a vector  $-\vec{0}\vec{u}$ .

Add this to both sides.

$$-\vec{0}\vec{u} + \vec{0}\vec{u} = -\vec{0}\vec{u} + \vec{0}\vec{u} + \vec{0}\vec{u}$$

$$\vec{0} = \vec{0} + \vec{0}\vec{u}$$

$$\vec{0} = \vec{0}\vec{u}.$$

This is the required conclusion.

# Subspaces

## Definition:

A **subspace** of a vector space  $V$  is a subset  $H$  of  $V$  for which

- a) The zero vector is in<sup>a</sup>  $H$
- b)  $H$  is closed under vector addition. (i.e.  $\mathbf{u}, \mathbf{v}$  in  $H$  implies  $\mathbf{u} + \mathbf{v}$  is in  $H$ )
- c)  $H$  is closed under scalar multiplication. (i.e.  $\mathbf{u}$  in  $H$  implies  $c\mathbf{u}$  is in  $H$ )

---

<sup>a</sup>This is sometimes replaced with the condition that  $H$  is nonempty.

**Remark:** A subspace is a vector space. If these three properties hold, it inherits the structure from its parent space.