

Section 15: Shift Theorems

We saw the first translation theorem.

Theorem: Shift in s

Suppose $\mathcal{L}\{f(t)\} = F(s)$. Then for any real number a

$$\mathcal{L}\{e^{at}f(t)\} = F(s - a).$$

Equivalently,

$$\mathcal{L}^{-1}\{F(s - a)\} = e^{at}\mathcal{L}^{-1}\{F(s)\}.$$

We'll see a similar result relating to translations in the variable t . This requires some preliminary ideas.


The Unit Step Function

Definition: Unit Step Function

Let $a > 0$. The unit step function *centered at a* is denoted $\mathcal{U}(t - a)$. It is defined by

$$\mathcal{U}(t - a) = \begin{cases} 0, & 0 \leq t < a \\ 1, & t \geq a \end{cases}$$

The name *unit step function* is derived from its graph which looks like a stair step of height 1.



we'll define
 $u(t) = u(t - 0)$
 $= 1$ for
 $t \geq 0$

Figure: A graph of $\mathcal{U}(t - a)$ which jumps from zero to one at $t = a$.

Piecewise Defined Functions

Verify that

$$f(t) = \begin{cases} g(t), & 0 \leq t < a \\ h(t), & t \geq a \end{cases} \quad a > 0$$
$$= g(t) - g(t)\mathcal{U}(t-a) + h(t)\mathcal{U}(t-a)$$

Consider t values such that $0 \leq t < a$.

Then $\mathcal{U}(t-a) = 0$. Then

$$g(t) - g(t)\mathcal{U}(t-a) + h(t)\mathcal{U}(t-a) =$$

$$g(t) - g(t) \cdot 0 + h(t) \cdot 0 = g(t)$$

so this is $f(t)$ for $0 \leq t < a$

$$\begin{cases} g(t), & 0 \leq t < a \\ h(t), & t \geq a \end{cases} = g(t) - g(t)\mathcal{U}(t-a) + h(t)\mathcal{U}(t-a)$$

If $t \geq a$, then $\mathcal{U}(t-a) = 1$

Then

$$g(t) - g(t)\mathcal{U}(t-a) + h(t)\mathcal{U}(t-a) =$$

$$g(t) - g(t) \cdot 1 + h(t) \cdot 1 = h(t)$$

So this is $f(t)$ when $t \geq a$.

$$\text{Example } f(t) = \begin{cases} e^t, & 0 \leq t < 2 \\ t^2, & 2 \leq t < 5 \\ 2t & t \geq 5 \end{cases}$$

Rewrite the function f in terms of the unit step function.

$$f(t) = e^t - e^t u(t-2) + t^2 u(t-2) - t^2 u(t-5) + 2t u(t-5)$$

Check: for $0 \leq t < 2$, $u(t-2) = 0$ $u(t-5) = 0$

$$f(t) = e^t - e^t \cdot 0 + t^2 \cdot 0 - t^2 \cdot 0 + 2t \cdot 0 = e^t$$

For $2 \leq t < 5$, $u(t-2) = 1$ $u(t-5) = 0$

$$f(t) = e^t - e^t \cdot 1 + t^2 \cdot 1 - t^2 \cdot 0 + 2t \cdot 0 = t^2$$

For $t \geq 5$, $u(t-2) = 1$ $u(t-5) = 1$

$$f(t) = e^t - e^t \cdot 1 + t^2 \cdot 1 - t^2 \cdot 1 + 2t \cdot 1 = 2t$$

Find $\mathcal{L}\{u(t-a)\}$ assume $a > 0$

By definition

$$\mathcal{L}\{u(t-a)\} = \int_0^{\infty} e^{-st} u(t-a) dt$$

$$u(t-a) = \begin{cases} 0, & 0 \leq t < a \\ 1, & t \geq a \end{cases}$$

$$= \int_0^a e^{-st} \cdot 0 dt + \int_a^{\infty} e^{-st} \cdot 1 dt$$

for $s > 0$

$$= \frac{-1}{s} e^{-st} \Big|_a^{\infty} = \frac{-1}{s} (0 - e^{-as})$$

$$= \frac{1}{s} e^{-as}$$

Translation in t

Given a function $f(t)$ for $t \geq 0$, and a number $a > 0$

$$f(t-a)\mathcal{U}(t-a) = \begin{cases} 0, & 0 \leq t < a \\ f(t-a), & t \geq a \end{cases}.$$




Figure: The function $f(t-a)\mathcal{U}(t-a)$ has the graph of f shifted a units to the right with value of zero for t to the left of a .

Theorem (translation in t)

If $F(s) = \mathcal{L}\{f(t)\}$ and $a > 0$, then

$$\mathcal{L}\{f(t-a)\mathcal{U}(t-a)\} = e^{-as}F(s).$$

A special case is $f(t) = 1$. We just found

$$\mathcal{L}\{\mathcal{U}(t-a)\} = e^{-as}\mathcal{L}\{1\} = \frac{e^{-as}}{s}.$$

We can state this in terms of the inverse transform as

$$\mathcal{L}^{-1}\{e^{-as}F(s)\} = f(t-a)\mathcal{U}(t-a).$$