

Section 18: Sine and Cosine Series

This section has two topics

- ▶ Fourier series of functions on $(-p, p)$ that have symmetry, and
- ▶ half-range sine and cosine series for functions defined on $(0, p)$.

Functions with Symmetry

Recall some definitions:

Suppose f is defined on an interval containing x and $-x$.

If $f(-x) = f(x)$ for all x , then f is said to be **even**.

If $f(-x) = -f(x)$ for all x , then f is said to be **odd**.

For example, $f(x) = x^n$ is even if n is even and is odd if n is odd. The trigonometric function $g(x) = \cos x$ is even, and $h(x) = \sin x$ is odd.

Even and Odd Symmetry




Figure: Graphical interpretation of even and odd symmetry.

Integrals on symmetric intervals

If f is an even function on $(-p, p)$, then

$$\int_{-p}^p f(x) dx = 2 \int_0^p f(x) dx.$$

If f is an odd function on $(-p, p)$, then

$$\int_{-p}^p f(x) dx = 0.$$

Products of Even and Odd functions

$$\text{Even} \times \text{Even} = \text{Even},$$

and

$$\text{Odd} \times \text{Odd} = \text{Even}.$$

While

$$\text{Even} \times \text{Odd} = \text{Odd}.$$

So, suppose f **is even** on $(-p, p)$. This tells us that $f(x) \cos(nx)$ is **even** for all n and $f(x) \sin(nx)$ is **odd** for all n .

And, if f **is odd** on $(-p, p)$. This tells us that $f(x) \sin(nx)$ is **even** for all n and $f(x) \cos(nx)$ is **odd** for all n .

Fourier Series of an Even Function

If f is even on $(-p, p)$, then the Fourier series of f has only constant and cosine terms. Moreover

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{p}\right)$$

where

$$a_0 = \frac{2}{p} \int_0^p f(x) dx$$

and

$$a_n = \frac{2}{p} \int_0^p f(x) \cos\left(\frac{n\pi x}{p}\right) dx.$$

$b_n = 0$
for all n

Fourier Series of an Odd Function

If f is odd on $(-p, p)$, then the Fourier series of f has only sine terms. Moreover

$$f(x) = \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{p}\right)$$

where

$$b_n = \frac{2}{p} \int_0^p f(x) \sin\left(\frac{n\pi x}{p}\right) dx.$$


$a_0 = 0$
 $a_n = 0$ for all n

Find the Fourier series of f

$$f(x) = \begin{cases} x + \pi, & -\pi < x < 0 \\ \pi - x, & 0 \leq x < \pi \end{cases}$$

$$y = x + \pi \\ n=1, b=\pi$$

$$y = \pi - x \\ n=-1, b=\pi$$



Symmetry can be determined by evaluating $f(-x)$.
or by graphing

$$P = \pi$$

$$\frac{n\pi x}{P} = \frac{n\pi x}{\pi} = nx$$

Find a_0 and a_n

$$a_0 = \frac{2}{\pi} \int_0^{\pi} f(x) dx = \frac{2}{\pi} \int_0^{\pi} (\pi - x) dx$$

$$= \frac{2}{\pi} \left[\pi x - \frac{x^2}{2} \right]_0^{\pi} = \frac{2}{\pi} \left[\pi^2 - \frac{\pi^2}{2} - 0 \right] = \frac{2}{\pi} \left[\frac{\pi^2}{2} \right]$$

$$a_0 = \pi$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos(nx) dx$$

$$= \frac{2}{\pi} \int_0^{\pi} (\pi - x) \cos(nx) dx$$

Int by parts $u = \pi - x$, $du = -dx$

$dv = \cos(nx) dx$, $v = \frac{1}{n} \sin(nx)$

$$a_n = \frac{2}{\pi} \left[\frac{\pi-x}{n} \sin(nx) \right]_0^\pi + \int_0^\pi \frac{1}{n} \sin(nx) dx$$

0

$$= \frac{2}{\pi} \left[\frac{-1}{n^2} \cos(nx) \right]_0^\pi$$

$$= -\frac{2}{n^2 \pi} [\cos(n\pi) - \cos(0)]$$

$$= \frac{-2}{n^2 \pi} [(-1)^n - 1]$$

$$a_n = \frac{2}{n^2 \pi} (1 - (-1)^n)$$


$$a_0 = \pi$$

The series is

$$f(x) = \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{2}{n^2\pi} (1 - (-1)^n) \cos(nx)$$

for $f(x) = \begin{cases} x + \pi, & -\pi < x < 0 \\ \pi - x, & 0 \leq x < \pi \end{cases}$

Let's plot the series over the interval
 $-3\pi < x < 3\pi$



Taking Advantage of Symmetry

$$f(x) = \begin{cases} x + \pi, & -\pi < x < 0 \\ \pi - x, & 0 \leq x < \pi \end{cases}$$

Notice the amount of work we saved by using the symmetry. The formulas for the coefficients give

$$a_0 = \frac{1}{\pi} \int_{-\pi}^0 (x + \pi) dx + \frac{1}{\pi} \int_0^{\pi} (\pi - x) dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^0 (x + \pi) \cos(nx) dx + \frac{1}{\pi} \int_0^{\pi} (\pi - x) \cos(nx) dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^0 (x + \pi) \sin(nx) dx + \frac{1}{\pi} \int_0^{\pi} (\pi - x) \sin(nx) dx$$

We only had to compute the integrals in **green**.

Half Range Sine and Half Range Cosine Series

Suppose f is only defined for $0 < x < p$. We can **extend** f to the left, to the interval $(-p, 0)$, as either an even function or as an odd function. Then we can express f with **two distinct** series.




Figure: Some function f shown in dark blue with even and odd extensions.

Half Range Cosine Series

For f defined on $(0, p)$ we can *pretend* that f is defined on $(-p, p)$ by setting

$$f(-x) = f(x) \quad 0 < x < p.$$

Then we can define the

Half range cosine series
$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{p}\right)$$

where
$$a_0 = \frac{2}{p} \int_0^p f(x) dx \quad \text{and} \quad a_n = \frac{2}{p} \int_0^p f(x) \cos\left(\frac{n\pi x}{p}\right) dx.$$

Extending a Function to be Even




Figure: $f(x) = p - x$, $0 < x < p$ together with its **even** extension.

Half Range Sine

Suppose f is defined on $(0, p)$. We can extend f to be defined on $(-p, p)$ by setting

$$f(-x) = -f(x) \quad 0 < x < p.$$

Then we can define the

Half range sine series $f(x) = \sum_{n=1}^{\infty} b_n \sin \left(\frac{n\pi x}{p} \right)$

where $b_n = \frac{2}{p} \int_0^p f(x) \sin \left(\frac{n\pi x}{p} \right) dx.$

Extending a Function to be Odd




Figure: $f(x) = p - x$, $0 < x < p$ together with its **odd** extension.

Find the Half Range Sine Series of f

$$f(x) = 2 - x, \quad 0 < x < 2$$

$$P = 2, \quad \frac{n\pi x}{P} = \frac{n\pi x}{2}$$

$$f(x) = \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{2}\right)$$

$\frac{2}{P}$



$$b_n = \frac{2}{2} \int_0^2 f(x) \sin\left(\frac{n\pi x}{2}\right) dx$$

$$= \int_0^2 (2-x) \sin\left(\frac{n\pi x}{2}\right) dx$$

Int by parts $u = z - x$, $du = -dx$

$$dv = \sin\left(\frac{n\pi x}{z}\right) dx \quad v = -\frac{z}{n\pi} \cos\left(\frac{n\pi x}{z}\right)$$

$$b_n = \left. -\frac{z}{n\pi} (z-x) \cos\left(\frac{n\pi x}{z}\right) \right|_0^z - \int_0^z \frac{+z}{n\pi} \cos\left(\frac{n\pi x}{z}\right) (+1) dx$$

$$= -\frac{z}{n\pi} (z-z) \cos(n\pi) - \frac{z}{n\pi} (z-0) \cos(0)$$

$$= \frac{z}{n\pi} (z) = \frac{4}{n\pi}$$

$$b_n = \frac{4}{n\pi}$$

The series is

$$f(x) = \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin\left(\frac{n\pi x}{2}\right)$$