

November 19 Math 2306 sec. 52 Fall 2021

Section 18: Sine and Cosine Series

We consider a function f defined on the interval $(0, p)$ and extend it to be even or odd.




Figure: Some function f shown in dark blue with even and odd extensions.

Half Range Cosine Series

For f defined on $(0, p)$ we can *pretend* that f is defined on $(-p, p)$ by setting

$$f(-x) = f(x) \quad 0 < x < p.$$

Then we can define the

Half range cosine series
$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{p}\right)$$

where
$$a_0 = \frac{2}{p} \int_0^p f(x) dx \quad \text{and} \quad a_n = \frac{2}{p} \int_0^p f(x) \cos\left(\frac{n\pi x}{p}\right) dx.$$

Half Range Sine

Suppose f is defined on $(0, p)$. We can extend f to be defined on $(-p, p)$ by setting

$$f(-x) = -f(x) \quad 0 < x < p.$$

Then we can define the

Half range sine series
$$f(x) = \sum_{n=1}^{\infty} b_n \sin \left(\frac{n\pi x}{p} \right)$$

where
$$b_n = \frac{2}{p} \int_0^p f(x) \sin \left(\frac{n\pi x}{p} \right) dx.$$

Find the Half Range Sine Series of f

$$f(x) = 2 - x, \quad 0 < x < 2$$

We worked this out and found the half-range sine series

$$f(x) = \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin\left(\frac{n\pi x}{2}\right).$$

Find the Half Range Cosine Series of f

$$f(x) = 2 - x, \quad 0 < x < 2$$

← $P=2$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{2}\right)$$

$$a_0 = \frac{2}{2} \int_0^2 f(x) dx = \int_0^2 (2-x) dx$$

$$= 2x - \frac{x^2}{2} \Big|_0^2 = 4 - 2 = 2$$

$$a_n = \frac{2}{2} \int_0^2 f(x) \cos\left(\frac{n\pi x}{2}\right) dx = \int_0^2 (2-x) \cos\left(\frac{n\pi x}{2}\right) dx$$

Int by parts

$$u = 2-x, \quad du = -dx$$

$$dv = \cos\left(\frac{n\pi x}{2}\right) dx, \quad v = \frac{2}{n\pi} \sin\left(\frac{n\pi x}{2}\right)$$

$$a_n = \frac{2}{n\pi} (2-x) \sin\left(\frac{n\pi x}{2}\right) \Big|_0^2 + \int_0^2 \frac{2}{n\pi} \sin\left(\frac{n\pi x}{2}\right) dx$$

$$= \frac{2}{n\pi} \left(\frac{-2}{n\pi} \right) \cos\left(\frac{n\pi x}{2}\right) \Big|_0^2$$

$$= \frac{-4}{n^2\pi^2} \left(\cos(n\pi) - \cos(0) \right)$$

$$a_n = \frac{-4}{n^2 \pi^2} ((-1)^n - 1)$$

$$a_n = \frac{4}{n^2 \pi^2} (1 - (-1)^n) , \quad a_0 = 2$$

The half range cosine series for f

is

$$f(x) = 1 + \sum_{n=1}^{\infty} \frac{4}{n^2 \pi^2} (1 - (-1)^n) \cos\left(\frac{n\pi x}{2}\right)$$

Example Continued...

We have two different half range series:

$$\text{Half range sine: } f(x) = \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin\left(\frac{n\pi x}{2}\right)$$

$$\text{Half range cosine: } f(x) = 1 + \sum_{n=1}^{\infty} \frac{4(1 - (-1)^n)}{n^2\pi^2} \cos\left(\frac{n\pi x}{2}\right).$$

We have two different series representations for this function each of which converge to $f(x)$ on the interval $(0, 2)$. The following plots show graphs of f along with partial sums of each of the series. When we plot over the interval $(-2, 2)$ we see the two different symmetries. Plotting over a larger interval such as $(-6, 6)$ we can see the periodic extensions of the two symmetries.

Plots of f with Half range series




Figure: $f(x) = 2 - x$, $0 < x < 2$ with 10 terms of the sine series.

Plots of f with Half range series




Figure: $f(x) = 2 - x$, $0 < x < 2$ with 10 terms of the sine series, and the series plotted over $(-6, 6)$

Plots of f with Half range series




Figure: $f(x) = 2 - x$, $0 < x < 2$ with 5 terms of the cosine series.

Plots of f with Half range series





Figure: $f(x) = 2 - x$, $0 < x < 2$ with 5 terms of the cosine series, and the series plotted over $(-6, 6)$


Plot the Half-Range Sine Series

$$f(x) = 2 - x, \quad 0 < x < 2 \quad f(x) = \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin\left(\frac{n\pi x}{2}\right)$$



Plot the Half-Range Cosine Series

$$f(x) = 2 - x, \quad 0 < x < 2 \quad f(x) = 1 + \sum_{n=1}^{\infty} \frac{4(1 - (-1)^n)}{n^2 \pi^2} \cos\left(\frac{n\pi x}{2}\right)$$



Solution of a Differential Equation

An undamped spring mass system has a mass of 2 kg attached to a spring with spring constant 128 N/m. The mass is driven by an external force $f(t) = 2t$ for $-1 < t < 1$ that is 2-periodic so that $f(t+2) = f(t)$ for all $t > 0$. Determine a particular solution x_p for the displacement for $t > 0$.

undamped

$$mx'' + kx = f(t)$$

Here, $m=2$, $k=128$

$$2x'' + 128x = f(t)$$

$$\Rightarrow x'' + 64x = \frac{1}{2}f(t)$$

We'll express f as a Fourier Series.

From Nov. 15

$$f(t) = 2 \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

The ODE is

$$x'' + 64x = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

This is

$$x'' + 64x = \frac{2}{\pi} \sin(\pi t) - \frac{1}{\pi} \sin(2\pi t) + \frac{2}{3\pi} \sin(3\pi t) + \dots$$

Suppose x_p has a sine series

$$x_p = \sum_{n=1}^{\infty} B_n \sin(n\pi t)$$

This is the method of undetermined coefficients. Assume that x_p can be differentiated term by term.

$$\begin{aligned} x_p' &= \sum_{n=1}^{\infty} \frac{d}{dt} (B_n \sin(n\pi t)) \\ &= \sum_{n=1}^{\infty} B_n (n\pi) \cos(n\pi t) \end{aligned}$$

$$X_p'' = \sum_{n=1}^{\infty} -B_n (n\pi)^2 \sin(n\pi t)$$

$$X_p = \sum_{n=1}^{\infty} B_n \sin(n\pi t)$$

$$X_p'' + 64 X_p = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

$$\sum_{n=1}^{\infty} -B_n (n\pi)^2 \sin(n\pi t) + \sum_{n=1}^{\infty} 64 B_n \sin(n\pi t) = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

$$\sum_{n=1}^{\infty} (-(n\pi)^2 + 64) B_n \sin(n\pi t) = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

Matching coefficients

$$(64 - n^2\pi^2)B_n = \frac{2(-1)^{n+1}}{n\pi}$$

Note $64 - n^2\pi^2 \neq 0$ for all n

Hence

$$B_n = \frac{2(-1)^{n+1}}{n\pi(64 - n^2\pi^2)}$$

The particular solution

$$X_p = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi(64 - n^2\pi^2)} \sin(n\pi t)$$