November 19 Math 2306 sec. 54 Fall 2021 Section 18: Sine and Cosine Series

We consider a function f defined on the interval (0, p) and extend it to be even or odd.



Figure: Some function *f* shown in dark blue with even and odd extensions.

November 17, 2021

1/27

Half Range Cosine Series

For *f* defined on (0, p) we can *pretend* that *f* is defined on (-p, p) by setting

$$f(-x) = f(x) \quad 0 < x < p.$$

Then we can define the

Half range cosine series
$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{p}\right)$$

where $a_0 = \frac{2}{p} \int_0^p f(x) dx$ and $a_n = \frac{2}{p} \int_0^p f(x) \cos\left(\frac{n\pi x}{p}\right) dx$.

э

イロト イポト イヨト イヨト

Half Range Sine

Suppose *f* is defined on (0, p). We can extend *f* to be defined on (-p, p) by setting

$$f(-x) = -f(x) \quad 0 < x < p.$$

Then we can define the

Half range sine series $f(x) = \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{p}\right)$ where $b_n = \frac{2}{p} \int_0^p f(x) \sin\left(\frac{n\pi x}{p}\right) dx.$

Find the Half Range Sine Series of f

$$f(x) = 2 - x, \quad 0 < x < 2$$

We worked this out and found the half-range sine series

$$f(x) = \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin\left(\frac{n\pi x}{2}\right).$$

November 17, 2021 4/27

イロト イポト イヨト イヨト

Find the Half Range Cosine Series of f

- P=2 $f(x) = 2 - x, \quad 0 < x < 2$

$$f(x) = \frac{\alpha_0}{2} + \sum_{n=1}^{\infty} \alpha_n \cos\left(\frac{n\pi x}{2}\right)$$

$$Q_{p} = \frac{2}{z} \int_{0}^{z} f(x) dx = \int_{0}^{z} (z-x) dx$$

$$= 2x - \frac{x^2}{2} \Big|_{0}^{2} = 4 - 2 = 2$$

$$\begin{aligned}
\Omega_n &= \frac{2}{2} \int_{0}^{z} f(x) G_{5}\left(\frac{n\pi x}{2}\right) dx &= \int_{0}^{z} (z-x) G_{5}\left(\frac{n\pi x}{2}\right) dx \\
&= \frac{1}{2} \int_{0}^{z} f(x) G_{5}\left(\frac{n\pi x}{2}\right) dx
\end{aligned}$$

November 17, 2021 5/27

$$lut b_{\gamma} \text{ parts};$$

$$u = 2-x, \quad du = -dx$$

$$du = Cos\left(\frac{n\pi x}{2}\right) dx \quad V = \frac{2}{n\pi} \sin\left(\frac{n\pi x}{2}\right)$$

$$Q_{n} = \frac{2}{n\pi} \left(2-x\right) \sin\left(\frac{n\pi x}{2}\right) \Big|_{0}^{2} + \int_{0}^{2} \frac{2}{n\pi} \sin\left(\frac{n\pi x}{2}\right) dx$$

$$Q_{n} = \frac{2}{n\pi} \left(\frac{-2}{n\pi}\right) G_{s}\left(\frac{n\pi x}{2}\right) \Big|_{0}^{2}$$

$$= \frac{-4}{n^{2}\pi^{2}} \left(C_{os}\left(n\pi\right) - C_{s}(o)\right)$$

November 17, 2021 6/27

 $= \frac{-4}{n^2 \pi^2} ((-1)^2 - 1)$ $Q_{n} = \frac{4}{n^{2}\pi^{2}} \left(1 - (-1)^{n} \right), \quad Q_{0} = Z$



November 17, 2021 7/27

Example Continued...

We have two different half range series:

Half range sine:
$$f(x) = \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin\left(\frac{n\pi x}{2}\right)$$

Half range cosine: $f(x) = 1 + \sum_{n=1}^{\infty} \frac{4(1-(-1)^n)}{n^2\pi^2} \cos\left(\frac{n\pi x}{2}\right)$.

We have two different series representations for this function each of which converge to f(x) on the interval (0, 2). The following plots show graphs of *f* along with partial sums of each of the series. When we plot over the interval (-2, 2) we see the two different symmetries. Plotting over a larger interval such as (-6, 6) we can see the periodic extensions of the two symmetries.

Plots of *f* with Half range series



Plots of *f* with Half range series



November 17, 2021 15/27

Plots of f with Half range series



Plots of *f* with Half range series



November 17, 2021 17/27

Plot the Half-Range Sine Series

$$f(x) = 2 - x, \quad 0 < x < 2$$
 $f(x) = \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin\left(\frac{n\pi x}{2}\right)$



Plot the Half-Range Cosine Series

$$f(x) = 2 - x, \quad 0 < x < 2$$
 $f(x) = 1 + \sum_{n=1}^{\infty} \frac{4(1 - (-1)^n)}{n^2 \pi^2} \cos\left(\frac{n\pi x}{2}\right)$



Solution of a Differential Equation

An undamped spring mass system has a mass of 2 kg attached to a spring with spring constant 128 N/m. The mass is driven by an external force f(t) = 2t for -1 < t < 1 that is 2-periodic so that f(t+2) = f(t) for all t > 0. Determine a particular solution x_p for the displacement for t > 0.

For
$$\beta = 0$$
 (no domping) the ODP is
 $mx'' + kx = f(t)$
 $m = z, k = 128 \implies 2x'' + 128x = f(t)$
 $\implies x'' + 64x = \frac{1}{2} f(t)$

November 17, 2021 20/27

Le converte f as a Fourier Series. It will be a sine series.

From Nov 15,

$$f(t) = 2 \sum_{n=1}^{\infty} \frac{Z(-1)^n}{n\pi} S_{in}(n\pi t)$$

$$X'' + 6Y = \sum_{n=1}^{\infty} \frac{2(-1)^n}{n\pi} S_{1n}(n\pi t)$$

This is

November 17, 2021 21/27

 $X^{H} + (GYX) = \frac{2}{\pi} S_{In}(\pi t) - \frac{1}{\pi} S_{In}(2\pi t) + \frac{2}{3\pi} S_{In}(3\pi t) + \dots$

- We can use the method of undetermined befficients and look for Xp of the
 - $f_{0}(m) = \sum_{n=1}^{\infty} B_n S_n (n\pi t)$
- Suppose Xp can be differentiated term by term $Xp' = \sum_{n=1}^{\infty} \frac{d}{dt} B_n Sin(n\pi t)$

◆ロ → ◆母 → ◆ 重 → ◆ 重 → ■ November 17, 2021

22/27

$$= \sum_{n=1}^{\infty} B_n(n\pi) C_{s}(n\pi t)$$

$$X \varphi'' = \sum_{n=1}^{\infty} -B_n (n\pi)^2 S_{nn} (n\pi t)$$

$$X_{p} = \sum_{n=1}^{\infty} \mathbb{E}_{n} S_{n}(n\pi t)$$

$$X_{p}'' + 64 X_{p} = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} S_{n}(n\pi t)$$

$$\sum_{n=1}^{\infty} -B_{n}(n\pi)S.n(n\pi t) + \sum_{n=1}^{\infty} GYB_{n}S.n(n\pi t) = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi}S.n(n\pi t)$$

$$\sum_{n=1}^{\infty} \left(-n^{2}\pi^{2} + 6Y \right) B_{n} S_{n} \left(n\pi t \right) = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} S_{n} \left(n\pi t \right)$$

Molehing coefficients

$$(64 - n^2 \pi^2) B_n = \frac{2(-1)^2}{n \pi^2}$$

Since
$$GY - n^{2}\pi^{2} \neq 0$$
 for all n
 $B_{n} = \frac{2(-1)^{n+1}}{n\pi (GY - n^{2}\pi^{2})}$

November 17, 2021 24/27

・ロト・西ト・ヨト・ヨー うへの

