

Sections 16, 17, & 18

Let's solve the following IVP using the Laplace transform.

$$y'' + 6y' + 34y = 0, \quad y(0) = 1, \quad y'(0) = 1$$

Determine $Y(s) = \mathcal{L}\{y(t)\}$ by taking the transform of the ODE, subbing in the IC, and isolating Y .

$$Y(s) = \frac{7 + s}{s^2 + 6s + 34}$$

$$y'' + 6y' + 34y = 0, \quad y(0) = 1, \quad y'(0) = 1$$

Now, do whatever algebra is necessary to write $Y(s)$ in a form from which the inverse transform can be taken.

$$Y(s) = \frac{s+7}{s^2+6s+34} = \frac{s+7}{(s+3)^2+25}$$

$$s^2+6s+34 = (s+3)^2+25$$

$$Y(s) = \frac{s+3}{(s+3)^2+5^2} + \frac{4}{5} \frac{5}{(s+3)^2+5^2}$$

$$y(t) = e^{-3t} \cos(5t) + \frac{4}{5} e^{-3t} \sin(5t)$$

$$y'' + 6y' + 34y = 0, \quad y(0) = 1, \quad y'(0) = 1$$

Evaluate the solution to the IVP $y(t) = \mathcal{L}^{-1}\{Y(s)\}$.

$$Y(s) = \frac{s+3}{(s+3)^2 + 5^2} + \frac{4}{5} \frac{5}{(s+3)^2 + 5^2}$$

$$y(t) = e^{-3t} \cos(5t) + \frac{4}{5} e^{-3t} \sin(5t)$$

$$y'' + 6y' + 34y = 0, \quad y(0) = 1, \quad y'(0) = 1$$

$$y(t) = e^{-3t} \cos(5t) + \frac{4}{5} e^{-3t} \sin(5t).$$

Are there any quick ways to verify that this is probably correct?

- Verify initial conditions
- Take \mathcal{L} to make sure it goes back to previous step.
- cross reference w/ roots of Characteristic polynomial

Example

A 1 kg mass is attached to a spring with spring constant 25 N/m. The mass is at rest at equilibrium. After $t = 2$ seconds, a unit impulse force $f(t) = \delta(t - 2)$ is applied. Assuming that there is no damping, which of the following IVPs governs the displacement of the mass?

(a) $25x'' + x = 0, \quad x(0) = \delta(t-2), \quad x'(0) = 0$

(b) $x'' + 25x = 2, \quad x(0) = 0, \quad x'(0) = 0$

(c) $x'' + 25x = \delta(t-2), \quad x(0) = 0, \quad x'(0) = 0$

(d) $x'' + 2x' + 25x = 0, \quad x(0) = 0, \quad x'(0) = 0$

Example Continued...

Letting $X(s) = \mathcal{L}\{x(t)\}$, take the Laplace transform and isolate $X(s)$.

$$x'' + 25x = \delta(t - 2), \quad x(0) = 0, \quad x'(0) = 0$$

$$\mathcal{L}\{\delta(t - a)\} = e^{-as} \quad \text{for } a \geq 0.$$

$$X(s) = \frac{e^{-2s}}{s^2 + 25}$$

$$\mathcal{L}\{x'' + 25x\} = \mathcal{L}\{\delta(t - 2)\}$$

$$s^2 X(s) - s x(0) - x'(0) + 25X = e^{-2s}$$

Example Continued...

Take the inverse Laplace transform to determine the displacement $x(t) = \mathcal{L}^{-1}\{X(s)\}$.

$$X(s) = \frac{e^{-2s}}{s^2 + 5^2}$$

$$\mathcal{L}^{-1}\{e^{-as}F(s)\} = f(t-a)u(t-a), \quad \mathcal{L}^{-1}\{F(s)\} = f(t)$$

$$F(s) = \frac{1}{s^2 + 5^2} = \frac{1}{5} \frac{5}{s^2 + 5^2} \xrightarrow{\mathcal{L}^{-1}} f(t) = \frac{1}{5} \sin(5t)$$

$$x(t) = \frac{1}{5} \sin(5(t-2))u(t-2)$$

$$x'' + 25x = \delta(t - 2), \quad x(0) = 0, \quad x'(0) = 0$$

$$x(t) = \frac{1}{5} \sin(5(t - 2)) \mathcal{U}(t - 2).$$

Are there any quick ways to verify that this is probably correct?


- Since $\delta(t-2)$ is in the ODE, $\mathcal{U}(t-2)$ should be in the solution.
- $m^2 + 25 \Rightarrow$ solutions would be $\cos(5t)$ and/or $\sin(5t)$

Fourier Series

Consider the function $f(x) = \begin{cases} x+1, & -1 < x < 0 \\ 0, & 0 < x < 1 \end{cases}$.

Identify p , and determine whether f is even, odd, or neither.

$p = 1$ symmetry? no symmetry



$$f(x) = \begin{cases} x + 1, & -1 < x < 0 \\ 0, & 0 < x < 1 \end{cases}$$

$p = 1$ and this function doesn't have symmetry.

Find a_0 .

$$\begin{aligned} a_0 &= \frac{1}{1} \int_{-1}^1 f(x) dx \\ &= \int_{-1}^0 (x+1) dx = \frac{1}{2} \end{aligned}$$

This is as far as we got. The a's and b's are on the following slides.

$$f(x) = \begin{cases} x + 1, & -1 < x < 0 \\ 0, & 0 < x < 1 \end{cases}$$

$$a_0 = \frac{1}{2}$$

Find a_n .

$$f(x) = \begin{cases} x + 1, & -1 < x < 0 \\ 0, & 0 < x < 1 \end{cases}$$

$$a_n = \frac{1 - (-1)^n}{n^2 \pi^2}$$

Find b_n .


$$f(x) = \begin{cases} x + 1, & -1 < x < 0 \\ 0, & 0 < x < 1 \end{cases}$$

$$a_0 = \frac{1}{2}, \quad a_n = \frac{1 - (-1)^n}{n^2 \pi^2}, \quad b_n = -\frac{1}{n\pi}$$

Write the Fourier series for f .

$$f(x) = \begin{cases} x+1, & -1 < x < 0 \\ 0, & 0 < x < 1 \end{cases} \quad \text{Let's plot its Fourier series.}$$

$$f(x) = \frac{1}{4} + \sum_{n=1}^{\infty} \left[\frac{1 - (-1)^n}{n^2 \pi^2} \cos(n\pi x) - \frac{1}{n\pi} \sin(n\pi x) \right]$$



The series would be the pink plot (plus the blue one)