

Section 17: Fourier Series: Trigonometric Series

Suppose f is piecewise continuous on the interval $(-p, p)$. Then we can write f as a Fourier series

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \left(\frac{n\pi x}{p} \right) + b_n \sin \left(\frac{n\pi x}{p} \right) \right)$$

where

$$a_0 = \frac{1}{p} \int_{-p}^p f(x) dx,$$

$$a_n = \frac{1}{p} \int_{-p}^p f(x) \cos \left(\frac{n\pi x}{p} \right) dx, \quad \text{and}$$

$$b_n = \frac{1}{p} \int_{-p}^p f(x) \sin \left(\frac{n\pi x}{p} \right) dx$$

Convergence of the Series

Theorem: If f is continuous at x_0 in $(-p, p)$, then the series converges to $f(x_0)$ at that point. If f has a jump discontinuity at the point x_0 in $(-p, p)$, then the series **converges in the mean** to the average value

$$\frac{f(x_0-) + f(x_0+)}{2} \stackrel{\text{def}}{=} \frac{1}{2} \left(\lim_{x \rightarrow x_0^-} f(x) + \lim_{x \rightarrow x_0^+} f(x) \right)$$

at that point.

Periodic Extension:

The series is also defined for x outside of the original domain $(-p, p)$. The extension to all real numbers is $2p$ -periodic.

Example

$$f(x) = \begin{cases} 1, & -1 < x < 0 \\ -2, & 0 \leq x < 1 \end{cases}, \quad f(x) = -\frac{1}{2} + \sum_{n=1}^{\infty} \frac{3((-1)^n - 1)}{n\pi} \sin(n\pi x).$$




Figure: Plot of the infinite sum, the limit for the Fourier series of f .

Find the Fourier Series for $f(x) = x$, $-1 < x < 1$

$$P=1 \quad \frac{n\pi x}{P} = \frac{n\pi x}{1} = n\pi x$$

$$a_0 = \frac{1}{P} \int_{-P}^P f(x) dx$$

$$= \frac{1}{1} \int_{-1}^1 x dx = \left. \frac{x^2}{2} \right|_{-1}^1 = \frac{1^2}{2} - \frac{(-1)^2}{2} = 0$$

$$a_n = \frac{1}{P} \int_{-P}^P f(x) \cos\left(\frac{n\pi x}{P}\right) dx$$

$$= \frac{1}{1} \int_{-1}^1 x \cos(n\pi x) dx$$

$$u = x, \quad du = dx$$

$$v = \frac{1}{n\pi} \sin(n\pi x) \quad \text{d} \quad \cos(n\pi x) dx$$

$$a_n = \frac{1}{n\pi} \times \sin(n\pi x) \Big|_{-1}^1 - \int_{-1}^1 \frac{1}{n\pi} \sin(n\pi x) dx$$

$$\sin(n\pi) = 0$$

$$= \frac{-1}{n\pi} \left(\frac{-1}{n\pi} \right) \cos(n\pi x) \Big|_{-1}^1$$

$$= \frac{1}{n^2 \pi^2} \cos(n\pi) - \frac{1}{n^2 \pi^2} \cos(-n\pi) = 0$$

$$\cos(-n\pi) = \cos(n\pi)$$

$$a_n = 0 \quad \text{for } n \geq 1$$

$$b_n = \frac{1}{p} \int_{-p}^p f(x) \sin\left(\frac{n\pi x}{p}\right) dx$$

$$= \frac{1}{1} \int_{-1}^1 x \sin(n\pi x) dx$$

$$u = x \quad du = dx$$

$$v = \frac{-1}{n\pi} \cos(n\pi x) \quad dv = \sin(n\pi x) dx$$

$$b_n = \frac{-1}{n\pi} x \cos(n\pi x) \Big|_{-1}^1 - \int_{-1}^1 \frac{-1}{n\pi} \cos(n\pi x) dx$$

$$= \frac{-1}{n\pi} (1) \cos(n\pi) - \frac{-1}{n\pi} (-1) \cos(-n\pi) + \frac{1}{n\pi} \left(\frac{1}{n\pi}\right) \sin(n\pi x) \Big|_{-1}^1$$

$$= \frac{-1}{n\pi} \cos(n\pi) - \frac{1}{n\pi} \cos(n\pi)$$

$$= \frac{-2}{n\pi} (-1)^n = \frac{2}{n\pi} (-1)^{n+1}$$

$$a_0 = 0, \quad a_n = 0, \quad b_n = \frac{2(-1)^{n+1}}{n\pi}$$

$$f(x) = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi x)$$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \left(\frac{n\pi x}{p} \right) + b_n \sin \left(\frac{n\pi x}{p} \right) \right)$$

$$f(x) = x, \quad -1 < x < 1$$

Symmetry

For $f(x) = x$, $-1 < x < 1$

$$f(x) = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi x)$$

Observation: f is an odd function. It is not surprising then that there are no nonzero constant or cosine terms (which have even symmetry) in the Fourier series for f .

The following plots show f , f plotted along with some partial sums of the series, and f along with a partial sum of its series extended outside of the original domain $(-1, 1)$.




Figure: Plot of $f(x) = x$ for $-1 < x < 1$




Figure: Plot of $f(x) = x$ for $-1 < x < 1$ with two terms of the Fourier series.




Figure: Plot of $f(x) = x$ for $-1 < x < 1$ with 10 terms of the Fourier series




Figure: Plot of $f(x) = x$ for $-1 < x < 1$ with the Fourier series plotted on $(-3, 3)$. Note that the series repeats the profile every 2 units. At the jumps, the series converges to $(-1 + 1)/2 = 0$.




Figure: Here is a plot of the series (what it converges to). We see the periodicity and convergence in the mean. **Note:** A plot like this is determined by our knowledge of the generating function and Fourier series, not by analyzing the series itself.

Solution of a Differential Equation

An undamped spring mass system has a mass of 2 kg attached to a spring with spring constant 128 N/m. The mass is driven by an external force $f(t) = 2t$ for $-1 < t < 1$ that is 2-periodic so that $f(t+2) = f(t)$ for all $t > 0$. Determine a particular solution x_p for the displacement for $t > 0$.

From the last example, we can express

f as a series

$$f(t) = 2 \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

The ODE is

$$2x'' + 128x = 2 \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

$$\Rightarrow x'' + 64x = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

We can look for x_p in the form

$$x_p = \sum_{n=1}^{\infty} B_n \sin(n\pi t)$$

We'll assume we can differentiate term by term.

$$x_p' = \sum_{n=1}^{\infty} B_n (n\pi) \cos(n\pi t)$$

$$x_p'' = \sum_{n=1}^{\infty} B_n (-n^2\pi^2) \sin(n\pi t)$$

$$X_p'' + 64 X_p = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

$$\sum_{n=1}^{\infty} -n^2 \pi^2 B_n \sin(n\pi t) + 64 \sum_{n=1}^{\infty} B_n \sin(n\pi t) =$$

$$\sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

Collect like terms

$$\sum_{n=1}^{\infty} (-n^2 \pi^2 B_n + 64 B_n) \sin(n\pi t) = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

$$\sum_{n=1}^{\infty} (64 - n^2 \pi^2) B_n \sin(n\pi t) = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi} \sin(n\pi t)$$

Equating like terms

$$(64 - n^2 \pi^2) B_n = \frac{2(-1)^{n+1}}{n\pi}$$

$$B_n = \frac{2(-1)^{n+1}}{n\pi (64 - n^2 \pi^2)}$$

The particular solution

$$x_p = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n\pi (64 - n^2 \pi^2)} \sin(n\pi t)$$