

Section 13: The Laplace Transform

A quick word about functions of 2-variables:

Suppose $G(s, t)$ is a function of two independent variables (s and t) defined over some rectangle in the plane $a \leq t \leq b$, $c \leq s \leq d$. If we compute an integral with respect to one of these variables, say t ,

$$\int_{\alpha}^{\beta} G(s, t) dt$$

- ▶ the result is a function of the remaining variable s , and
- ▶ the variable s is treated as a constant while integrating with respect to t .

For Example...

Assume that $s \neq 0$ and $b > 0$. Compute the integral

$$\int_0^b e^{-st} dt$$

Treat s like a
constant

$$= \frac{1}{-s} e^{-st} \Big|_0^b$$

$$\int e^{at} dt = \frac{1}{a} e^{at} + C$$

a constant

$$= \frac{1}{-s} e^{-s(b)} - \frac{1}{-s} e^{-s(0)}$$

$$= -\frac{1}{s} e^{-bs} + \frac{1}{s}$$

Integral Transform

An **integral transform** is a mapping that assigns to a function $f(t)$ another function $F(s)$ via an integral of the form

$$\int_a^b K(s, t) f(t) dt.$$

- ▶ The function K is called the **kernel** of the transformation.
- ▶ The limits a and b may be finite or infinite.
- ▶ The integral may be improper so that convergence/divergence must be considered.
- ▶ This transform is **linear** in the sense that

$$\int_a^b K(s, t)(\alpha f(t) + \beta g(t)) dt = \alpha \int_a^b K(s, t) f(t) dt + \beta \int_a^b K(s, t) g(t) dt.$$

The Laplace Transform

Definition: Let $f(t)$ be defined on $[0, \infty)$. The Laplace transform of f is denoted and defined by

$$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt = F(s).$$

The domain of the transformation $F(s)$ is the set of all s such that the integral is convergent.

Note: The **kernel** for the Laplace transform is $K(s, t) = e^{-st}$.

Limits at Infinity e^{-st}


If $s > 0$, evaluate

$$\lim_{t \rightarrow \infty} e^{-st}$$

$$= 0$$

$$s > 0$$

$$\Rightarrow -st < 0$$



If $s < 0$, evaluate

$$\lim_{t \rightarrow \infty} e^{-st}$$

$$= \infty$$

$$\text{If } s < 0, \text{ then } -st > 0$$

Find the Laplace transform of $f(t) = 1$

By definition

$$\mathcal{L}\{1\} = \int_0^{\infty} e^{-st} \cdot 1 \, dt$$

Consider separately $s=0$, $s \neq 0$.

If $s=0$, the integral is

$$\int_0^{\infty} dt = \lim_{b \rightarrow \infty} \int_0^b dt = \lim_{b \rightarrow \infty} t \Big|_0^b = \lim_{b \rightarrow \infty} (b-0) = \infty$$

The integral diverges. So $s=0$ is not in the domain of $\mathcal{L}\{1\}$.

For $s \neq 0$

$$\begin{aligned}\mathcal{L}\{1\} &= \int_0^{\infty} e^{-st} dt = \lim_{b \rightarrow \infty} \int_0^b e^{-st} dt \\ &= \lim_{b \rightarrow \infty} \left(\frac{-1}{s} e^{-bs} + \frac{1}{s} \right)\end{aligned}$$

$$\text{for } s > 0 \quad = \frac{-1}{s} \cdot 0 + \frac{1}{s} = \frac{1}{s}$$

The integral diverges if $s \leq 0$.

$$\mathcal{L}\{1\} = \frac{1}{s} \text{ with domain } s > 0.$$

Find the Laplace transform of $f(t) = t$

By definition

$$\mathcal{L}\{t\} = \int_0^{\infty} e^{-st} t \, dt$$

If $s=0$, the integral diverges. For $s \neq 0$.

$$\int_0^{\infty} e^{-st} t \, dt$$

$$\begin{aligned} u &= t & du &= dt \\ v &= \frac{-1}{s} e^{-st} & dv &= -e^{-st} dt \end{aligned}$$

$$= \left. \frac{-1}{s} t e^{-st} \right|_0^{\infty} - \int_0^{\infty} \frac{-1}{s} e^{-st} \, dt$$

$$= \frac{-1}{s}(0 - 0) + \frac{1}{s} \int_0^{\infty} e^{-st} \, dt$$

* for $s > 0$

$$\lim_{t \rightarrow \infty} t e^{-st} = 0$$

$$\mathcal{L}\{t\} = \frac{1}{s} \int_0^{\infty} \underbrace{e^{-st}}_{\mathcal{L}\{1\}} dt \quad \text{for } s > 0$$


$$= \frac{1}{s} \left(\frac{1}{s} \right) = \frac{1}{s^2}$$

$$\mathcal{L}\{t\} = \frac{1}{s^2} \quad \text{w/ domain } s > 0.$$

A piecewise defined function

Find the Laplace transform of f defined by

$$f(t) = \begin{cases} 2t, & 0 \leq t < 10 \\ 0, & t \geq 10 \end{cases}$$



By definition

$$\begin{aligned} \mathcal{L}\{f(t)\} &= \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^{10} e^{-st} f(t) dt + \int_{10}^{\infty} e^{-st} f(t) dt \\ &= \int_0^{10} e^{-st} (2t) dt + \int_{10}^{\infty} \cancel{e^{-st}} (0) dt \end{aligned}$$

When $s=0$, we have $\int_0^{10} 2t dt = t^2 \Big|_0^{10} = 100$

For $s \neq 0$

$$2 \int_0^{10} e^{-st} t dt$$

$$= 2 \left[-\frac{1}{s} t e^{-st} \right]_0^{10} + \frac{1}{s} \int_0^{10} e^{-st} dt$$

$$= 2 \left[-\frac{1}{s} t e^{-st} \right]_0^{10} - \frac{1}{s^2} e^{-st} \Big|_0^{10}$$

$$= 2 \left[-\frac{10}{s} e^{-10s} - 0 - \frac{1}{s^2} (e^{-10s} - e^0) \right]$$

$$= -\frac{20}{s} e^{-10s} - \frac{2}{s^2} e^{-10s} + \frac{2}{s^2}$$

$$\mathcal{L}\{f(t)\} = \begin{cases} 100, & s=0 \\ \frac{2}{s^2} - \frac{20}{s} e^{-10s} - \frac{2}{s^2} e^{-10s}, & s \neq 0 \end{cases}$$

For $f(t) = \begin{cases} 2t, & 0 \leq t < 10 \\ 0, & 10 \leq t \end{cases}$

The Laplace Transform is a Linear Transformation

Some basic results include:

- ▶ $\mathcal{L}\{\alpha f(t) + \beta g(t)\} = \alpha F(s) + \beta G(s)$
- ▶ $\mathcal{L}\{1\} = \frac{1}{s}, \quad s > 0$
- ▶ $\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}, \quad s > 0 \text{ for } n = 1, 2, \dots$
- ▶ $\mathcal{L}\{e^{at}\} = \frac{1}{s-a}, \quad s > a$
- ▶ $\mathcal{L}\{\cos kt\} = \frac{s}{s^2+k^2}, \quad s > 0$
- ▶ $\mathcal{L}\{\sin kt\} = \frac{k}{s^2+k^2}, \quad s > 0$

Evaluate the Laplace transform $\mathcal{L}\{f(t)\}$ if

(a) $f(t) = \cos(\pi t)$

$$\mathcal{L}\{\cos kt\} = \frac{s}{s^2 + k^2}, \quad s > 0$$

Here, $k = \pi$

$$\mathcal{L}\{\cos(\pi t)\} = \frac{s}{s^2 + \pi^2}, \quad s > 0$$

Evaluate the Laplace transform $\mathcal{L}\{f(t)\}$ if

(b) $f(t) = 2t^4 - e^{-5t} + 3$

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{2t^4 - e^{-5t} + 3\}$$

$$= 2\mathcal{L}\{t^4\} - \mathcal{L}\{e^{-5t}\} + 3\mathcal{L}\{1\}$$

$$= 2\left(\frac{4!}{s^{4+1}}\right) - \frac{1}{s - (-5)} + 3\left(\frac{1}{s}\right)$$

$$= \frac{2(4!)}{s^5} - \frac{1}{s+5} + \frac{3}{s}$$

$$\mathcal{L}\{1\} = \frac{1}{s}, \quad s$$

$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}},$$

$$\mathcal{L}\{e^{at}\} = \frac{1}{s-a},$$