

Section 13: The Laplace Transform

Definition: Let $f(t)$ be defined on $[0, \infty)$. The Laplace transform of f is denoted and defined by

$$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt = F(s).$$

The domain of the transformation $F(s)$ is the set of all s such that the integral is convergent.

The Laplace Transform is a Linear Transformation

Some basic results include:

- ▶ $\mathcal{L}\{\alpha f(t) + \beta g(t)\} = \alpha F(s) + \beta G(s)$
- ▶ $\mathcal{L}\{1\} = \frac{1}{s}, \quad s > 0$
- ▶ $\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}, \quad s > 0 \text{ for } n = 1, 2, \dots$
- ▶ $\mathcal{L}\{e^{at}\} = \frac{1}{s-a}, \quad s > a$
- ▶ $\mathcal{L}\{\cos kt\} = \frac{s}{s^2+k^2}, \quad s > 0$
- ▶ $\mathcal{L}\{\sin kt\} = \frac{k}{s^2+k^2}, \quad s > 0$

Evaluate the Laplace transform $\mathcal{L}\{f(t)\}$ if

(d) $f(t) = \sin^2 5t$

$$\mathcal{L}\{\cos kt\} = \frac{s}{s^2 + k^2},$$

$$\mathcal{L}\{\sin kt\} = \frac{k}{s^2 + k^2},$$

$$f(t) = \frac{1}{2} - \frac{1}{2} \cos(10t)$$

$$\mathcal{L}\{\sin^2(5t)\} = \mathcal{L}\left\{\frac{1}{2} - \frac{1}{2} \cos(10t)\right\}$$

$$= \frac{1}{2} \mathcal{L}\{1\} - \frac{1}{2} \mathcal{L}\{\cos(10t)\}$$

$$= \frac{1}{2} \left(\frac{1}{s}\right) - \frac{1}{2} \frac{s}{s^2 + 10^2} = \frac{1}{2s} - \frac{1}{2} \frac{s}{s^2 + 100}$$

How would we evaluate $\int \sin^2(st) dt$?

wed use $\sin^2 \theta = \frac{1}{2} - \frac{1}{2} \cos(2\theta)$

Sufficient Conditions for Existence of $\mathcal{L}\{f(t)\}$

Definition: Let $c > 0$. A function f defined on $[0, \infty)$ is said to be of *exponential order c* provided there exists positive constants M and T such that $|f(t)| < Me^{ct}$ for all $t > T$.

f grows no faster than an exponential as $t \rightarrow \infty$

Definition: A function f is said to be *piecewise continuous* on an interval $[a, b]$ if f has at most finitely many jump discontinuities on $[a, b]$ and is continuous between each such jump.

Sufficient Conditions for Existence of $\mathcal{L}\{f(t)\}$

Theorem: If f is piecewise continuous on $[0, \infty)$ and of exponential order c for some $c > 0$, then f has a Laplace transform for $s > c$.

An example of a function that is NOT of exponential order for any c is $f(t) = e^{t^2}$. Note that

$$f(t) = e^{t^2} = (e^t)^t \implies |f(t)| > e^{ct} \quad \text{whenever } t > c.$$

This is a function that doesn't have a Laplace transform. We won't be dealing with this type of function here.

Section 14: Inverse Laplace Transforms

Now we wish to go *backwards*: Given $F(s)$ can we find a function $f(t)$ such that $\mathcal{L}\{f(t)\} = F(s)$?

If so, we'll use the following notation

$$\mathcal{L}^{-1}\{F(s)\} = f(t) \quad \text{provided} \quad \mathcal{L}\{f(t)\} = F(s).$$

We'll call $f(t)$ an **inverse Laplace transform** of $F(s)$.

A Table of Inverse Laplace Transforms

- ▶ $\mathcal{L}^{-1} \left\{ \frac{1}{s} \right\} = 1$
- ▶ $\mathcal{L}^{-1} \left\{ \frac{n!}{s^{n+1}} \right\} = t^n$, for $n = 1, 2, \dots$
- ▶ $\mathcal{L}^{-1} \left\{ \frac{1}{s-a} \right\} = e^{at}$
- ▶ $\mathcal{L}^{-1} \left\{ \frac{s}{s^2+k^2} \right\} = \cos kt$
- ▶ $\mathcal{L}^{-1} \left\{ \frac{k}{s^2+k^2} \right\} = \sin kt$

The inverse Laplace transform is also linear so that

$$\mathcal{L}^{-1} \{ \alpha F(s) + \beta G(s) \} = \alpha f(t) + \beta g(t)$$

Using a Table

When using a table of Laplace transforms, the expression must match exactly. For example,

$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}$$

so

$$\mathcal{L}^{-1}\left\{\frac{3!}{s^4}\right\} = t^3.$$

Note that $n = 3$, so there must be $3!$ in the numerator and the power $4 = 3 + 1$ on s .

Find the Inverse Laplace Transform

$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}$$

(a) $\mathcal{L}^{-1}\left\{\frac{1}{s^7}\right\}$ we need $\frac{6!}{s^7}$ to match exactly.

Note: $\frac{1}{s^7} = \frac{1}{s^7} \cdot \frac{6!}{6!} = \frac{1}{6!} \frac{6!}{s^7}$

$$\mathcal{L}^{-1}\left\{\frac{1}{s^7}\right\} = \mathcal{L}^{-1}\left\{\frac{1}{6!} \frac{6!}{s^7}\right\} = \frac{1}{6!} \mathcal{L}^{-1}\left\{\frac{6!}{s^7}\right\} = \frac{1}{6!} t^6$$

Example: Evaluate

$$(b) \quad \mathcal{L}^{-1} \left\{ \frac{s+1}{s^2+9} \right\}$$

$$\mathcal{L}^{-1} \left\{ \frac{s+1}{s^2+9} \right\} = \mathcal{L}^{-1} \left\{ \frac{s}{s^2+9} + \frac{1}{s^2+9} \right\}$$

$$= \mathcal{L}^{-1} \left\{ \frac{s}{s^2+3^2} \right\} + \mathcal{L}^{-1} \left\{ \frac{1}{s^2+3^2} \right\}$$

$$= \mathcal{L}^{-1} \left\{ \frac{s}{s^2+3^2} \right\} + \mathcal{L}^{-1} \left\{ \frac{1}{3} \cdot \frac{3}{s^2+3^2} \right\}$$

$$= \mathcal{L}^{-1} \left\{ \frac{s}{s^2+3^2} \right\} + \frac{1}{3} \mathcal{L}^{-1} \left\{ \frac{3}{s^2+3^2} \right\}$$

$$= \cos(3t) + \frac{1}{3} \sin(3t)$$

$$\mathcal{L}^{-1} \left\{ \frac{s}{s^2+k^2} \right\} = \cos kt$$

$$\mathcal{L}^{-1} \left\{ \frac{k}{s^2+k^2} \right\} = \sin kt$$

Example: Evaluate

(c) $\mathcal{L}^{-1} \left\{ \frac{s-8}{s^2-2s} \right\}$ Use a partial fraction decomp.

$$\frac{s-8}{s^2-2s} = \frac{s-8}{s(s-2)} = \frac{A}{s} + \frac{B}{s-2} \quad \text{Clear fractions}$$

$$s-8 = A(s-2) + Bs$$

$$\text{Set } s=0 \quad 0-8 = A(0-2) + B(0) \Rightarrow -8 = -2A \Rightarrow A=4$$

$$s=2 \quad 2-8 = A(2-2) + B(2) \Rightarrow -6 = 2B \Rightarrow B=-3$$

$$\mathcal{L}^{-1} \left\{ \frac{s-8}{s^2-2s} \right\} = \mathcal{L}^{-1} \left\{ \frac{4}{s} - \frac{3}{s-2} \right\}$$

$$= 4 \mathcal{L}^{-1} \left\{ \frac{1}{s} \right\} - 3 \mathcal{L}^{-1} \left\{ \frac{1}{s-2} \right\}$$

$$= 4(1) - 3e^{2t}$$

$$= 4 - 3e^{2t}$$