

Section 15: Shift Theorems

Theorem: (translation in s) Suppose $\mathcal{L}\{f(t)\} = F(s)$. Then for any real number a

$$\mathcal{L}\{e^{at}f(t)\} = F(s - a).$$

In other words, if $F(s)$ has an inverse Laplace transform, then

$$\mathcal{L}^{-1}\{F(s - a)\} = e^{at}\mathcal{L}^{-1}\{F(s)\}.$$

Example: Evaluate

$$\mathcal{L} \left\{ e^{4t} \cos(\pi t) \sin(\pi t) \right\}$$

We need $\mathcal{L} \{ \cos(\pi t) \sin(\pi t) \} = F(s)$

our transform will be $F(s-4)$

Recall $\sin 2\theta = 2 \sin \theta \cos \theta$

so $\cos(\pi t) \sin(\pi t) = \frac{1}{2} \sin(2\pi t)$

$$\mathcal{L} \{ \sin kt \} = \frac{k}{s^2 + k^2}$$

$$F(s) = \mathcal{L} \{ \cos(\pi t) \sin(\pi t) \} = \frac{1}{2} \mathcal{L} \{ \sin(2\pi t) \} = \frac{1}{2} \frac{2\pi}{s^2 + (2\pi)^2}$$

Hence

$$\mathcal{L}\{e^{4t} \cos(\pi t) \sin(\pi t)\} = \frac{1}{2} \frac{2\pi}{(s-4)^2 + (2\pi)^2}$$
$$= \frac{\pi}{(s-4)^2 + 4\pi^2}$$

Evaluate

$$\mathcal{L}^{-1} \left\{ \frac{s}{(s+4)^4} \right\}$$

We need to decompose $\frac{s}{(s+4)^4}$.

The correct form is

$$\frac{s}{(s+4)^4} = \frac{A}{s+4} + \frac{B}{(s+4)^2} + \frac{C}{(s+4)^3} + \frac{D}{(s+4)^4}$$

We can use a shortcut

$$\begin{aligned} \frac{s}{(s+4)^4} &= \frac{s+4-4}{(s+4)^4} = \frac{s+4}{(s+4)^4} - \frac{4}{(s+4)^4} \\ &= \frac{1}{(s+4)^3} - \frac{4}{(s+4)^4} \end{aligned}$$

Hence

$$\mathcal{L}^{-1}\left\{\frac{s}{(s+4)^4}\right\} = \mathcal{L}^{-1}\left\{\frac{1}{(s+4)^3}\right\} - 4 \mathcal{L}^{-1}\left\{\frac{1}{(s+4)^4}\right\}$$

we need $\mathcal{L}^{-1}\left\{\frac{1}{s^3}\right\} = \mathcal{L}^{-1}\left\{\frac{1}{2!} \frac{2!}{s^3}\right\} = \frac{1}{2!} t^2$

$$\mathcal{L}^{-1}\left\{\frac{1}{s^4}\right\} = \mathcal{L}^{-1}\left\{\frac{1}{3!} \frac{3!}{s^4}\right\} = \frac{1}{3!} t^3$$

If $s-a = s+4$, then $a = -4$

$$\begin{aligned}\mathcal{L}^{-1}\left\{\frac{s}{(s+4)^4}\right\} &= \mathcal{L}^{-1}\left\{\frac{1}{(s+4)^3}\right\} - 4 \mathcal{L}^{-1}\left\{\frac{1}{(s+4)^4}\right\} \\ &= \frac{1}{2!} t^2 e^{-4t} - \frac{4}{3!} t^3 e^{-4t}\end{aligned}$$

The Unit Step Function

Let $a \geq 0$. The unit step function $\mathcal{U}(t - a)$ is defined by

$$\mathcal{U}(t - a) = \begin{cases} 0, & 0 \leq t < a \\ 1, & t \geq a \end{cases}$$




Figure: We can use the unit step function to provide convenient expressions for piecewise defined functions.

Unit Step Function Notation

The unit step function is sometimes referred to as the *Heaviside step function*¹. However, many reserve that name for the version of this function defined on the interval $(-\infty, \infty)$.

- ▶ An alternative notations include

$$\mathcal{U}(t - a), \quad u_a(t), \quad u(t - a), \quad \text{and} \quad H(t - a).$$

- ▶ Restricting our focus to functions defined on $[0, \infty)$, $f(t) = 1$ and $f(t) = \mathcal{U}(t)$ are indistinguishable.

¹Named after English mathematician Oliver Heaviside.

Piecewise Defined Functions

Verify that

$a \geq 0$

$$f(t) = \begin{cases} g(t), & 0 \leq t < a \\ h(t), & t \geq a \end{cases} = g(t) - g(t)\mathcal{U}(t-a) + h(t)\mathcal{U}(t-a)$$

We'll show these are equal for all $t \geq 0$.

We'll have to consider $0 \leq t < a$ and $t \geq a$.

Suppose $0 \leq t < a$. Then $\mathcal{U}(t-a) = 0$

$$\begin{aligned} g(t) - g(t)\mathcal{U}(t-a) + h(t)\mathcal{U}(t-a) &= g(t) - g(t) \cdot 0 + h(t) \cdot 0 \\ &= g(t) \quad \text{as required} \end{aligned}$$

$$f(t) = \begin{cases} g(t), & 0 \leq t < a \\ h(t), & t \geq a \end{cases} = g(t) - g(t)\mathcal{U}(t-a) + h(t)\mathcal{U}(t-a)$$

Now, suppose $t \geq a$. Then

$$\mathcal{U}(t-a) = 1$$

$$g(t) - g(t)\mathcal{U}(t-a) + h(t)\mathcal{U}(t-a) =$$

$$g(t) - g(t) \cdot 1 + h(t) \cdot 1 = h(t)$$

as required.

Piecewise Defined Functions in Terms of \mathcal{U}

Write f on one line in terms of \mathcal{U} as needed

$$f(t) = \begin{cases} e^t, & 0 \leq t < 2 \\ t^2, & 2 \leq t < 5 \\ 2t & t \geq 5 \end{cases}$$

We can use $\mathcal{U}(t-a)$ as a switch to turn pieces of the function on and off along with adding them in or subtracting them off.

$$f(t) = e^t - e^t \mathcal{U}(t-2) + t^2 \mathcal{U}(t-2) - t^2 \mathcal{U}(t-5) + 2t \mathcal{U}(t-5)$$

↑
on ↑
~~off~~ ↑
on ↑
~~off~~ ↑
on

$$f(t) = e^t (u(t-0) - u(t-2)) + e^2 (u(t-2) - u(t-5)) + 2e (u(t-5) - 0)$$

Translation in t

Given a function $f(t)$ for $t \geq 0$, and a number $a > 0$

$$f(t-a)\mathcal{U}(t-a) = \begin{cases} 0, & 0 \leq t < a \\ f(t-a), & t \geq a \end{cases}.$$




Figure: The function $f(t-a)\mathcal{U}(t-a)$ has the graph of f shifted a units to the right with value of zero for t to the left of a .

Theorem (translation in t)

If $F(s) = \mathcal{L}\{f(t)\}$ and $a > 0$, then

$$\mathcal{L}\{f(t-a)\mathcal{U}(t-a)\} = e^{-as}F(s).$$

In particular,

$$\mathcal{L}\{\mathcal{U}(t-a)\} = \frac{e^{-as}}{s}.$$

← here
 $f(t) = 1$
 $\mathcal{L}\{f\} = \frac{1}{s}$

As another example,

$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}} \implies \mathcal{L}\{(t-a)^n\mathcal{U}(t-a)\} = \frac{n!e^{-as}}{s^{n+1}}.$$